

DRIVERS OF CHANGE

Coordinating Lead Authors:

Semia Cherif (*Tunisia*), Enrique Doblas-Miranda (*Spain*), Piero Lionello (*Italy*)

Lead Authors:

Carlos Borrego (*Portugal*), Filippo Giorgi (*Italy*), Ana Iglesias (*Spain*), Sihem Jebari (*Tunisia*), Ezzeddine Mahmoudi (*Tunisia*), Marco Moriondo (*Italy*), Olivier Pringault (*France*), Gil Rilov (*Israel*), Samuel Somot (*France*), Athanassios Tsikliras (*Greece*), Montserrat Vila (*Spain*), George Zittis (*Cyprus*)

Contributing Authors:

Giovanni Argenti (*Italy*), Marie-Anne Auger-Rozenberg (*France*), Ernesto Azzurro (*Italy*), Corina Basnou (*Spain*), Sophie Bastin (*France*), Mustapha Béjaoui (*Tunisia*), Lorenzo Brilli (*Italy*), Martina Carrete (*Spain*), Emma Cebrian (*Spain*), Hanene Chaabane (*Tunisia*), Sílvia Coelho (*Portugal*), Renato Colucci (*Italy*), Styliani Dafka (*Germany*), Sofia Darmaraki (*Greece/Canada*), Camilla Dibari (*Italy*), Donna Dimarchopoulou (*Greece*), Jean-Claude Dutay (*France*), Monia El Bour (*Tunisia*), Antonietta Elia (*Spain*), Elena Georgopoulou (*Greece*), Sylvaine Giakoumi (*France*), Juan Jesús González Alemán (*Spain*), Pablo González-Moreno (*UK*), Madeleine Goutx (*France*), Olivier Grünberger (*Tunisia*), Ivan Güttler (*Croatia*), Nathalie Hilmi (*Monaco*), Gabriel Jordà (*Spain*), Stelios Katsanevakis (*Greece*), Mehdi Lahlou (*Morocco*), Manfred A. Lange (*Cyprus*), Luisa Leolini (*Italy*), Myriam Lopes (*Portugal*), Annarita Mariotti (*USA*), Ana Isabel Miranda (*Portugal*), Meryem Mojtabid (*France*), Alexandra Monteiro (*Portugal*), Samuel Morin (*France*), Pierre Nabat (*France*), Anika Obermann-Hellhund (*Germany*), Tuğba Öztürk (*Turkey*), Androniki Pardalou (*Greece*), Sandra Rafael (*Portugal*), Francesca Raffaele (*Italy*), Lena Reimann (*Germany*), Alain Roques (*France*), Asma Sakka Hlaili (*Tunisia*), Alberto Santini (*Italy*), Giuseppe Scarcella (*Italy*), Katrin Schroeder (*Italy*), Isla Simpson (*USA*), Nicolina Staglioniò (*Italy*), Meryem Tanharte (*Morocco*), Rob Tanner (*France*), Rémi Thiéblemont (*France*), Yves Tramblay (*France*), Marco Turco (*Italy*), Nassos Vafeidis (*Germany*), Martin Wild (*Switzerland*), Elena Xoplaki (*Germany*), Argyro Zenetos (*Greece*)

This chapter should be cited as: Cherif S, Doblas-Miranda E, Lionello P, Borrego C, Giorgi F, Iglesias A, Jebari S, Mahmoudi E, Moriondo M, Pringault O, Rilov G, Somot S, Tsikliras A, Vila M, Zittis G 2020 Drivers of change. In: Climate and Environmental Change in the Mediterranean Basin – Current Situation and Risks for the Future. First Mediterranean Assessment Report [Cramer W, Guiot J, Marin K (eds.)] Union for the Mediterranean, Plan Bleu, UNEP/MAP, Marseille, France, pp. 59–180.



Table of contents

2. Drivers of change	64
Executive summary.....	64
2.1 Introduction	66
2.2 Climate change	67
2.2.1 Framing	67
2.2.1.1 <i>Observations and reanalyses</i>	68
2.2.1.2 <i>Modelling</i>	68
2.2.2 General circulation and dynamics	70
2.2.2.1 <i>General circulation</i>	70
<i>Observed trends</i>	70
<i>Future changes</i>	70
2.2.2.2 <i>Teleconnection patterns</i>	72
2.2.2.3 <i>Extratropical cyclones and medicanes</i>	72
2.2.2.4 <i>Regional winds</i>	73
2.2.3 Radiation, clouds and aerosols	74
2.2.3.1 <i>Observed change in surface radiation</i>	74
2.2.3.2 <i>Projected change in surface radiation</i>	75
2.2.4 Temperature and related extremes	75
2.2.4.1 <i>Observed temperature changes</i>	75
2.2.4.2 <i>Future temperatures</i>	76
2.2.5 Precipitation, related extremes and the water cycle	79
2.2.5.1 <i>Observed trends in precipitation</i>	79
2.2.5.2 <i>Future precipitation</i>	81
<i>Mean precipitation changes</i>	81
<i>Variability and extremes</i>	82
2.2.5.3 <i>Changes in evaporation, net water losses over sea and over land</i>	83
2.2.6 The cryosphere	84
2.2.6.1 <i>Observed trends in the cryosphere</i>	84
2.2.6.2 <i>Future conditions in the cryosphere</i>	85
2.2.7 Ocean hydrology	86
2.2.7.1 <i>Observed change in marine waters</i>	87
2.2.7.2 <i>Future change in marine waters</i>	88
<i>Air-sea and land-sea exchanges</i>	88
<i>Sea surface temperature</i>	89
<i>Sea surface salinity</i>	90
<i>Surface circulation and exchanges across straits</i>	90
<i>Deep water characteristics</i>	90
2.2.8 Sea level, storm surges and wave heights	91
2.2.8.1 <i>Observed change in sea-level</i>	91
2.2.8.2 <i>Future sea-levels</i>	91
2.2.9 Acidification of the Mediterranean Sea	92
2.2.9.1 <i>Observed change in acidity</i>	94
2.2.9.2 <i>Future change in acidity</i>	94
2.3 Pollution.....	94
2.3.1 Introduction	94

2.3.2 Physical pollutants.....	94
2.3.2.1 Particulate matter (PM) levels and sources	95
2.3.2.2 Particulate matter (PM) chemical profiles	96
2.3.2.3 Plastics (macro/micro/nano)	97
2.3.3 Chemical pollutants.....	98
2.3.3.1 Nutrients	98
2.3.3.2 Gaseous pollutants: nitrogen dioxide, sulphur dioxide, ozone.....	100
2.3.3.3 Trace metallic elements	101
2.3.3.4 Organic pollutants	103
Polycyclic Aromatic Hydrocarbons (PAHs)	103
Pesticides	104
2.3.3.5 Emerging contaminants	105
2.3.4 Biological pollutants.....	107
2.4 Land and sea use changes.....	109
2.4.1 Land use changes.....	109
2.4.1.1 Past trends and recent dynamics	109
Mediterranean landscapes	109
Recent changes in Mediterranean landscapes.....	110
2.4.1.2 Principal impacts of land use changes	110
Forestry and other natural resources	110
Crops and livestock.....	112
Different land uses as a single driver of change	113
2.4.1.3 Future projections.....	113
2.4.2 Sea use changes.....	114
2.4.2.1 Trends in fisheries exploitation	114
2.4.2.2 Current status of marine fisheries resources	115
2.4.2.3 The future of marine resources	115
2.5 Non-indigenous species.....	116
2.5.1 Non-indigenous species in the Mediterranean Sea.....	116
2.5.1.1 Spatiotemporal trends, sources and vectors of introduction.....	117
2.5.1.2 Non-indigenous species as drivers of biodiversity and ecosystem change	117
2.5.1.3 Further introductions, monitoring and managing non-indigenous species	119
2.5.2 Terrestrial non-indigenous species and pests	120
2.5.2.1 Spatial patterns and temporal trends	120
Degree of introduction across Mediterranean-type ecosystems and geographical areas	120
Temporal trends of non-indigenous species and pests	122
Pathways of introduction (intentional and accidental) of non-indigenous species and pests	123
2.5.2.2 Non-indigenous species as drivers of biodiversity and ecosystem change	123
2.5.2.3 Further introductions, spread and impacts of non-indigenous species and pests	125
Existing tools for predicting the risk of introduction and research needs	125
Non-indigenous species likely to be introduced into the Mediterranean in the next 20-50 years.....	125
2.6 Interaction among drivers	126
2.6.1 Drivers impacting other drivers	126
2.6.2 Pairs of interacting drivers.....	127
2.6.2.1 Climate change effects on pollution	127
2.6.2.2 Pollution effects on climate change.....	127
2.6.2.3 Impact of climate on land and sea use	127
Effects of climate change on land use	127
Climate change and variability drives dynamics of marine species	128
2.6.2.4 Effects of land use on climate change	129

2.6.2.5	<i>Links between trends in non-indigenous species and climate change</i>	129
	<i>Impact of climate change on marine non-indigenous species.....</i>	129
	<i>Impacts of climate change on terrestrial non-indigenous species.....</i>	131
2.6.2.6	<i>Impacts of pollution on land and sea use</i>	131
2.6.2.7	<i>Impacts of land and sea use change on pollution.....</i>	131
2.6.2.8	<i>Pollution effects on non-indigenous species.....</i>	132
2.6.2.9	<i>Effects of land and water use on non-indigenous species.....</i>	132
2.6.2.10	<i>Effects of non-indigenous species on land and sea use</i>	132
2.6.3	More complex interactions among drivers.....	132
2.6.3.1	<i>Floods</i>	132
2.6.3.2	<i>Desertification.....</i>	132
2.6.3.3	<i>Wildfires.....</i>	132
2.7	Mediterranean socioeconomic scenarios	134
Box 2.1	How much has the Mediterranean Basin warmed since the pre-industrial period?	137
Box 2.2	Representative Concentration Pathways (RCPs).....	138
References.....		139
Information about authors.....		179

2 Drivers of change

Executive summary

Climate drivers

During recent decades, observations of several variables provide evidence of the ongoing anthropogenic climate change in the Mediterranean region, particularly increase of mean and extreme temperatures, and dry environmental conditions. Climate projections show that the region will among the most affected regions by climate change, specifically regarding precipitation and the hydrological cycle, but also mean warming and heat extremes (in both the terrestrial and marine environment), sea level rise and sea water acidification.

Basin-wide, annual mean temperatures are now 1.5°C above the preindustrial level. In the last decades dry conditions have become more frequent and a large reduction of glaciers across high mountains of the Mediterranean has occurred at a progressively increasing pace. Mediterranean Sea waters have become warmer and saltier, Mediterranean sea level has risen at a rate (1.4 mm yr⁻¹) similar to the global trend at centennial scale.

In the future, the regional average warming will exceed the global mean value by 20% and it might reach 5.6°C at the end of the 21st century in the RCP8.5 high emission scenario. Heat waves and warm temperature extremes will intensify. Total annual precipitation is expected to decrease over most of the region (the average reduction rate is approximately 4% per each degree of global warming). However, magnitude and spatial distribution of changes are uncertain, because of differences among models. Dry conditions will be further enhanced by increasing evapotranspiration over land. At the same time, the inter-annual variability of the hydrological cycle will increase, with longer dry spells especially in the southern areas. Extreme precipitation events will become more intense over large parts of the northern Mediterranean areas.

Mediterranean mean sea level is projected to be at the end of the 21st century in the range from 20 to 110 cm higher than at the end of the 20th century, depending on the level of anthropogenic emissions. Sub-regional and local relative sea level rise will be further modulated by vertical land motions and regional circulation features (with deviations in the

order of 10 cm from the basin average). Therefore, though in the future milder marine storms are expected, coastal hazards, floods and erosion will increase, because of mean sea level rise.

Widespread seawater warming will continue. Annual mean surface temperature will increase 2.7-3.8°C and 1.1-2.1°C in one century under the RCP8.5 and the RCP4.5 scenarios, respectively. Marine heat waves will become longer, more intense than today and their spatial extent will increase. Seawater acidification will continue, with a pH reduction that might larger than 0.4 units at the end of the 21st century.

Pollution

Across the Mediterranean Basin, ocean and inland pollution are ubiquitous, diverse and increasing in both quantity and in the number of pollutants, due to demographic pressure, enhanced industrial and agricultural activities, and climate change.

Mediterranean seawater is generally oligotrophic (low nutrient), with decreasing levels from Gibraltar eastwards to the Levantine Sea. Several coastal regions are hotspots of human-induced nutrient inputs. This nutrient enrichment causes eutrophication and may provoke harmful and toxic algal blooms, whose frequency will likely increase. Harmful algal blooms may cause negative impacts on ecosystems and may represent serious economic threats for fisheries, aquaculture, tourism and human health.

Emerging contaminants are well present across the Mediterranean Basin, and enhanced by increasing inflow of untreated wastewater. These substances may cause disorders of the nervous, hormonal and reproductive system. And the increasing frequency of extreme precipitation events in the north of the Mediterranean increases the supply of fecal bacteria and viruses to the coastal zone. The Mediterranean Sea is one of the most polluted large water bodies globally in terms of plastic and the level of this pollution is expected to increase in the future.

The Mediterranean Basin is among the regions in the world with the highest concentrations of gaseous air pollutants (NO₂, SO₂ and O₃). Its dry and sunny climate, and specific atmospheric circulation patterns enhance air pollution levels. Ships are among the major causes of increasing SO₂ and NO_x emissions in this region.

Emissions of aerosols and particulate matter (PM) into the atmosphere arise from a variety of anthropogenic activities. Particular meteorological conditions and natural sources, including the proximity of the Sahara Desert, create particular patterns of aerosol concentrations that may influence particulate matter PM concentrations. The occurrence of critically high PM concentrations associated with dust outbreaks is higher in the southern Mediterranean (>30% of the annual days) than in the northern area (<20% of the annual days).

Land and sea use change

Landscapes and their use have changed over millennia in the Mediterranean Basin, however the rate of change has increased substantially since the second half of the 20th century, with rapid growth of urban and peri-urban areas leading to loss of biodiversity and habitats. Abandonment of agro-pastoralism (which will likely continue in the future) is causing unmanaged shrubs forest development in marginal lands, arid and mountain areas in European countries, while land overexploitation is causing widespread forest degradation in areas of North Africa and the Middle East. Future land use trends depend strongly on regional policies for urbanization, agriculture, forestry and nature conservation.

Marine resource overexploitation and unsustainable fishing practices have increased in time and are the main drivers of the population decline of several species. Presently, more than 60% of marine stocks have collapsed or are overexploited. Sustainable management of marine resources requires the reduction of fishing pressure.

Non-indigenous species

The Mediterranean Sea (and particularly the Levantine Basin) is a hotspot for the establishment of many non-indigenous species (invertebrates, primary producers, and vertebrates), whose arrival and increase are linked to the decrease or collapse in populations of native species. Most marine non-indigenous species enter the Mediterranean Sea from the Red Sea and Atlantic Ocean, but those introduced by ships and aquaculture produce the largest impact on the environment. The number and spread of non-indigenous species will likely further increase with increasing shipping activity and impacts of climate change on the Mediterranean water masses.

Mediterranean land areas currently host a high number of non-indigenous species (mostly plants and invertebrates) in human-modified ecosystems and in regions with high infrastructure development. Most invertebrate species are phytophagous pests that cause damages to crops and forests. Future warming is predicted to induce a northward shift at a speed of $37\text{--}55 \text{ km decade}^{-1}$ of current major non-indigenous species and determine a window of opportunity for new non-indigenous species adapted to dry environments. The presently increasing trend of the numbers of introduced invertebrates and vertebrates (the latter generally caused by accidental escapes) will very likely continue, as they can be easily transported also as stowaways in air and maritime cargo.

Interaction among drivers

When ecosystems and societal sectors are threatened by multiple, co-occurring drivers, climate change, pollution, land and sea use change, and non-indigenous species can interact. Interactions cause effects that can be additive/cumulative, synergistic or antagonistic and result in alteration, intensification, and even in generation of new impacts. Examples of new threats are increase of flood events, due to a combination of climatic and land use changes, desertification, which is the result of increasing aridity and exploitation of resources, and wildfires, affected by forest encroachment and heat waves, among many other interactions.

2.1 Introduction

This chapter describes characteristics and evolution of human-induced and natural factors that cause changes in the Mediterranean Basin ecosystems and human systems. In order to cover most major risks for people and biodiversity, four broad domains of change drivers are considered: climate change and variability, pollution, land and sea use changes and non-indigenous species. These factors correspond to the concept of “direct drivers”, which was introduced in the Millennium Ecosystem Assessment (MEA 2005; Nelson et al. 2006), that unequivocally influence processes in ecosystems and can be identified and measured to differing degrees of accuracy.

Anthropogenic climate change is already affecting the environment and societies in the Mediterranean region. Warming is unequivocal, and there are emerging signs of changes of the hydrological cycle and other climate variables (*Section 2.2*). Climate models indicate a trend towards a warmer and drier environment, seawater warming, with more intense warm extremes both over land and in the sea, and regional increase of sea level (*Section 2.2*). The Mediterranean region is likely very vulnerable to climate change and many components of its terrestrial and marine environment are already under stress (*Section 2.2*).

Atmospheric and water pollution can be driven by many factors, which affect all the compartments of the environment: water, air and soil/sediments. Pollutants can migrate from one media to another. There is a wide range of pollutants that can be biological (e.g., bacteria or insects), chemical (e.g., pesticides, trace metals) or physical (e.g., particulate matter) (*Section 2.4-6* and *Chapters 3 and 4* of this report).

Changes in land and sea use changes are considered among the major direct drivers of environmental change worldwide, but their characteristics vary, depending on each region, even at very local scale. Mediterranean terrestrial landscapes and ecosystems show different patterns of change on northern and southern shores, due to urbanization, coastal development, evolving agricultural and farming practices, including their abandonment. The overexploitation of the Mediterranean Sea resources poses a particular threat due to its intrinsic geographical limits (*Section 2.4*).

Non-indigenous species are profoundly affecting terrestrial and marine ecosystems in the

Mediterranean and their impact is not only measurable in biodiversity alterations, but also in human health and economic damages (*Section 2.5*).

There is no strict consensus of the grouping of the drivers into the categories that have been adopted in this report. More condensed or more articulated lists can be found in the scientific literature. Our four categories include all physical, chemical and biological factors that directly act on the Mediterranean environment, with a substantial correspondence with those used in the recent IPBES 2018 regional reports (Bustamante et al. 2018; Elbakidze et al. 2018; Nyingi et al. 2018; Wu et al. 2018b).

The level or rate of change of direct drivers can be influenced or altered by indirect drivers (MEA 2005; Nelson et al. 2006). Indirect drivers are grouped in categories such as demographic, economic, sociopolitical, cultural, religious, technological, legislation and financial drivers. In turn, indirect drivers are distinguished in “exogenous” and “endogenous” drivers, whose magnitude can and cannot be influenced/alterred by the decision-makers, respectively. Whether a driver is exogenous or endogenous depends on the organizational level and on the spatial and temporal scale. The concepts of indirect and direct drivers roughly match those of driving forces and pressures in the Drivers–Pressures–State–Impacts–Responses (DPSIR) framework, which was initially developed by the European Environmental Agency (EEA 1999).

Understanding of indirect drivers is essential for the benefit of the environment. It is the action on indirect drivers by policymakers and stakeholders that can effectively manage the risks posed to the environment and human societies by climate change, pollution, land and sea use changes and arrival of non-indigenous species. However, this is a quite different topic with respect to the content of this chapter. Mechanism and tools influencing direct drivers are considered in other parts of this report. The scope of this chapter is the assessment of the state of knowledge of physical, chemical, and biological factors, of their present status, past and future evolution in the Mediterranean Basin.

2.2 Climate change

2.2.1 Framing

There is no universal definition of the land boundaries of the Mediterranean region (*Chapter 1*). In this section, and also for much of this report, we adopt a simple regular latitude-longitude box (29°N to 47.5°N and 10°W to 39°E, *Fig. 2.1*), which includes some regions with other than Mediterranean climates, such as in the Alps, the Eastern Balkans or part of the Sahara. This definition of Mediterranean region is similar to the MED zone adopted in IPCC-AR4 (IPCC 2007), and slightly larger than in IPCC-AR5 (IPCC 2013a).

delimitations (*Fig. 2.1*), the Western Mediterranean Sea (MEDW) between the Gibraltar Strait and the Sicily Strait, the Adriatic Sea (ADR) north of the Otranto Strait, the Aegean Sea (AEG) north of the Cretan Arc Straits and the Eastern Mediterranean Sea (MEDE) for the remaining.

When assessing future climate change, it is important to specify the reference period to which climate projections are compared, along with future “time slices”. In MAR1 we use 20-year long time periods following standard IPCC practice. This length of time period is sufficient to smooth part




Figure 2.1 | Mediterranean coastline, topography over land and bathymetry over the sea plus the box definition. Relief data are derived from the ETOP01 1 Arc-Minute Global Relief Model⁵. Sub-regions are defined as West Maghreb (WM), East Maghreb (EM), Levant (LE), Anatolia (AN), Central Mediterranean (CM), Iberia (IB), France (FR), Alps (AL) and Balkans (BA). The Mediterranean Sea is divided into 4 sub-basins, the Western Mediterranean Sea (MEDW), the Eastern Mediterranean Sea (MEDE), the Aegean Sea (AEG) and the Adriatic Sea (ADR).

In order to provide a spatially refined assessment, we define sub-regions over land and over sea by using smaller rectangular boxes. Over land, the European sub-regions follow the definition proposed during the PRUDENCE European project (Christensen et al. 2002) (*Fig. 2.1*), and we define new sub-regions (inspired by Nabat et al. 2015b) for the Middle East and Africa. Over the sea, 4 sub-regions are defined using the natural strait

of the high-frequency natural climate variability that can otherwise mask the forced trend, but it is short enough to assume that climate does not change much during the 20 years covered. For the reference period, we choose the latest years of the 20th century (1980-1999). This choice is a compromise related to the observation and model data availability at the Mediterranean scale. IPCC has traditionally chosen the pre-industrial

⁵ <http://www.ngdc.noaa.gov/mgg/global/global.html>

period (around 1850 or 1900) as a reference but regional climate models and regional high-quality observations are not available for that period. This choice (2 full decades at the end of the 20th century) also targets to facilitate the repeatability of the MAR1 computations made in future Mediterranean studies or reports.

In addition to the reference period, we define also the “present-climate” period (1995–2014), which defines the current climate conditions. The reference period is fixed in different reports to be able to intercompare results of simulations across different model generations, whereas the present-climate will move from one report to another. For example, the IPCC-AR4 (IPCC 2007) defined 1981–2000 as present climate, whereas the IPCC-AR5 (IPCC 2013a) used 1986–2005. For the assessment of past changes, the longest period available (generally 1950–2019 or 1900–2019, if possible) in the observations is used in this report and units such as °C per decade are used in order to compare past trends computed over different past periods.

For the future, we keep 20-year time slices in order to sample the same level of internal variability as in the reference period. We divide the 21st century in 20-year time slices with a present-climate period (2000–2019), a near-future period (2020–2039), a mid-term period centered in 2050 (2040–2059) and a far-future period close to the end of the 21st century (2080–2099). The mid-21st century period is arguably of particular interest for many stakeholders, especially for mid-term adaptation. The end of the 21st century period is also of interest for stakeholders working on mitigation targets and involved in very long-term planning (e.g., for the design and planning of dams, forests, cities).

For the future climate change assessment, an important part of the uncertainty is related to the future evolution of socio-economic development. To be able to propose future climate projections considering various possible socio-economic trajectories and climate policy pathways, we follow the IPCC scenario approach. Where more recent studies are not available, the assessment also considers studies based on the older IPCC SRES approach (Nakićenović 2000). Although results based on multiple IPCC scenarios are reported in the MAR1, we mostly focus on two options which encompass the range of IPCC-AR5, CMIP5 and CORDEX simulations: the “business as usual” scenario (RCP8.5, for an explanation of the RCPs see *Box 2.2*) and the optimistic scenario closest to the UNFCCC Paris Agreement target (RCP2.6). These scenarios have been chosen also due to

model projection availability constraints at the regional scale.

Detecting trends and attributing regional climate change to human influence is challenging due to natural climate variability and the strong spatio-temporal dependency of different climate variables. For projections, uncertainty estimates are provided where this is possible. For small model ensembles, the total range is also given. For larger ensemble, 90% confidence intervals are used as much as possible or else “whisker plots” describing the various statistics of the distribution (median, 25th and 75th percentile, 90% interval, minimum and maximum values).

2.2.1.1 Observations and reanalyses

More and more observation datasets have become available at regional scales, either from satellite or *in-situ* observations, or from reconstructions and reanalyses. This new generation of observation-based products are (1) long and homogeneous enough to allow trend studies (ESA-CCI) (Ribes et al. 2019); (2) of sufficiently high spatial resolution to capture complex topography and land-sea mask (SAFRAN, EURO-4M), thereby allowing regional to local studies; and (3) of sufficiently high temporal resolution (daily or hourly, COMEPHORE) (Fumière et al. 2019), to allow the study of regionally-relevant extreme events. High-resolution gridded products have also become available for southern Mediterranean countries e.g., Cyprus (Camera et al. 2017) or Tunisia (Tramblay et al. 2019).

Finding the best fit-for-purpose observation dataset is becoming a new challenge, given the large number of available products, often characterized by substantial differences. Results of past trend studies and model evaluations are sensitive to the choice of the reference dataset (Flaounas et al. 2012; Prein and Gobiet 2017; Zittis 2018; Fumière et al. 2019; Kotlarski et al. 2019; Peña-Angulo et al. 2020). Long-term, accessible, gridded, well-calibrated and homogeneous in time and space *in-situ* data are nonetheless still lacking, especially for the ocean or the high-frequency variables over land. In addition, regional model-based reanalyses are still rare.

Various observation datasets are used to assess the past evolution of the different components of the Mediterranean climate system. Atmospheric dynamics are mostly assessed against atmospheric reanalyses (ERA-Interim, ERA20C, 20CR) (Section 2.2.2). For aerosols, clouds and surface radiation, both satellite products and station data (BRSN,

GEBA) are used to estimate past evolutions (*Section 2.2.3*). The trend evaluation over land relies on high-resolution observation-based gridded products, i.e., CRU, E-OBS for temperature (*Section 2.2.4*) and CRU, E-OBS, U. Del, GPCC for precipitation (*Section 2.2.5*). The other water cycle components are evaluated against reconstructed products blending *in-situ* observations, satellite and models of river networks (Ludwig et al. 2009; Pellet et al. 2019; Wang and Polcher 2019) (*Section 2.2.6*), while satellite data are used for sea surface temperature (Marullo et al. 2010; Pisano et al. 2016; Pastor et al. 2018) (*Section 2.2.7*), and *in-situ* data for the deep water characteristics (Houpert et al. 2016; Schroeder et al. 2017; Testor et al. 2018; von Schuckmann et al. 2018).

2.2.1.2 Modelling

Complex and realistic global and regional climate models (GCMs and RCMs), based on fundamental physics, chemistry and biology equations are currently the standard tools to simulate the future evolution of the regional climate system. Different types of climate models are available to study the Mediterranean climate (past and future), often organized in large coordinated multi-model initiatives under the World Climate Research Programme (WCRP) umbrella CORDEX (Giorgi et al. 2009) and CMIP5 (Taylor et al. 2012). Combining the various sources of information or extracting the most credible (actionable) information is a new challenge, sometimes called the “distillation” problem (Hewitson et al. 2014; Fernández et al. 2019).

The MAR1 climate assessment concerning future climate evolution is based on four climate model ensembles, chosen for their good representation of the Mediterranean climate and for their good coverage of the various sources of uncertainty in future climate projections:

- CMIP3 and CMIP5 GCM ensembles with resolution ranging approximately from 300 to 100 km: they are the largest GCM multi-model ensembles available so far. They cover at a relatively low resolution all the uncertainty sources and can provide data for all the components of the climate system (atmosphere, land, ocean, marine biogeochemistry and aerosols). Some of the participating models share components and this may result in a redundancy in the ensemble results (Knutti et al. 2017).
- The Euro-CORDEX RCM ensemble is a large high-resolution ensemble at 12 km resolution (Jacob et al. 2014), which has clearly improved the representation of climate variables compared

to coarse resolution GCMs over land, e.g., for extreme precipitation (Fantini et al. 2018), regional winds (Obermann et al. 2018), mountain climate (Torma et al. 2015; Torma 2019), and over the sea, e.g., for regional winds (Herrmann et al. 2011) and extreme winds (“medicanes”) (Gaertner et al. 2018). An example for the strong modification of the future climate change signal by high-resolution RCM compared to GCMs has been found for summer precipitation over the Alps (Giorgi et al. 2016). Although the EURO-CORDEX ensemble is a large high-resolution multi-model dataset, it does not cover the entire uncertainty space of the CMIP5 ensemble.

- The Med-CORDEX RCM ensemble (Ruti et al. 2016) is a relatively small ensemble which does not cover the CMIP5 uncertainty range particularly well, but is the best data source available to study the future evolution of the Mediterranean Sea (Darmaraki et al. 2019b; Soto-Navarro et al. 2020), its ecosystems (Moullec et al. 2019) and atmosphere-ocean interactions. The models of this ensemble have high-resolution in both the atmosphere (resolution range: 25–50 km) and the ocean (resolution range: 6–30 km) component of the regional climate system (Somot et al. 2018).
- The CORDEX FPS-convection CPRCM ensemble (Coppola et al. 2020): this mini-ensemble provides the highest spatial resolution (2–3 km) for the greater Alpine region, reaching convection resolving scales. It yields, in particular, a strong improvement in the representation of extreme precipitation at sub-daily time scales (Kendon et al. 2014; Ban et al. 2015; Fosser et al. 2015; Berthou et al. 2018; Fumière et al. 2019). Convection resolving models are very promising tools to study the future evolution of extreme precipitation associated with thunderstorms, medicanes or mesoscale convective systems in the Mediterranean region (Lenderink et al. 2019) or urban-climate interactions.

A large variety of downscaling methods are available to study the Mediterranean climate (COST VALUE) (Maraun et al. 2019), including on-line tools on web processing servers (Cofiño et al. 2007). Among these methods, correcting climate change simulations using statistical tools (bias correction methods) allows to improve the present-climate statistics of climate simulations, with clear benefits for studying threshold-dependent extremes or for forcing impact models. All statistical methods require long-term observations (stations, gridded-products or satellite data calibrated for climate applications) for the learning phase and their application is therefore limited in regions where observations are lacking.

In the MAR1 report, CMIP ensembles are mostly used for the assessment of atmospheric dynamics, aerosol, cloud and radiation, water cycle, sea hydrology, sea level and acidification, whereas the Euro-CORDEX ensemble is used for the assessment of wind, clouds, temperature, precipitation and the cryosphere. The Med-CORDEX coupled regional models are used for sea hydrology and sea level, and the FPS-convection ensemble for the assessment of extreme precipitation.

Despite the continuous improvement of climate model ensembles by increased resolution and ensemble size, increased complexity and improved model physics, these still suffer from deficiencies and systematic errors. In particular their poor representation of some key regional phenomena may limit confidence for some aspects of the MAR1 assessment. This is especially true for coastal sea level, medicanes, tornadoes, hail phenomena, lightning, city climate, sub-daily precipitation, glaciers, clouds or cloud-aerosol interactions, human influence on land and water use.

2.2.2 General circulation and dynamics

2.2.2.1 General circulation

The proximity of the Mediterranean to the Atlantic and Indian Oceans and the surrounding massive land areas, with diverse climatic characteristics, places the area at the crossroads of many global climate patterns and processes of tropical and extra-tropical origin (Xoplaki et al. 2003a; Lionello et al. 2006; Lelieveld et al. 2012; Lionello et al. 2012a; Lionello 2012; Ulbrich et al. 2012). Its location on the eastern edge of the Atlantic Ocean means it is particularly affected by variability and change in the North Atlantic jet stream (or Polar Front Jet) in both winter and summer (Dunkeloh and Jacobbeit 2003; Hurrell et al. 2003; Athanasiadis et al. 2010; Bladé et al. 2012) and by stationary blocking patterns (Tyrilis and Hoskins 2008). The Mediterranean Basin is also influenced by semi-permanent large-scale anticyclones (e.g., the Azores anticyclone in the west during summer and the cold Siberian anticyclone in the northeast during winter), while mobile anticyclones play also important role throughout the whole year (Hatzaki et al. 2014). During the summer, the climate of the Mediterranean is further influenced by circulation patterns set up by the Asian summer monsoon (Rodwell and Hoskins 1996) and local orography (Simpson et al. 2015).

Observed trends

The wintertime large-scale circulation has exhibited a long-term trend toward increased sea-level pressure and anticyclonic circulation over the Mediterranean (*Fig. 2.2a and b*) (Mariotti and Dell'Aquila 2012). Aside from this long-term trend, the historical record has also exhibited sizable multi-decadal variability. This is illustrated for the sea-level pressure anomalies in *Fig. 2.2e* and has also been discussed widely in the contexts of trends in the North Atlantic Oscillation (NAO) and associated Mediterranean drying that occurred over the latter half of the 20th century (Hurrell 1995) in which internal variability is thought to play an important role (Kelley et al. 2012). CMIP5 models suggest that the externally forced contribution to sea-level pressure trends since 1900 consist of a pattern that resembles that of the observed trends but with a magnitude that is considerably smaller (*Fig. 2.2c*). It is likely that both external forcing and internal variability have contributed to the observed long-term trends (Hoerling et al. 2012). During summer, it is challenging to assess the observed long-term trends, as there is no strong agreement in the pattern of sea-level pressure change (*Fig. 2.2f and g*). There are indications of a summertime decline in sea-level pressure over North Africa and the southern Mediterranean and, indeed, the CMIP5 models suggest that external forcings have contributed to a decline in sea-level pressure in this region over the 20th century (*Fig. 2.2h*).

Future changes

Under rising greenhouse gas concentrations, climate models project that the Hadley Cell circulation will change, the tropics will expand and the mid-latitude westerlies and associated storm tracks will likely shift poleward (*medium/high confidence*) (Yin 2005; Lu et al. 2007b, 2007a; Chang et al. 2012; Barnes et al. 2013; Shaw et al. 2016; D'Agostino et al. 2017, 2020). This is expected to enhance subsidence and reduce storminess at the latitudes of the Mediterranean region, with a resulting reduction in precipitation (*medium confidence*). While there is considerable inter-model spread in the magnitude of these projected changes and the forced signal can be small compared to internal variability (Woollings and Blackburn 2012; Barnes et al. 2013; Zappa et al. 2015; Quan et al. 2018; Grise et al. 2019), the Mediterranean could be influenced by additional local circulation anomalies, leading to pronounced hydroclimate changes (Seager et al. 2014; D'Agostino and Lionello 2020). Future projections suggest that the wintertime trend toward increased anticyclonic circulation




Figure 2.2 | Trends in sea level pressure (SLP). [a]-[e] show the DJF (December-January-February) season. [a] and [b] show 1900-2018 trends in SLP from ERA20C and 20CR reanalyses, respectively. Grey = not significantly different from zero at the 95% level. Significance is calculated by resampling, with replacement, the residuals of the linear trend, adding the resampled residuals to the linear trend and re-calculating the linear trend. This is repeated 1,000 times to obtain the probability (p-value) at each grid point that the trend is significantly different from zero. Spatial autocorrelation is accounted for using the False Discovery Rate method of Wilks (2016) with control value = 0.1. Stippling depicts grid points where the magnitude of the trend is larger than in any of the individual CMIP5 ensemble members. [c] shows the CMIP5 multi-model mean trend for 1900-2018. The ensemble mean for each model is calculated, then the linear trend is obtained before calculating the average trend across all models. Grey depict regions where less than 3/4 of the models agree on the sign of the change. [d] is as [c] but for future trends from 2019-2099. [e] shows time series of 20 year running mean SLP averaged over the Mediterranean (green box in c) for the two re-analyses, all individual ensemble members from all models and the CMIP5 multi-model mean. [f]-[j] are as [a]-[e] but for JJA (June-July-August).

over the Mediterranean will continue at an accelerated pace (*Fig. 2.2d*) (Giorgi and Coppola 2007). This is accompanied by a strengthening of the westerly winds and increased storminess over northern Europe, reduced westerlies over north Africa

and decreasing storminess over the Mediterranean (Woollings and Blackburn 2012; Rojas et al. 2013; Zappa et al. 2013). Climate models predict a summertime poleward shift of the North Atlantic jet (Simpson et al. 2015) and a summertime de-

crease/wintertime increase in sea-level pressure centered over the Mediterranean (*Fig. 2.2i*) (Giorgi and Coppola 2007; Bladé et al. 2012), with the reduction in Mediterranean sea-level pressure continuing at an accelerated pace over the coming decades (*Fig. 2.2i*), albeit with a large inter-model spread (*Fig. 2.2j*). This was argued to be dominated by a heat-low response to drier soils in the Mediterranean region (Haarsma et al. 2009).

2.2.2.2 Teleconnection patterns

The influence of teleconnection patterns (linkages between weather changes occurring in widely separated regions of the globe) on Mediterranean climate has been extensively studied (Corte-Real et al. 1995; Hurrell and Van Loon 1997; Wibig 1999; Pozo-Vázquez et al. 2001; Quadrelli et al. 2001; Xoplaki et al. 2003a, 2003b, 2004; Hatzaki et al. 2009; Toreti et al. 2010; Ulbrich et al. 2012; Tyrlis and Lelieveld 2013; Rousi et al. 2015; Sen et al. 2019). Particularly during winter, the region is prone to the impacts of the westerly flow and the teleconnection patterns of North Atlantic Oscillation (NAO), Eastern Atlantic/Western Russia (EA/WR) and Scandinavian (SCAN) (Barnston and Livezey 1987).

The NAO is, for parts of the region, one of the most important modes of internal climate variability. It affects especially the storm-tracks and cyclogenesis over parts of the basin (Trigo et al. 2000, 2004; Reale and Lionello 2013) and mainly precipitation over the western-central Mediterranean in winter (Lamb and Peppler 1987; Rodríguez-Fonseca and de Castro 2002; Xoplaki et al. 2004). NAO is also found to have some influence on winter precipitation in parts of the eastern Mediterranean, however this influence is smaller (Dunkeloh and Jacobit 2003; Xoplaki et al. 2004; Feliks et al. 2010; Felis and Rimbu 2010; Nissen et al. 2010). A lesser but distinct influence is detected between NAO and the Mediterranean surface air temperature (Hurrell 1995; Cullen and DeMenocal 2000; Ben-Gai et al. 2001; Pozo-Vázquez et al. 2001; Sáenz et al. 2001; Castro-Díez et al. 2002; Trigo et al. 2002b; Türkeş and Erlat 2003; Xoplaki et al. 2003b; Toreti et al. 2010).

Observed trends of NAO are not monotonic and are difficult to assess since decadal oscillations are too large to reach a conclusion with an acceptable level of confidence. Nevertheless, mostly negative trends prevail since the early 1990s following a positive trend starting in the 1960s (Ulbrich and Christoph 1999; Mariotti et al. 2002b; Türkeş and Erlat 2003; Trigo et al. 2004, 2006; Xoplaki et al.

2004; Pinto and Raible 2012; Saffioti et al. 2016; Iles and Hegerl 2017).

Climate projections mostly suggest a weak positive NAO trend in a warmer future climate (*low/medium confidence*), accompanied by a small northeastward displacement of its centers-of-action by the end of the 21st century (Ulbrich and Christoph 1999; Gillett et al. 2003, 2013; Hu and Wu 2004; Stephenson et al. 2006; Bacer et al. 2016; Deser et al. 2017; Barcikowska et al. 2020). Some studies indicate no significant trends (Fyfe et al. 1999; Dorn et al. 2003; Rauthe et al. 2004; Fischer-Brunns et al. 2009), or even decreasing trends (Osborn et al. 1999).

Studies on the effect of El Niño Southern Oscillation (ENSO) phenomenon on Mediterranean precipitation have shown that links exist, particularly during autumn and spring in the western Mediterranean and during winter in the Eastern Mediterranean. However, results are not conclusive concerning their evolution and robustness. In fact, the ENSO signal is difficult to be isolated, because of the dominating mid-latitude dynamics, the sign of its correlation with total precipitation depends on season and it is not stationary (Rodó et al. 1997; Rodó 2001; Mariotti et al. 2002b, 2005; Knippertz et al. 2003; Hasanean 2004; Alpert et al. 2006; López-Parages and Rodríguez-Fonseca 2012; Kalimeris et al. 2017).

2.2.2.3 Extratropical cyclones and medicanes

The Mediterranean is one of the main cyclogenetic areas of the world (Petterssen 1956; Hoskins and Hodges 2002; Wernli and Schwierz 2006), with much of the high-impact weather (e.g., strong winds and heavy precipitation) associated with cyclonic structures. Cyclogenesis areas such as the north-western Mediterranean, North Africa, the north shore of the Levantine Basin, the seasonality (Alpert et al. 1990a, 1990b; Trigo et al. 1999, 2002a; Lionello et al. 2006, 2016; Campins et al. 2011), as well as the occurrence of explosive cyclogenesis (Kouroutzoglou et al. 2011; Reale et al. 2019) are well documented in the literature. Within Mediterranean cyclones, there is a sub-group of hybrid depressions of extratropical cyclogenesis, the so-called 'medicanes' (Mediterranean hurricanes) or tropical-like cyclones (Rasmussen and Zick 1987; Reale and Atlas 2001; Emanuel 2005). These are mesoscale maritime extratropical cyclones that can physically emulate tropical characteristics at a certain point of their life cycle (Emanuel 2005; Miglietta 2019). Such features can include a cloud-

free area at the center (the “eye”), spiral bands with deep convection around it, intense surface winds and a warm-core and symmetric structure (Miglietta et al. 2015). These events can pose serious societal and ecological threat to the affected coastal regions (Nastos et al. 2018).

During the recent past there is an absence of strong trends in cyclone numbers affecting the Mediterranean (Lionello et al. 2016), however when trends are detected these are mostly negative (*low/medium confidence*) (Trigo et al. 2000; Maheras et al. 2001; Flocas et al. 2010; Nissen et al. 2010). Similarly, the number of explosive Mediterranean cyclones has likely decreased, but this reduction is not statistically significant (Kouroutzoglou et al. 2010). The statistical record of medicanes has limited reliability and sample size, given their maritime characteristics, small size and infrequent occurrence. Thus, it has not been possible to derive an objective climatology. Observational studies cannot be used to identify trends because the identification is commonly subjective (Miglietta et al. 2013; Tous and Romero 2013; Nastos et al. 2018). Dynamical downscaling methods have been used to build a medicane climatology, but only negligible trends were obtained (Cavicchia et al. 2014).

For the future, climate models project a reduction in the number of cyclones (*medium/high confidence*) especially in winter (Lionello et al. 2002; Geng and Sugi 2003; Bengtsson et al. 2006; Leckebusch et al. 2006; Lionello and Giorgi 2007; Pinto et al. 2007; Löptien et al. 2008; Ulbrich et al. 2009; Raible et al. 2010; Zappa et al. 2013; Nissen et al. 2014). There is some uncertainty, as the spread in the model responses appears to be quite large (Ulbrich et al. 2008, 2009; Harvey et al. 2012). With respect to cyclone intensity, climate projections are more controversial, as some models suggest a decrease in the frequency of the most intense systems (Pinto et al. 2007; Raible et al. 2010), while other models show more extreme events or increases in the intensity of extreme cyclones (Lionello et al. 2002; Gaertner et al. 2007). For medicanes, climate projections indicate a decreasing response in frequency but increasing intensity (Gaertner et al. 2007; Romero and Emanuel 2013, 2017; Cavicchia et al. 2014; Walsh et al. 2014; Tous et al. 2016; Romera et al. 2017; González-Alemán et al. 2019).

2.2.2.4 Regional winds

Surface wind speed and its changes on different temporal and spatial scales are governed by driving

and drag forces, where all relevant contributions are difficult to estimate and disentangle (Wu et al. 2018a). Surface wind climate studies are less common than air-temperature and precipitation studies for example, and more work is needed to explain historical wind speed evolution and precisely estimate different sources of uncertainty in the future projections. This variable is now becoming more important, since parts of the region, both inland and offshore, have high potential for the production of wind energy (Balog et al. 2016; Onea et al. 2016) (*Chapter 3.3*).

Observation-based studies of winds over the Mediterranean are rare, and depend on the availability of homogenized and long time series. In most regions, wind trends were found non-monotonic over the past decades and, concrete conclusions are difficult to be established (Pirazzoli and Tomasin 2003; Vautard et al. 2010; Azorin-Molina et al. 2014). An additional source of information are reanalysis datasets, but robust trends have been identified over only a few regions in the Mediterranean (Nissen et al. 2010; Donat et al. 2011; Bett et al. 2013). Climate model simulations over historical periods can also be used in assessing and understanding past trends (Knippertz et al. 2000).

Despite the uncertainties in future projections (Shepherd 2014; Belušić Vozila et al. 2019), there is a general agreement for a limited wind speed reduction over most of the Mediterranean, with the exception of the Aegean Sea and north eastern land areas (*Fig. 2.3, Section 2.2.8*) (*medium confidence*) (Somot et al. 2006; McInnes et al. 2011; Dobrynin et al. 2012; Planton et al. 2012; Belušić Vozila et al. 2019). Changes in the local winds (such as Bora, Mistral, Tramontane, Sirocco and Etesians) may have more complex responses involved, depending on the changes in their underlying feedbacks (Grisogono and Belušić 2009; Ulbrich et al. 2012). Regional projections over the Adriatic reveal strong sensitivity in the climate change signal of the local Bora and Sirocco winds (Belušić Vozila et al. 2019). In particular, the frequency of winter Bora events is projected to increase while the frequency of Sirocco events is expected to decrease. Overall, the mean wind speed during Bora and Sirocco events is expected to be reduced, with the exception of Bora in northern Adriatic. RCM projections of Mistral and Tramontane show small changes in the former and significant decrease in the frequencies of the latter (Obermann-Hellund et al. 2018). Etesian winds over the Aegean Sea is one of the few exceptions since increases in the wind speed are expected for the future (Ezber 2018; Dafka et al. 2019). In general, RCM projections have the




Figure 2.3 | Projected changes (%) in surface wind speed based on Med-CORDEX simulations (Ruti et al. 2016) for the end of the 21st century (2071–2100) relative to the period 1961–1990, for pathways RCP4.5 (top panel) and RCP8.5 (bottom panel). Seven RCMs were used for RCP4.5 and eight RCMs were used for RCP8.5. Dotted areas show differences that passed a 95% significance test.

tendency to simulate decrease of the wind energy density over the Mediterranean, with the exception of the Aegean Sea (Hueging et al. 2013; Tobin et al. 2015; Moemken et al. 2018).

Regional climate simulations indicate changes in wind speed over land regions as well (*Fig. 2.3*). Most pronounced changes, consistent for both RCP4.5 and RCP8.5 pathways, are an increase of wind speeds over the Balkans and a decrease over the Alps.

2.2.3 Radiation, clouds and aerosols

The amount of solar radiation reaching the Earth's surface is a key determinant of the spatio-temporal variations of climate on our planet and is the ultimate energy source for all processes relevant for climate and life. The main factors explaining the variability of surface solar radiation over the Mediterranean region for different time scales (daily, seasonal, interannual, past and future trends) are clouds (Pyrina et al. 2015) and aerosols (Nabat et al. 2015a, 2015b). The daily variability in clouds and aerosols is strongly influ-

enced by weather regimes (Rojas et al. 2013; Nabat et al. 2020) and climate oscillations such as the North Atlantic Oscillation, NAO (Chiacchio and Wild 2010).

Aerosols in the Mediterranean come from various and numerous sources (Lelieveld 2002), both natural, notably dust and sea-salt, and anthropogenic, notably sulfates, nitrates and black carbon (*Section 2.3*). Their interactions with radiation and clouds are essential in understanding climate in this region.

2.2.3.1 Observed change in surface radiation

The long-term solar radiation records taken at widespread locations around the globe underwent substantial multidecadal variations, characterized by a reduction of surface solar radiation from the 1950s to the 1980s, known as "global dimming" and a partial recovery, thereafter, referred to as "brightening" (Wild 2009, 2012; Wild et al. 2017). This dimming/brightening pattern is also observed




Figure 2.4 | Past and future evolution of Surface Downwelling Shortwave Radiation (RSDS in W m⁻²), Total Cloud cover (CLT in %) and Aerosol Optical Depth (AOD) from 1950 to 2100 averaged over the Mediterranean area in CMIP5 Global Climate Models (GCM) and 12-km Euro-CORDEX Regional Climate Models (RCM) ensembles. Figures based on Bartok et al. (2017), redrawn and extended using published datasets.

in the Mediterranean area (*very high confidence*), both in all-sky and clear-sky conditions, documented with many ground-based and satellite observations after careful data quality assessment and homogenization (Sánchez-Lorenzo et al. 2007, 2013, 2017; Zerefos et al. 2009; Kambezidis et al. 2016; Manara et al. 2016; Alexandri et al. 2017; Pfeifroth et al. 2018), as well as climate simulations (Folini and Wild 2011; Zubler et al. 2011; Nabat et al. 2014). The surface solar radiation trends averaged over the Mediterranean have been estimated in climate model simulations between -3.5 and -5.2 W m⁻² per decade for the dimming period (1953-1968), against for the brightening period between +0.9 and +4.6 W m⁻² per decade in 1989-2004 (Folini and Wild 2011) and 2.3 W m⁻² per decade in 1980-2012 (Nabat et al. 2014). The surface solar radiation anomalies calculated by global climate models for the periods 1950-1969 and 2000-2019 against the reference period of 1980-1999 are positive, showing respectively the dimming and the brightening effects, since the reference period refers to the period where surface solar radiation was the lowest (Fig. 2.4).

In parallel to the brightening period, a decrease in aerosol loads has been observed since 1980 both in ground-based stations (Li et al. 2014) and in satellite data (Floutsi et al. 2016). This decrease, corresponding to a trend in aerosol optical depth of -0.03 decade⁻¹, is mainly due to reductions in anthropogenic emissions, leading to a decrease in anthropogenic aerosol concentrations such as sulfate (Nabat et al. 2013). These aerosol trends have been shown to be the main explanation of the dimming-brightening phenomenon in the Mediterranean area (*high confidence*) through attribution model studies (Folini and Wild 2011; Zubler et al. 2011; Nabat et al. 2014), and with the direct aerosol effect responsible for about 80% of the simulated brightening. This phenomenon is qualitatively reproduced by most GCMs, but only by a few regional climate models, due to different treatments of aerosols in models (Fig. 2.4). The evolution of natural aerosols is more uncertain over the same period.

Concerning clouds, a decrease in cloud cover of 0.63% per decade since the 1970s has also been detected from different observations datasets over

the Mediterranean area (Sánchez-Lorenzo et al. 2017). This trend mainly concerns low and mid cloud layers (Kambezidis et al. 2016) (*medium confidence*). The spread between models that capture these trends is high, because of their difficulties to capture cloud characteristics (*Fig. 2.4*). Clouds may also have played a significant role in the past trend of surface solar radiation, at least locally (*low confidence*). Stronger positive trends in surface solar radiation are detected in spring over western Mediterranean Basin, explained by a decrease in cloud optical depth for this season over this basin (Kambezidis et al. 2016), and despite an averaged positive trend of surface solar radiation observed over the eastern basin, more uncertainty exists due to the lack of observations of both clouds and aerosols (Alexandri et al. 2017).

2.2.3.2 Projected change in surface radiation

In future climate projections, anthropogenic aerosol loads over the Mediterranean are expected to keep on decreasing (*high confidence*) because of decreases in anthropogenic emissions in Europe (Shindell et al. 2013). The decrease is expected to be more pronounced in the near future with an aerosol optical depth anomaly between -0.12 and -0.03 (5-95% uncertainty range) for the 2040-2059 period with respect to the reference period 1980-1999 (*Fig. 2.4*), and will slow down in the far future (between -0.18 and -0.04, 2080-2099 vs 1980-1999). The evolution of natural aerosols is more uncertain, due to current unknown future evolution of the desert dust (*Section 2.3.2*).

Total cloud cover is also expected to decrease during the 21st century over the Mediterranean (*medium confidence*) (Boé and Terray 2014; Enriquez-Alonso et al. 2016; Bartók et al. 2017; Hentgen et al. 2019). This is consistent with the northward expansion of the Hadley cell (Sánchez-Lorenzo et al. 2017; D'Agostino et al. 2020) (*Section 2.2.2*) and with enhanced lower tropospheric drying (Hentgen et al. 2019). The expected anomaly in cloud cover for the mid-21st century ranges from -4.9 to -0.2% in the RCP8.5 (5-95% uncertainty range, *Fig. 2.4*), because of the difficulty of models to capture the spatial variability of the cloudiness evolution (Bartók et al. 2017).

As projected by GCMs, surface solar radiation is expected to continue increasing in the 21st century, especially in the near future (*medium confidence*). The anomaly is between 0.6 and 7.7 W m⁻² for the period 2020-2039 in the RCP8.5 (5-95% uncertainty

range), in line with the decrease in anthropogenic aerosols (Boé et al. 2020; Gutiérrez et al. 2020) (*Fig. 2.4*). However, this evolution is not shared by all RCMs (Bartók et al. 2017; Gutiérrez et al. 2020). Only regional climate models which consider aerosol dynamics simulate the increase in surface solar radiation as the global models (Boé et al. 2020; Gutiérrez et al. 2020). This increase in future surface solar radiation is reinforced by an expected decrease in cloud cover (Enriquez-Alonso et al. 2016), despite a compensational effect of increased absorption in clear sky conditions due to higher water vapor content in the atmosphere (Haywood et al. 2011) (*Fig. 2.4*).

2.2.4 Temperature and related extremes

2.2.4.1 Observed temperature changes

Climate reconstructions, ground-based observations, reanalysis and remote-sensing datasets all corroborate the transition to warmer conditions during the 20th century and that warming has accelerated during the last decades (*high confidence*). Basin-wide, annual mean temperatures are now 1.5°C above late 19th century levels (*Box 2.1, Fig. 2.33*). Particularly after the 1980s, regional warming has accelerated and increases at a higher rate than the global average (Lelieveld et al. 2012; Lionello et al. 2012a; Zittis and Hadjinicolaou 2017; Cramer et al. 2018; Lionello and Scarascia 2018; Zittis et al. 2019). These studies present a strong consensus that the recent observed warming is robust throughout the region analysis, though magnitude and level of significance of the observed temperature trends in the Mediterranean varies depending (a) on the region, country or station under consideration, (b) on the type of data set investigated and (c) on the season and period of analysis.

Solar forcing and large volcanic eruptions are found to have a strong influence on the Mediterranean temperature variability over the last centuries (Trouet 2014). A combination of climate reconstructions, documentary sources and observed data suggests that looking at the long-term timescale (e.g., over the last 500 years), warm periods are not exceptional for the Mediterranean, which is characterized by a sequence of warming-cooling cycles (Luterbacher and Xoplaki 2003; Camuffo et al. 2010; Lelieveld et al. 2012). A study of summer temperature since Roman times shows that although the mean 20th century European (including the northern part of the Mediterranean Basin) was not significantly warmer than some




Figure 2.5 | Observed temperature trends (left panels) and time-series of temperature over land for the Mediterranean based on the Climatic Research Unit (CRU) [Harris et al. 2020] gridded observations. Time-series refer to the whole Mediterranean as defined in the left panels and for three Mediterranean sub-regions (*Fig. 2.1*): North (FR, ALBA), Central (IB, CM, AN) and South Mediterranean (WM, EM, LE).

earlier centuries, there are no earlier 30-year periods found to be warmer than the most recent 3 decades (Luterbacher et al. 2016).

Recent climatic trends are clear, particularly after the 1980s. Over these last decades, according to different type of observations, significant positive trends of the order of 0.1–0.5°C per decade have been identified (*Fig. 2.5*) (Nasrallah and Balling 1993; Saaroni et al. 2003; Feidas et al. 2004; Brunetti et al. 2006; El Kenawy et al. 2009; Tanarhte et al. 2012; Lelieveld et al. 2012; Lionello 2012; Donat et al. 2014; Mariotti et al. 2015; Lionello and Scarascia 2018; Bilbao et al. 2019). In addition, for parts of the basin, there is some evidence that the diurnal temperature range has also changed (Price et al. 1999; Bilbao et al. 2019; Sun et al. 2019).

Besides mean values, hot and cold extremes have also become warmer, while in particular there is strong evidence and consensus that heat waves have become more frequent and severe. Various climatic indicators show significant increasing trends of extreme heat events characteristics (e.g., duration, frequency and intensity). The number of warm and tropical nights has also increased over most Mediterranean locations including Iberia,

north Africa, Italy, Malta, Greece, Anatolia and the Levant (Kostopoulou and Jones 2007; Bartolini et al. 2008; Kuglitsch et al. 2010; El Kenawy et al. 2011; Galdies 2012; Donat et al. 2014; Filahi et al. 2015; Lelieveld et al. 2016; Ceccherini et al. 2017; Nashwan et al. 2018; Tolika 2019). Parts of the region were impacted by some of the most severe record-breaking weather events of the last decade, mainly related with summer heat extremes (Coumou and Rahmstorf 2012). Considering only winter, some studies that suggest a different behavior of hot and cold extremes between the eastern and western parts of the Mediterranean, with negative temperature trends in the former and positive trends in the latter (Hertig et al. 2010; Efthymiadis et al. 2011), but these studies do not include the most recent warm decades.

2.2.4.2 Future temperatures

According to future projections, the greater Mediterranean Basin is among the most responsive regions to global warming. Previous studies have identified the region as one of the most prominent climate change hot spots (Giorgi 2006; Lionello et al. 2006; Giorgi and Lionello 2008; Diffenbaugh and Giorgi 2012; Lionello 2012; Lionello and Scarascia




Figure 2.6 | Projected changes in annual temperature between the recent past reference period (REF: 1980-1999) and three future sub-periods (NEAR: 2020-2039, MID: 2040-2059, END: 2080-2099), based on the ensemble mean of EURO-CORDEX 0.11° simulations for pathways RCP2.6 (top panels) and RCP8.5 (bottom panels).

2018). Multi-model ensembles of climate simulations indicate that widespread warming will almost certainly occur in the Mediterranean in the 21st century (*high confidence*), though, climate models might overestimate actual values in warm and dry conditions (Boberg and Christensen 2012).

The warming level of warming strongly depends on the reference period definition, the future time horizon and the level of greenhouse gas forcing (Christensen et al. 2007; Giorgi and Lionello 2008; Collins et al. 2013; Dubrovský et al. 2014; Jacobbeit et al. 2014; Mariotti et al. 2015; Ozturk et al. 2015; Lionello and Scarascia 2018; Zittis et al. 2019). A quantitative estimation based on state-of-the-art EURO-CORDEX regional simulations is presented in *Table 2.1* and *Fig. 2.6-2.7*. Over land regions, a robust and significant warming of the range of 0.9-5.6°C (with respect to the reference period 1980-1999) is suggested for the future. The robustness and significance of the climate signal is much higher for air temperature rather than other variables such as precipitation (Knutti and Sedláček 2012; Lelieveld et al. 2016). There are strong indications and a general consensus that regional warming will continue faster than the global average and will exceed the global mean value by 20% on an annual basin and 50% in summer (*high confidence*) (Vautard et al. 2014; Dosio and Fischer 2018; Lionello and Scarascia 2018; Nikulin et al. 2018). Daytime temperatures are expected to increase more than nighttime temperatures, indicating an increase of the amplitude of the diurnal temperature range (Lionello and Scarascia 2018).

Changes in the occurrence of extreme events closely follow changes in inter-annual variability. Therefore, such changes can be also considered as a proxy measure of seasonal extremes (Schär et al. 2004; Giorgi 2006). The intensity of extreme temperature is projected to increase more rapidly than the intensity of more moderate temperatures over the continental interior due such increases in temperature variability (Beniston et al. 2007).

Projected changes in extreme temperature indicators suggest that the frequency and severity of heat waves will increase (*high confidence*) (Diffenbaugh et al. 2007; Goubanova and Li 2007; Giorgi and Lionello 2008; Fischer and Schär 2010; Diffenbaugh and Giorgi 2012; Sillmann et al. 2013; Russo et al. 2014; Jacob et al. 2014; Kostopoulou et al. 2014; Zittis et al. 2016; Lelieveld et al. 2016; Ouzeau et al. 2016; Lionello and Scarascia 2020). According to projections for a business-as-usual scenario, summer daily maximum temperature is expected to increase up to 7°C by the end of the 21st century in comparison with the recent past (Sillmann et al. 2013; Lelieveld et al. 2016). Besides warmer daytime temperature maxima, parts of the Mediterranean will likely face an increase of more than 60% in the number of tropical nights. Increase of warm temperature extremes will be dramatic particularly in summer and with a 4°C global warming almost all nights will be warm and there will be no cold days (Sillmann et al. 2013; Dosio and Fischer 2018; Lionello and Scarascia 2020).




Figure 2.7 | Time-series of simulated mean annual (top panel), summer (middle panel) and winter (bottom panel) temperature averaged over the Mediterranean based on EURO-CORDEX 0.11° simulations for historical times (black curve) and future pathways RCP2.6 (blue curve) and RCP8.5 (red curve). Solid lines indicate the ensemble means and shaded areas the spread of the simulations. Box-plots represent the averages over the decade 2091-2100 in terms of model spread.

Warming is projected to be milder in winters and much stronger during summers. This is mainly attributed to land-atmosphere interactions and the transition to drier conditions (Seneviratne et al. 2006; Jaeger and Seneviratne 2011; Quesada et al. 2012; Zittis et al. 2014). Another important feedback, particularly for southern Mediterranean, is the coupling of longwave radiation between the desert soil surface and lower atmosphere which

amplifies warming and intensifies the summertime heat low over the Sahara (Cook and Vizy 2015; Evan et al. 2015). The exceptional summertime warming over parts of the region is also likely associated with a thermal low, which is explained by the widening of the Persian trough that extends from South Asia to the eastern Mediterranean, and is projected to expand westward and combine with the intensifying thermal low over the Sahara (Lelieveld et al. 2016).

2.2.5 Precipitation, related extremes and the water cycle

2.2.5.1 Observed trends in precipitation

Observed precipitation trends during the full or portions of the 20th century up to present day, covering the full or portions of the Mediterranean Basin, are available in gridded format from various sources, such as CRU, UDEL, E-OBS, EURO4M. Annual, DJF (December-January-February) and JJA (June-July-August) precipitation trends from the CRU dataset for different time periods, 1950-2018 and 1980-2018 are shown in Fig. 2.8. Fig. 2.9 shows the temporal evolution of land precipitation (1901-2018) averaged over the full Mediterranean area and its northern, central and southern portions (defined as the sum of the three northern, central and southern regions of Fig. 2.8, respectively).

The sign of the observed precipitation trend exhibits pronounced spatial variability and depends on the time period and season considered (Fig. 2.8). For example, the period 1950-2018 shows a prevailing decreasing trend over most of the Mediterranean Basin of annual and winter precipitation, which is reversed over large portions of the basin if we only consider the period 1980-2018. This is because of the marked multidecadal variability of precipitation in the Mediterranean, which may actually mask trends induced by greenhouse gas emissions. The prominent role of multidecadal variability is also evident when precipitation is regionally averaged (Fig. 2.9). In this case the most evident trend is a decrease of winter precipitation over the central and southern portions of the basin since the second half of the 20th century. Overall, because of the marked multidecadal variability of precipitation and the small magnitude of trends, the confidence in the detection of trends from greenhouse gas emissions for the historical past is low (Lelieveld et al. 2012; Lionello et al. 2012a; Peña-Angulo et al. 2020; Vicente-Serrano et al. 2020).




Figure 2.8 | Observed annual, DJF, JJA precipitation trends from the CRU dataset. Left and right panels consider the 1950-2018 and 1980-2018 periods, respectively.




Figure 2.9 | Time series of annual, DJF, and JJA precipitation over land from the CRU dataset. Time-series refer to the whole Mediterranean as defined in the left panels and three Mediterranean sub-regions (Fig. 2.1): North (FR, ALBA), Central (IB, CM, AN) and South Mediterranean (WM, EM, LE).

2.2.5.2 Future precipitation

Mean precipitation changes

21st century precipitation projections for the Mediterranean region have been produced based on GCM and RCM ensembles of experiments. Analyses of GCM projections over the Mediterranean region have been conducted for CMIP3 (Giorgi and Coppola 2007; Giorgi and Lionello 2008; Mariotti et al. 2008) and CMIP5 (Mariotti et al. 2015; Lionello and Scarascia 2018). Several generations of RCM-based projections for the EURO-Mediterranean region are also available from projects such as PRUDENCE (Déqué 2007), ENSEMBLES (Déqué et al. 2012) and EURO-CORDEX (Jacob et al. 2014). In addition, projections based on coupled regional atmosphere-ocean models have been conducted as part of the CIRCE (Gualdi et al. 2013) and MEDCORDEX (Ruti et al. 2016) projects.

A consistent dominant signal emerges from these projections, consisting of a predominant drying through the entire Mediterranean Basin in the warm seasons (April through September, with largest magnitude in JJA), drying in most central and southern areas along with wetting in the northernmost regions (e.g., the Alps) in the winter season (*medium confidence*). This large-scale pattern of change is illustrated for the Euro-CORDEX dataset, at the annually averaged scale, in Fig. 2.10 for the RCP2.6 and RCP8.5 scenarios

and different future time slices. Table 2.1 provides quantitative values of precipitation change for different scenarios and model ensembles over the Mediterranean region.

In general, the patterns of change intensify in magnitude from the near future to the far future time slices and from the low to high greenhouse gas emission scenarios, i.e., they intensify with the anthropogenic forcing and resulting global warming. As a result, for example, at the Mediterranean scale, the CMIP5 ensemble yields a decrease of annual precipitation over the Mediterranean area of about 4% per degree of global warming (Lionello and Scarascia 2018).

The magnitude and pattern of precipitation decrease vary widely across models. For example, the summer precipitation reduction in the CMIP3 and CMIP5 datasets for the high-end greenhouse gas emission scenarios (roughly equivalent to the RCP8.5) varied from less than 10% to over 40% across models (IPCC 2007; Giorgi and Lionello 2008; Lionello and Scarascia 2018). Although qualitatively consistent, different ensembles show different sensitivities over the Mediterranean. The CMIP5 GCM ensemble produced a less pronounced summer drying than the CMIP3 one, when expressed in terms of change per degree of global warming (IPCC 2013b). RCM-based projections, e.g., as part of Euro-CORDEX yield a lower drying than GCM-based ones, with reduced areas




Figure 2.10 | Maps showing EURO-CORDEX-based change in annual, winter and summer precipitation change, for the RCP2.6 and RCP8.5 scenarios and the near-future, mid-term and far-future with respect to the reference period.

of precipitation decrease in the future compared to the GCMs. Considering a range of global warming of 0.9–5.6°C (*Section 2.2.4.2*) at the end of the 21st century (with respect to the reference period 1980–1999) and a decrease of 4% per degree of global warming, this gives a reduction between 4 and 22% for the Mediterranean annual precipitation (on land).

Fig. 2.11 shows the temporal evolution of Mediterranean scale precipitation (RCP2.6 and RCP8.5) from 1950 to 2100 in the EURO-CORDEX dataset, and presents both mean and inter-model range of data. The large inter-model spread includes some positive values and two scenarios start to separate, at least in an ensemble mean sense, only after the mid of the 21st century. Therefore, while it is possible to assess that precipitation will likely decrease over the Mediterranean Basin, at least under the higher end scenarios, it is difficult to assign robust quantitative values, especially at sub regional scale.

The uncertainty in projections is even larger as we move to the local scale because of the effects of local forcings, e.g., topography and coastlines. For example, focusing on the high elevations of the Alpine chain, Giorgi et al. (2016) found that the RCM projections at 12 km resolution exhibit an increase of summer precipitation in areas where the GCMs project a decrease. This is due to the occurrence of increased convection related to high elevation warming and heating. As another example, increases of cold season precipitation in the upwind side of mountain chains and decreases in the lee side have been found by the high-resolution simulations of Gao et al. (2006) in response to the topographically-forced precipitation shadowing effect. These results were confirmed by the analysis of the EURO-CORDEX ensemble projections (Kotlarski et al. 2019). In other words, high resolution RCM projections suggest that care needs to be taken when extending large scale patterns to the local scale, since local scale changes can be heavily affected by topography and coast lines.

Variability and extremes

Several studies have assessed changes in interannual variability of precipitation over the Mediterranean region in GCM-based projections, mostly using as a measure of variability the coefficient of variation (i.e., the interannual standard deviation divided by the mean), which removes the strong dependence of precipitation standard deviation from the mean (Räisänen 2002;




Figure 2.11 | Time-series of simulated mean annual (top panel), wet (middle panel) and dry (bottom panel) season over the Mediterranean land areas based on EURO-CORDEX 0.11° simulations for historical times (black curve) and future pathways RCP2.6 (blue curve) and RCP8.5 (red curve). Solid lines indicate the ensemble means and shaded areas the spread of the simulations. Box-plots represent the averages over the decade 2091–2100 in terms of model spread.

Giorgi and Bi 2005; Giorgi and Coppola 2009; Giorgi et al. 2019]. They all found a prevailing increase in precipitation variability over the Mediterranean, especially over areas showing strong precipitation decreases, which thus appears to be a robust response in global climate projections. However, this result may not apply if the metrics used to measure variability is the standard deviation.

Giorgi et al. (2011, 2014) carried out an analysis of changes in different hydroclimatic indices from an ensemble of CMIP5 projections and, on an annual basis, consistently found an increase in mean daily precipitation intensity and 95th percentile of daily precipitation, a decrease in precipitation frequency and wet spell length and an increase in the number of dry days and dry spell length (*medium confidence*). The hydroclimatic intensity index introduced by Giorgi et al. (2011), which is essentially the product of precipitation intensity and mean dry spell length, shows a consistent increase throughout the Mediterranean Basin. These conclusions based on GCM projections, were essentially confirmed by an analysis of high-resolution EURO-CORDEX projections by Jacob et al. (2014), except for a slight decrease of 95th percentile daily precipitation over some areas of the Iberian, Italian and Hellenic peninsulas and southern France in summer. A recent study (Lionello and Scarascia 2020), based on CMIP5, shows that global warming will further increase the existing difference in intensity of precipitation and hydrological extremes between North and South Mediterranean areas. Both the daily precipitation intensity the total precipitation during extreme events are already larger in the North than in the South Mediterranean areas, and differences will increase with global warming. The projected increase of dry spell length is larger in the south than in the North Mediterranean (*medium confidence*).

In conclusion, (*high confidence*) both global and regional climate projections indicate a predominant shift towards a precipitation regime of higher interannual variability (when measured by the coefficient of variation), higher intensity of precipitation and greater extremes (especially in winter, spring and fall, but not in the southern areas), decreased precipitation frequency and longer dry spells (especially in summer). This hydroclimatic response to global warming is greater for the RCP8.5 than the RCP2.6 scenario and for the far future vs. the near future time slices (*high confidence*).

2.2.5.3 Changes in evaporation, net water losses over sea and over land

Evaporation in the Mediterranean not only provides moisture locally, but also results in a net export of water to neighboring areas, primarily to the South and East (Mariotti et al. 2002a; Nieto et al. 2006). The Mediterranean Sea is the dominant regional evaporation source, and changes in Mediterranean Sea evaporation impact the

sea's water, salt and heat budgets. Large-scale internal climate variability and greenhouse gas forced global change have been primary drivers of Mediterranean evaporation changes during the 20th century and into the 21st century, while local processes have acted to modulate those effects (Mariotti and Dell'Aquila 2012). Overall, the net surface water loss (evaporation minus precipitation over the sea) has increased over most of the Mediterranean surface, mainly due to a decrease of precipitation during the period 1960-1990 and a strong evaporation increase since the mid-seventies due to local warming (Mariotti 2010; Sevault et al. 2014; Mariotti et al. 2015; Skliris et al. 2018). The freshwater discharge due to the river runoff has also decreased (Ludwig et al. 2009). Projected regional warming trends point to continuing increases in Mediterranean Sea evaporation, land drying in southern areas during summer and a net regional water loss.

Observations-based estimates of Mediterranean Sea evaporation from the OAFlux Programme starting in 1958 point to decadal variations with a minimum around 1965-1975, and an overall positive trend of about $10\% \text{ decade}^{-1}$ ($0.06 \text{ mm day}^{-1} \text{ decade}^{-1}$) (Mariotti 2010). Since the mid-1970s, there is a substantial evaporation increase ($0.1\text{--}0.2 \text{ mm day}^{-1} \text{ decade}^{-1}$) with a tendency toward higher rates of increase during the 1990s. Much of the evaporation increase since the mid-1970s has been in early winter, especially in the Ligurian Sea, Adriatic Sea, and southeastern Mediterranean. The evaporation increase has resulted in a rate of increase in freshwater fluxes during 1979-2006 estimated in the range of $0.1\text{--}0.3 \text{ mm day}^{-1} \text{ decade}^{-1}$. Increases in sea surface temperatures have primarily driven these evaporation changes via changes in the surface humidity gradient. Based on OAFlux data, the estimated Mediterranean mean rate of evaporation change in relation to the warming is about $0.7 \text{ mm day}^{-1} \text{ K}^{-1}$ (or $25\% \text{ K}^{-1}$) over the period of 1958-2006. An increase in net Gibraltar water flux to compensate for the overall increase in fresh water loss has been derived (Fenoglio-Marc et al. 2013).

For the land surrounding the Mediterranean Sea, past evapotranspiration changes are regionally and seasonally dependent and largely follow precipitation trends, since soil moisture availability is a primary limiting factor. Increasing soil-moisture limitations seem to have driven recent global evapotranspiration decline and increased drought tendencies over the Mediterranean region (Sheffield and Wood 2008; Vicente-Serrano et al. 2014; Samaniego et al. 2018). Evapotranspiration

estimates from French National Centre for Meteorological Research (CNRM) (Douville et al. 2013) display a tendency for evapotranspiration to increase during winter since the 1970. For the summer there is a progressive decrease (Mariotti et al. 2015).

The future evolution of the Mediterranean Sea physical characteristics is strongly related to the evolution of the air-sea and land-sea exchanges of water and heat. For the Mediterranean Sea, the net surface water loss by the sea is constituted by the combination of the evaporation over the sea, the precipitation over the sea, the river runoff and the Bosphorus Strait net transport. Increase in the net surface water loss by the sea is expected in the future due to a decrease in precipitation and in river runoff and an increase in evaporation (Mariotti et al. 2008, 2015; Sánchez-Gómez et al. 2009; Elguindi et al. 2011; Dubois et al. 2012; Planton et al. 2012; Adloff et al. 2015). Relative to the 20th century, this increase ranges from +8 to +35% for the mid-21st century (2020-2049) and from +20 to +60% at the end of the 21st century (2070-2099) in the medium-range A1B socio-economic scenarios (Planton et al. 2012).

To a first order, CMIP5 projections are largely similar to those based on CMIP3 (Mariotti et al. 2008) and consistent with those based on regional model downscaling (Sánchez-Gómez et al. 2009; Dell'Aquila et al. 2018). By 2071-2098, the Mediterranean Sea evaporation is projected to increase during all seasons and especially in winter (projected annual-mean increase is $0.25 \pm 0.08 \text{ mm day}^{-1}$) (Mariotti et al. 2015). Note that future change in the Nile freshwater inflow remains unknown due to the impossibility so far to accurately model the influence of regional water- and land-use anthropogenic activities on its past and future evolution (Somot et al. 2006; Dubois et al. 2012).

Over land, evapotranspiration projections present mixed changes, with a precipitation-driven increase in winter over Northern areas, and a decrease in summer over many land areas, especially over Spain, western Northern Africa and Turkey. Evapotranspiration increase will be also driven by increase of atmospheric evaporative demand (Vicente-Serrano et al. 2015). These evapotranspiration changes have been linked to a projected northward expansion of the Mediterranean land type (Alessandri et al. 2015) and regional surface vegetation changes (Anav and Mariotti 2011).

Changes in precipitation and evaporation over the Mediterranean Basin will lead to changes in

drought occurrence. Drought can be of different types, such as meteorological, hydrological and agricultural drought, which can often be difficult to separate. Here we focus on meteorological drought, essentially measured by indices of monthly, seasonal up to annual precipitation deficits, such as the precipitation index (PI) or the standardized precipitation index (SPI). The Mediterranean Basin, is impacted by frequent drought episodes due to the strong inter-annual variability of rainfall in this region, and a trend towards drier conditions and increased meteorological drought occurrence after the 1970s over the Mediterranean Basin was found based on analyses of observations (Vicente-Serrano et al. 2011; Hoerling et al. 2012; Spinoni et al. 2015; Caloiero et al. 2018). Due to the pronounced interannual and decadal variability of Mediterranean precipitation, the robustness of this result needs to be confirmed, and may differ for different areas of the Mediterranean.

Concerning projections, since most model simulations indicate a trend towards drier conditions over the Mediterranean, especially in the warm season and over the southern areas, it is expected that the frequency and intensity of meteorological drought will increase under warmer climates. This has been confirmed (*high confidence*) by extensive analyses of precipitation projections with both global and regional climate models (Giorgi and Lionello 2008; Mariotti et al. 2008; Dai 2013; Dubrovský et al. 2014; Spinoni et al. 2015, 2018; Stagge et al. 2015; Quintana-Seguí et al. 2016; Naumann et al. 2018; Lionello and Scarascia 2020).

2.2.6 The cryosphere

2.2.6.1 Observed trends in the cryosphere

After the peak of the "little ice age" (~1,400-1,860 AD, Ivy-Ochs et al. 2009) increasing summer and mean annual air temperature led to a dramatic reduction in the area and volume of glaciers across high mountains of the Mediterranean (Hughes 2018). Short glacier readvances were observed in the 1890s, 1920s, 1970s and 1980s (Zemp et al. 2008). Deglaciation rate generally accelerated in recent decades (Rabaté et al. 2013), although the patterns of glacier retreat were complicated by the sensitivities of glaciers to different climatic regimes (Hughes 2018). As glacier retreats to cirque headwalls, it becomes more dominated by local topoclimatic controls, especially avalanching snow. Nevertheless, a complete loss of glaciers in some low-latitude mountain ranges has already

occurred (Rabaté et al. 2013), accompanied by a shorter duration of seasonal snow cover (Brown and Mote 2009). Several small cirque glaciers existing in the southern Dinaric Alps, Balkan Peninsula, Turkey, Pyrenees, Sierra Nevada and the Apennines disappeared across the 20th century and in the last decades (Hughes 2018). In the Alps, glaciers covered 4,470 km² in 1850, 2,909 km² in the 1970s and 2,270 km² in 2000, meaning a 50% loss from 1850 to 2000 (Zemp et al. 2008). Few very small glaciers still exist in mountains of Montenegro and Albania. Elsewhere, perennial ice and snow patches still survive and attest to how close some Mediterranean mountains are to supporting small glaciers even where the equilibrium line altitude (ELA) is located above the highest peaks (Hughes 2018). ELA raised by about 170 m in the western Alps over the period 1984-2010 (Rabaté et al. 2013) while in the southeastern Alps change in the ELA was in the order of about +250 m between the 1980s and 2010 (Colucci and Žebre 2016).

Temperature increase led to a shift of periglacial processes to higher elevations as well as degradation of mountain permafrost in high mountain environments (Oliva et al. 2018). In the western and central Mediterranean, permanently frozen ground is now rarely found below 2,500 m. Alpine permafrost belt is detected above 2,630 m in northern aspects and 2,800 m in southern ones and in the Pyrenees, above 2,400 m in the Southern Alps, above ~2,350 m on Rila Mountain and ~2,700 m on Mount Olympus and above 2,800-3,400 m in north-eastern Turkey and central Anatolia (Oliva et al. 2018). No permafrost belt is found in the highest mountains in southern Europe (Sierra Nevada) and northern Africa (Atlas) where permanent frozen conditions are only found in the form of isolated patches at the highest elevations at 3,000-3,100 m (Oliva et al. 2016) and 3,800 m (Vieira et al. 2017), respectively. Certain climate conditions (i.e., reduced snow cover) can favour the presence of permafrost patches at relatively low elevations in the Central Apennines or by lithological conditions (i.e., volcanic sediments, karst lithology), as detected in the highest active European volcano (Mt. Etna) at elevations above 2,900 m (Maggi et al. 2018) or in limestone dominated mountains across the Mediterranean (Colucci and Guglielmin 2019).

2.2.6.2 Future conditions in the cryosphere

Mountain glaciers in the Mediterranean region are projected to continue losing mass in the 21st century until complete disappearance of most

mountain glaciers by the end of the century (*very high confidence*). A recent multi-model projection exercise (GlacierMIP, Hock et al. 2019) indicates that relative volume losses by 2100 (average of model runs ± 1 standard deviation) are of the order of 69 $\pm 19\%$ for RCP2.6 and 93 $\pm 10\%$ for RCP8.5. This indicates that, even under scenarios with strong reduction in greenhouse gas emissions, only glaciers at the highest elevation will persist at the end of the 21st century. For mid-century, changes depend far less on the climate scenario, with reductions of the order of 50 $\pm 20\%$ for RCP2.6 and 60 $\pm 20\%$ for RCP8.5.

Projected changes of the mountain snow cover are studied based on climate model experiments, either directly from GCM or RCM output, or following downscaling and the use of snowpack models. Future changes in snow conditions are mostly driven by changes in meteorological drivers. The projections generally do not specifically account for future changes in the deposition rate of light absorbing particles on snow and associated changes in snow albedo. At lower elevation, under the current multi-annual mean rain/snow transition elevation, the water mass of the snow cover is projected to decline by 25% likely range: (10-40%, between the recent past period (1986-2005) and the near future (2031-2050), regardless of the climate scenario). By the end of the 21st century (2081-2100), reductions of up to 80% (likely range 50-90%) are expected under RCP8.5, 50% (likely range from 30-70%) under RCP4.5 and 30% (likely range 10-40%) under RCP2.6 (Beniston et al. 2018; Hanzer et al. 2018; Verfaillie et al. 2018). At higher elevations, projected reductions are smaller (*high confidence*), as temperature increases at higher elevations affect the ablation component of snow mass evolution (in particular, melt and sublimation), rather than the onset and accumulation component. The strong interannual variability of snow conditions is projected to remain a key feature of this cryospheric component throughout the 21st century (*high confidence*).

In the Mediterranean domain, permafrost is only located in the mountains, often patchy and confined to areas of rugged topography, including cliffs. In contrast to glacier and snow cover, climate projections of the ground thermal regime have not been performed in a comprehensive manner, using a cascade of climate models and impact models. Evidence stems from small-scale studies, but all studies point towards increased permafrost thaw in mountain environments, following surface air temperature changes (Marmy et al. 2016; Beniston et al. 2018). Future changes in mountain

permafrost have major implication for natural hazards (slope instabilities).

2.2.7 Ocean hydrology

The Mediterranean Sea can be considered as a laboratory of the global ocean (Lacombe 1990; Béthoux et al. 1999) as it shows many key and interesting oceanic physical processes, such the open-sea deep convection occurring in some areas (Gulf of Lion, South Adriatic, and the Cretan Sea) leading to the formation of cold and salty deep-water masses (Tsimplis et al. 2006; Schroeder et al. 2012). The dominance of evaporation in the Mediterranean Sea and intermediate water formation (in the Rhodes Gyre) leads to an anti-estuarine

thermohaline circulation with a surface layer with the Atlantic Water (AW, comparatively fresh and warm) and a layer with the Levantine Intermediate Water (LIW, very salty and comparatively cold) entering and exiting across the Gibraltar Strait, simultaneously.

In addition, the Mediterranean Sea is surrounded by various and complex topography channeling regional winds (Mistral, Tramontane, Bora, Meltem, Sirocco, Etesians) that define local circulations. The presence of complex coastlines, islands, narrow and shallow straits require adapted observation strategies and high-resolution modeling tools. Further, the Mediterranean Sea is also known to impact the Atlantic Ocean through the Mediter-




Figure 2.12 | A) Local amplification factor of the Mediterranean sea surface warming using 1980-1999 as reference period. Local sea surface warming values are divided by the basin-averaged warming value. Reddish (resp. blueish) colors mean that the area is warming more (resp. less) than the basin-average. The top panel is the ensemble mean of five Med-CORDEX coupled regional climate system models, further averaging warming rates of four 20-year long time periods (2000-2019, 2020-2039, 2040-2059, 2080-2099) and 2 scenarios (RCP4.5, RCP8.5). The bottom is based on the CMEMS observations (2000-2017). **B) Fraction of the Mediterranean Sea surface (in %) experiencing a given sea surface temperature change value (in °C),** compared to the reference period (1980-1999) for various periods (2000-2019 in blue, 2020-2039 in orange, 2040-2059 in purple, 2080-2099 in red) and for the scenarios RCP4.5 and RCP8.5 using the envelope of the results of 5 Med-CORDEX coupled regional climate system models. The change in the CMEMS observations is added for the period 2000-2017 (in black). Information is first aggregated at the yearly scale.

ranean Outflow Waters that flow into the Atlantic at about 1,000 m depth and are considered as a source of salt and heat for the Atlantic Ocean (Artale et al. 2006).

2.2.7.1 Observed change in marine waters

There is increasing evidence that Mediterranean water masses are becoming warmer and deep water masses saltier. This assertion is supported both by direct measurements (Béthoux and Gentili 1999; Rixen et al. 2005; Vargas-Yáñez et al. 2010, 2017) and by numerical simulations (Beuvier et al. 2010; Harzallah et al. 2018; Somot et al. 2018).

Since the 1980s upper layer temperature has increased (Rivetti et al. 2017; Vargas-Yáñez et al. 2017) as well as sea surface temperature (Marullo et al. 2010; Pastor et al. 2018), with acceleration since the 1990s (Macías et al. 2013). Since the beginning of the 1980s, the sea surface warming rate ranges between +0.29 and +0.44°C decade⁻¹ on average over the whole Mediterranean Sea, depending on the studied period and on the reference data sets (Nabat et al. 2014; CEAM 2019; Darmaraki et al. 2019a). In the period 2000–2017 with respect to 1980–1999, all Mediterranean Sea areas show a positive yearly-mean sea surface temperature anomaly of at least +0.2°C. The sea surface warming has not been uniform, but mostly bimodal (*Fig. 2.12b*) with stronger trends in the eastern basin (Adriatic, Aegean, Levantine and North-East Ionian Seas, *Fig. 2.12a*), where some areas warmed by +1.2°C. Very local places in the Levantine Basin have warmed 50% more rapidly than the Mediterranean Sea average whereas a spot in the Ionian Sea has warmed 50% less than the basin average (*Fig. 2.12a*). Note that the climate models currently underestimate the observed sea surface warming (blue and black lines in *Fig. 2.12b*) (Nabat et al. 2014; Sevault et al. 2014; Dell'Aquila et al. 2018).

In the Mediterranean Sea, periods of abnormally warm sea surface, also called "marine heat waves" have become more frequent, more intense, spatially more extended and more severe over the last decades (Oliver et al. 2018; Darmaraki et al. 2019a). To illustrate this trend, the most severe marine heat waves detected since 1982 are 2003, 2012, 2015, and 2017 (Bensoussan et al. 2019). In addition, 14 marine heat waves occurred during the 2008–2017 10-year period whereas only 2 occurred during the 1982–1991 period (Darmaraki et al. 2019a). Contrary to sea surface temperature, a corresponding sea surface salinity evolution has

not been reported (Rixen et al. 2005; Sevault et al. 2014), except for specific locations (Ozer et al. 2017; Vargas-Yáñez et al. 2017).

Long-term trends in the Mediterranean Sea intermediate and deep hydrology have been detected, though they are affected by marked variability on decadal time scales, the Eastern Mediterranean Transient (EMT, Roether et al. 2007), Western Mediterranean Transition (WMT, Schroeder et al. 2016) and BiOS (Gačić et al. 2010) being probably the most known manifestations of this.

Since the mid 1990s the temperature and the salinity of the Levantine Intermediate Water (LIW) have increased by 0.53°C (*Fig. 2.13*, "Sicily Channel" panel) and 0.13 psu, i.e., with rates of 0.024°C yr⁻¹ and 0.006 psu yr⁻¹, respectively (Schroeder et al. 2017). Such trends are at least one order of magnitude greater than those reported for the global ocean intermediate layer (Schroeder et al. 2017). The western basin deep waters are shown to have gradually increased their temperature and salinity since the 1950s (Rohling and Bryden 1992; Béthoux et al. 1998; Rixen et al. 2005; Marty and Chiavérini 2010), with an acceleration after the mid 1980s and an even stronger rate since 2005 due to an abrupt WMT (Marty and Chiavérini 2010; Borghini et al. 2014; Schroeder et al. 2016). Deep-water trends of 0.04°C ±0.001°C decade⁻¹ and 0.015±0.003 psu decade⁻¹ (since 1961) have been reported by comparing time series of deep CTD (conductivity-temperature-depth) casts (Borghini et al. 2014) (*Fig. 2.13*, bottom panel).

Changes in the Mediterranean water mass characteristics have a signature also in the water outflowing from the Mediterranean Sea through the Strait of Gibraltar (Millot et al. 2006; Naranjo et al. 2017). Mooring observations collected since 2004 (*Fig. 2.13*, "Gibraltar Strait" panel) show a positive trend in temperature and salinity of 7.7x10⁻³ °C yr⁻¹ and 0.63x10⁻³ psu yr⁻¹, respectively (von Schuckmann et al. 2018). Since 2012 a noticeable increase of these trends is interpreted as the signal of the WMT (Naranjo et al. 2017). In addition, no significant changes in the strait transports (net exchange: Fenoglio-Marc et al. 2013; Boutov et al. 2014; Soto-Navarro et al. 2015) and surface circulation (Pascual et al. 2014) have been detected.

No significant trends in frequency of dense water formation events have been detected (Beuvier et al. 2010; Houpert et al. 2016; Somot et al. 2018; Dunić et al. 2019), although a strong interannual




Figure 2.13 | Warming stripes in the Intermediate Water (from east to west) and the Deep Water (in the Gulf of Lion). Each stripe refers to a single year and covered periods differ depending on variable. Values have been computed using yearly potential temperature averages in different locations where long-term mooring data are available at different depths. Data from the eastern Ionian come from the HCMR Pylos deep Observatory (Velaoras et al. 2013), and have been downloaded from CMEMS. Data from the Sicily Channel come from the CNR-ISMAR mooring (Schroeder et al. 2017) and can be downloaded from CMEMS. Data from the Gulf of Lion (500 m and 2,300 m) come from the LION Observatory of the MOOSE Network (Houpert et al. 2016) and can be downloaded from SEANOE (Testor et al. 2019). Data from the Gibraltar Strait come from the IEO (Spanish Oceanographic Institute) mooring (von Schuckmann et al. 2018) and can be downloaded from the Copernicus Marine Environment Monitoring Service (CMEMS). Note that colour scales are different for each panel.

variability is reported for each of the dense water formation sites, the northwestern Mediterranean (Marty and Chiavérini 2010; Houpert et al. 2016; Somot et al. 2018; Waldman et al. 2018), the Adriatic Sea (Dunić et al. 2019), and the Aegean Sea (Roether et al. 2007; Beuvier et al. 2010).

2.2.7.2 Future change in marine waters

Air-sea and land-sea exchanges

The future evolution of the Mediterranean Sea physical characteristics is strongly related to the evolution of the air-sea and land-sea exchanges of water and heat. For the Mediterranean Sea, the net surface water loss by the sea is constituted by the combination of the evaporation over the sea, the precipitation over the sea, the river runoff and the Bosphorus Strait net transport. The net surface heat loss by the sea consists of the shortwave radiation, the longwave radiation, the latent heat and the sensible heat fluxes (these variables are assessed in more detail in Sections 2.2.3, 2.2.4 and 2.2.5, as well as in Section 3.1.3.2).

In addition to the changes in radiation, the future evolution of the Mediterranean Sea physical characteristics is strongly related to the evolution of the other air-sea heat fluxes. Under present-climate conditions, the net heat loss by the Mediterranean Sea surface (namely the sum of shortwave radiation, longwave radiation, latent heat and sensible heat fluxes) is positive meaning that sea is losing heat by its surface over a long period of time. In the future, the net heat loss by the sea surface is expected to decrease (Somot et al. 2006, 2008; Dubois et al. 2012; Gualdi et al. 2013; Adloff et al. 2015; Soto-Navarro et al. 2020), because the increase in shortwave, net longwave and sensible heat fluxes will dominate the increase in latent heat loss (Dubois et al. 2012). With respect to the end of the 20th century and based on coupled regional climate models, the decrease in the net heat loss could reach between -1.8 and -5.5 W m⁻² by 2050 following the medium-range A1B scenario (Dubois et al. 2012) and between -2.1 and -6.4 W m⁻² (resp. -1.0 and -3.7 W m⁻²) at the end of the 21st century following high-range RCP8.5 (resp. medium-range RCP4.5) scenarios (Soto-Navarro et al. 2020). This implies that the atmosphere could even start to warm the Mediterranean Sea from the mid-21st century instead of cooling it in the present-day climate according to some models (*medium confidence*). Changes in the Mediterranean Sea surface heat budget depend to a great extent on the socio-econom-

ic scenario chosen: the higher the greenhouse gas emissions, the greater the response of the budget.

To summarize, an increase in the net surface water loss by the sea is expected in the future due to a decrease in precipitation and in river runoff and an increase in evaporation (*Sections 2.2.5 and 3.1.1*). In addition, a decrease in the net surface heat loss by the sea is expected in the future (*Sections 2.2.3 and 2.2.5*) because the increase in shortwave, net longwave and sensible heat will dominate the increase in latent heat loss. In particular, this means that, from the middle of the 21st century, some models predict that the atmosphere could, in average, warm the Mediterranean Sea instead of cooling it in the present-day climate.

Future changes in the wind strength over the sea will likely remain low even at the end of the 21st century in pessimistic scenarios (*Section 2.2.2*).

Sea surface temperature

In future climate change scenarios based on both GCMs and RCMs including the Mediterranean Sea representation, a significant warming of the Mediterranean Sea surface temperature is projected (*very high confidence*) (Somot et al. 2006; Planton et al. 2012; Shaltout and Omstedt 2014; Adloff et al. 2015; Mariotti et al. 2015; Alexander et al. 2018; Darmaraki et al. 2019b). The warming rate depends at the first order on both the temporal horizon and the greenhouse gas emission scenario (*very high confidence*) (Adloff et al. 2015; Mariotti et al. 2015; Darmaraki et al. 2019b). The sea warming will generally remain below that of the air over surrounding land (*high confidence*) due to ocean thermal inertia, probably leading to an increase in land-sea temperature contrast (Somot et al. 2008). With respect to the end of the 20th century, the annual-mean and basin-mean sea surface temperature is expected to increase by 0.6-1.3°C before the mid-21st century and by 2.7-3.8°C (resp. 1.1-2.1°C) at the end of the 21st century period under the pessimistic RCP8.5 (resp. medium RCP4.5) scenario (Darmaraki et al. 2019b). The upper values of those warming ranges are possibly underestimated as higher warming are obtained in CMIP5 GCMs (Mariotti et al. 2015; Darmaraki et al. 2019b).

Future warming will be roughly homogeneous in space (*medium confidence*) with the Balearic Sea, the North Ionian Sea, the Northeast Levantine Sea and the Adriatic Sea identified as potential hotspots of maximum warming (*low confidence*) (*Fig. 2.12a*)

(Adloff et al. 2015; Darmaraki et al. 2019b; Soto-Navarro et al. 2020). This hotspot pattern however does not match well with the observed warming pattern (*Fig. 2.12a*), illustrating that climate change related evolution is likely still hidden by natural variability. Spatially, the future warming of the sea surface is bimodal as it has been in the past (*Fig. 2.12b*). For the near-future (2020-2039) with respect to the end of the 20th century, local annual-mean sea surface temperature change is everywhere positive and can reach locally +1.6°C at maximum whatever the scenarios whereas at the end of the 21st century, the local annual-mean warming spreads from 2 to 5°C for scenario RCP8.5 (resp. from 0.5-3.0°C for RCP4.5).

Warming is not projected to be constant all year round. Stronger warming is expected in summer and weaker warming in winter (*medium confidence*), resulting in substantial increase in warm extremes and a decrease in cold extremes (Alexander et al. 2018). As an illustration, under RCP8.5, maximum monthly-mean sea surface temperature anomalies could reach +3°C over 2040-2059 and more than +5°C over 2080-2099 (median of the CMIP5 models) averaged over the Mediterranean Sea (Alexander et al. 2018). In addition, from the period 2040-2069, the 30-year mean sea surface temperature will always be warmer than the warmest year during the period 1976-2005 (*medium confidence*). This will already be the case in about 50% of the years for the 2010-2039 period (Alexander et al. 2018).

Marine heat waves will very likely increase in spatial coverage, become longer, more intense and more severe than today (*medium confidence*). The intensity of this evolution strongly depends on the temporal horizon and on the socio-economic scenario (Frölicher et al. 2018; Darmaraki et al. 2019b). By 2021-2050, it is expected that marine heat wave frequency increases by a factor 1.5, duration by 2.4-2.7, mean intensity by 1.5 and severity by 5-7 with values largely independent from the socio-economic scenarios (Darmaraki et al. 2019b). By 2100, models project at least one long-lasting marine heat wave occurring every year under RCP8.5 up to 3 months longer, and about 4 times more intense and 42 times more severe than today's events. Their occurrence is expected between June and October, affecting at peak, the entire Mediterranean Basin (Darmaraki et al. 2019b). Under a RCP8.5 scenario, the 2003 marine heatwaves may become a normal event for the period 2021-2050 and a weak event at the end of the 21st century (*medium confidence*) (Darmaraki et al. 2019b).

The warm extreme sea surface temperature changes at the end of the 21st century is likely due to a combination of three factors: a mean sea surface warming, an amplification of the seasonal cycle and an increase in the interannual and day-to-day variability [Alexander et al. 2018; Darmaraki et al. 2019b].

Sea surface salinity

The future evolution of sea surface salinity of the Mediterranean Sea remains largely uncertain as its sign of change [Adloff et al. 2015; Soto-Navarro et al. 2020]. Any change will likely be spatially and temporally inhomogeneous (*medium confidence*) due to the primary role of the river and near-Atlantic freshwater inputs [Adloff et al. 2015; Soto-Navarro et al. 2020]. For the end of the 21st century, basin-scale surface salinity anomalies range from -0.18 to +0.16 psu (resp. -0.25 to 0.25 psu) for the pessimistic RCP8.5 (resp. RCP4.5) scenario [Soto-Navarro et al. 2020]. However, a surface salinity increase in the eastern Mediterranean Basin is more likely than not whereas the western basin may see an increase or a decrease in its surface salinity [Adloff et al. 2015; Soto-Navarro et al. 2020].

Surface circulation and exchanges across straits

Change in sea surface circulation has not been deeply assessed yet in the literature [Adloff et al. 2015; Macías et al. 2018], despite their strong capacity to locally modulate the future sea surface temperature and salinity anomalies. In particular, it is likely that the surface circulation changes affect the local sea surface warming hotspots listed above. Noticeable surface circulation changes have been reported for the end of the 21st century in the Balearic Sea and in the North Ionian Sea independently from the scenario choice [Adloff et al. 2015].

At the strait of Gibraltar, the net heat transport towards the Mediterranean Sea will likely increase due to near-Atlantic warming as well as the net mass transport due to increased sea surface water deficit [Somot et al. 2006; Marcos and Tsimplis 2008; Carillo et al. 2012; Adloff et al. 2015; Soto-Navarro et al. 2020]. The future evolution of the net salt transport at the strait is unclear, because it depends on the salinity change in the near-Atlantic Ocean surface layer entering the Mediterranean Sea [Marcos and Tsimplis 2008; Adloff et al. 2015; Soto-Navarro et al. 2020]. This means that it is currently unclear if the salt transport from the

Atlantic will increase or decrease in the future leading to large uncertainty for the salinity change in the Mediterranean Sea.

Deep water characteristics

Due to the contrasting effects of increase in sea surface temperature and salinity, the future evolution of the sea surface density is uncertain. Generally, scenarios with strong greenhouse gas concentration increase project a decrease in surface density associated to an increase in vertical stratification of the water column. Increase in density (thus a decrease in stratification) is still possible in scenarios with low level of warming [Adloff et al. 2015]. Due to its active thermohaline circulation, the surface climate change signal may be propagated efficiently towards the deeper layers of the Mediterranean Sea [Somot et al. 2006; Carillo et al. 2012] and lead to larger deep warming rates than in other oceans in the world.

The warming and salting rates of the deep layers is very uncertain as it depends on various factors such as the surface signal, the intensity of the present and future Mediterranean thermohaline circulation (MTHC). This means in particular that the socioeconomic scenario is not the main source of uncertainty in future changes of the deep layers [Adloff et al. 2015]. At the end of the 21st century, water masses deeper than 600 m may warm between +0.03 and +1.38°C, and their salinity may increase or decrease with a large uncertainty range, depending on the model (-0.05; +0.51) psu [Adloff et al. 2015; Soto-Navarro et al. 2020].

All published studies agree on a long-term weakening of the open-sea deep convection, the winter deepwater formation and the related branch of the thermohaline circulation in the western Mediterranean Sea in high emission scenarios [Thorpe and Bigg 2000; Somot et al. 2006; Adloff et al. 2015; Soto-Navarro et al. 2020]. However, natural variability may lead to increase in deep water formation with respect to today's situation during short periods in the future [Macías et al. 2018]. For the end of the 21st century and the A2 scenario, decrease of the maximum mixed layer depth reached in the northwestern Mediterranean Sea reach between -17% and -82% depending on the model choice [Somot et al. 2006; Adloff et al. 2015]. The picture in the eastern Mediterranean Sea is more contrasted with weakening in some simulations but enhanced convection and thermohaline circulation in others [Somot et al. 2006; Adloff et al. 2015; Soto-Navarro et al. 2020]. Some simulations (but not all) project that EMT-like

situation may become the new normal situation for the eastern basin (Adloff et al. 2015).

2.2.8 Sea level, storm surges and wave heights

A particular characteristic of the Mediterranean Sea in terms of sea level variability is that it is a semi-enclosed domain linked to the global ocean through the Strait of Gibraltar. This implies that changes in the nearby Atlantic are quickly transferred into the Mediterranean as a basin-wide barotropic signal. At the same time, basin-wide sea level anomalies caused by local forcing (e.g., thermal expansion, evaporation) tend to be transferred to the global ocean in a way that the Mediterranean is in balance with the nearby Atlantic. As a consequence, the Mediterranean basin-wide variations, especially at low frequencies, closely follow the variations in the nearby Atlantic (Calafat et al. 2012; Adloff et al. 2018). The exception to this is the part of the variability related to changes in the atmospheric mechanical forcing (i.e., wind and atmospheric pressure), which can produce Mediterranean-Atlantic differences of few cm even at multidecadal time scales (Menemenlis et al. 2007; Jordà et al. 2012).

The Mediterranean Sea is a microtidal region, with tidal range mostly below 15 cm and relatively low sea levels with 50-year return values below 60 cm over most of the basin (Marcos et al. 2009). These values are small compared to other European Seas. The two exceptions are the North Adriatic and the Gulf of Gabes, where storm surge levels are estimated to be several times per year above 50 cm, with 5-year return values around 90 cm and 70 cm, respectively (Conte and Lionello 2013).

The wave climate in the region is milder than in the Atlantic with smaller mean wave heights (1-1.5 m) and shorter periods (5-6 s) and presents an important spatial variability due to the complex orography and coastline surrounding the basin (Menéndez et al., 2014). Its variability is connected to NAO and the Indian Monsoon index (Lionello and Sanna 2005) and other northern hemisphere teleconnection indices, particularly the East Atlantic Pattern (Lionello and Galati 2008). Annual maxima along the coastlines are largest (above 5 meters) at the northwestern coast of Africa, but high values well above 4 meters occur in several parts (Lionello et al. 2017).

2.2.8.1 Observed change in sea-level

During the 20th century, coastal tide gauges around the Mediterranean have recorded a rise

in the mean sea level. Once tide gauge data have been corrected for the vertical land motion, the sea level trend is very consistent among sites being $\sim 1.4 \text{ mm yr}^{-1}$ (Wöppelmann and Marcos 2012). This trend is superimposed on interannual and decadal variability that can temporarily mask the sea level rise. The clearest example is the period 1960-1980 during which Mediterranean sea level showed a decreasing trend because a higher than usual atmospheric pressure (Tsimpis et al. 2005). After that period, the atmospheric pressure returned to the typical values and sea level continued to follow the global evolution. For the more recent period, in which sea level has been monitored by satellite altimetry (1993-2018), Mediterranean sea level trend has increased up to $2.8 \pm 0.1 \text{ mm yr}^{-1}$, consistent with global sea level trend ($3.1 \pm 0.4 \text{ mm yr}^{-1}$) (Cazenave and WCRP Global Sea Level Budget Group 2018). The rise at global scale is mainly the result of a combination of water thermal expansion and land-based ice melting. During the 20th century both factors contributed equally, although during the last decades, glacier melt is dominating (Cazenave and WCRP Global Sea Level Budget Group 2018).

Analyses of tide gauge data have revealed an increase in the magnitude and duration of the extreme sea level events in the region during the last decades, caused by the rise in the relative mean sea level (for the northern Adriatic Sea: Lionello et al. 2012b; Marcos et al. 2015). In general, wave observational records are too short for assessing multidecadal trends, with the exception of the Northern Adriatic Sea, where one among the worldwide longest instrumental time series (1979 to present) shows an increase in the number of storms, but a decrease of the extreme wave heights (Pomaro et al. 2017).

2.2.8.2 Future sea-levels

The modeling of Mediterranean mean sea level future variations is not straightforward. With their coarse spatial resolution, present-day global climate models (GCMs) are not able to reproduce the regional processes in the basin, although, they are better suited to represent the connection to the global ocean (Calafat et al. 2012). Conversely, regional climate models (RCMs) can capture part of the regional variability but are usually not designed to reproduce the connection with the global ocean, and thus missing a key part of the variability (Adloff et al. 2018). Therefore, sea level rise projections solely based on RCMs have missed that component and only should be considered for the regional patterns, which can cause local

spatial deviations from the basin average by up to +10 cm (Carillo et al. 2012; Adloff et al. 2015, 2018). In conclusion, accounting for all components shows that the Mediterranean sea level rise will be close to the northeastern Atlantic, where future sea level will be similar (difference lower than 5%) to the global mean sea level because regional differences produced by changes in the circulation and mass redistribution almost compensate each other (Slangen et al. 2017). This leads to estimate that the basin mean sea level will likely be 37-90 cm higher than at the end of the 20th century, with a small probability to be above 110 cm. Main contributions to basin-average sea level changes are coming from terrestrial ice melting and the northeastern Atlantic dynamics (Jordà et al. 2020, Fig. 2.14). A different computation based on the sea level projections in the SROCC (Oppenheimer et al. 2019) and accounting for the uncertainty calculation method of the AR5 (Church et al. 2013), confirms that the likely range of the Mediterranean Sea level will be approximately in the range from

a function of time and of the emission scenario, reaching a value in the range from -5% and -10% at the end of the 21st century in the RCP8.5 scenario (Lionello et al. 2017). In any case future sea level rise will become the dominant factor and it will lead to an increase frequency and intensity level of coastal floods (Lionello et al. 2017; Vousdoukas et al. 2017).

Regarding future changes in waves, they will be determined by changes in the wind field over the Mediterranean Sea (Section 2.2.2). Published studies point towards a generalized reduction of the mean significant wave height field over a large fraction of the Mediterranean Sea, especially in winter (Lionello et al. 2008, 2017; Perez et al. 2015). Similarly, the wave extremes are expected to decrease in number and intensity, although there is no consensus whether very large extreme events, associated with very strong winds, would also decrease (Gaertner et al. 2007; Romero et al. 2017; Romero and Emanuel 2017).




Figure 2.14 | Projected Mediterranean sea level rise averaged in (2080-2099) with respect to present climate (1980-1999) under scenario RCP8.5. Results based on CMIP5 and Med-CORDEX outputs for the dynamical components and Slangen et al. (2017) for other components. [a] Sum of all contributors [b] Range of projected values for the different contributors: NE Atlantic dynamics (Atl Dyn), Mediterranean dynamics (Med Dyn), Atmospheric Loading (AL), Glacial Isostatic Adjustment (GIA), Ground Water (GW) and Terrestrial Ice Melting (TIM) and Total.

20 to 110 cm higher (depending on scenario) at the end of the 21st century than at the end of the 20th century (Le Cozannet et al. 2019; Thiéblemont et al. 2019).

RCMs and GCMs do not model extreme sea level events and specific 2D simulations forced by high frequency atmospheric forcing are needed. Published studies point towards a reduction on the average number of positive surges throughout the 21st century (Marcos et al. 2011; Conte and Lionello 2013; Lionello et al. 2017). Overall, the results indicate small progressive reduction in comparison with their present-day magnitude as

2.2.9 Acidification of the Mediterranean Sea

Human activities are responsible for an increase in atmospheric CO₂ since the beginning of the industrial era. The input of anthropogenic carbon in the Mediterranean is caused by the flux at the air-sea interface, but also results for the Mediterranean Sea, from exchange with the Atlantic Ocean across the Strait of Gibraltar. Approximately 30% of anthropogenic carbon is absorbed by the oceans (Sabine et al. 2004) and leads to decrease of pH in ocean water masses. The Mediterranean Sea is able to absorb relatively more anthropogenic CO₂ per unit area than the

	RECENT CHANGE	PROJECTED CHANGES RCP2.6										PROJECTED CHANGES RCP8.5																								
		20-YEAR PERIODS					20-YEAR PERIODS					20-YEAR PERIODS					20-YEAR PERIODS																			
	2000-2019	NEAR-FUTURE (2020-2039)			MID-CENTURY (2040-2059)			END OF THE 21ST CENTURY (2080-2099)			NEAR-FUTURE (2020-2039)			MID-CENTURY (2040-2059)			END OF THE 21ST CENTURY (2080-2099)																			
CHANGE IN SURFACE TEMPERATURE (°C, MEAN, VALUE, LAND-ONLY)																																				
	BASED ON CRU			BASED ON EURO_CORDEX 12 KM ENSEMBLE																																
		ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA	ANN	DJF	JJA														
MED	0.8	0.41	1	1.1	1.1	1.2	1.3	1.2	1.5	1.3	1.4	1.3	1.3	1.3	1.5	2.2	2	2.5	4.6	4.2	5.3															
WEST MAGHREB	0.5	0.1	0.8	1.3	1.1	1.3	1.4	1.1	1.6	1.3	1.3	1.4	1.4	1.2	1.6	2.4	2.1	2.7	5	4.2	4.6															
EAST MAGHREB	0.6	0.3	0.6	1.2	1	1.2	1.4	1.1	1.6	1.4	1.3	1.4	1.4	1.2	1.6	2.3	2	2.6	4.6	4	5.3															
LEVANT	0.9	0.7	1	1.2	1	1.2	1.3	1.3	1.4	1.3	1.3	1.3	1.3	1.3	1.4	2.1	1.9	2.4	4.3	3.9	4.8															
ANATOLIA	0.9	0.4	1.2	1.2	1.2	1.3	1.4	1.4	1.6	1.5	1.6	1.5	1.4	1/4	1.6	2.3	2.1	2.6	4.6	4.3	5.3															
CENTRAL MED	0.8	0.3	1	1.1	1	1.3	1.3	1.2	1.4	1.2	1.4	1.2	1.3	1.3	1.5	2.1	1.9	2.5	4.3	3.9	5.1															
IBERIA	0.5	0.2	0.8	1	0.9	1.2	1.1	0.9	1.3	1	1	1.1	1.1	1	1.3	2	1.7	2.4	4.3	3.6	5.3															
FRANCE	0.7	0.4	0.7	0.9	0.9	1	1.1	0.9	1.1	0.9	1.1	0.8	1	1	1.1	1.8	1.7	2	3.9	3.6	4.7															
ALPS	0.8	0.5	1	1.1	1.1	1.1	1.3	1.2	1.3	1.1	1.4	1	1.2	1.3	1.3	2.1	2.1	2.2	4.5	4.5	5															
BALKANS	1.1	0.8	1.4	1.1	1.1	1.1	1.3	1.4	1.3	1.2	1.7	1	1.3	1.4	1.4	2.1	2.3	2.2	4.4	4.6	4.8															
CHANGE IN SURFACE TEMPERATURE (°C, MEAN, VALUE, LAND-ONLY)																																				
	BASED ON CRU			BASED ON EURO_CORDEX 12 KM ENSEMBLE																																
		ANN	WET	DRY	ANN	WET	DRY	ANN	WET	DRY	ANN	WET	DRY	ANN	WET	DRY	ANN	WET	DRY	ANN	WET	DRY														
MED	1.3	2.5	-0.4	0.8	2.3	-1	1	2.3	-0.6*	3.9	6	1.1*	0	1.5	-2.1*	-1.9	0	-4.5	-6.8	-2	-13.3															
WEST MAGHREB	2.7	1.3	6.2	-6.7	-7.8	-5.2	-6.2	-5.7	-6.9	-1.7*	0.8	-4.9	-11.2	-9.4	-13.7	-16.8	-20.3	-11.8	-31.2	-33	-28.7															
EAST MAGHREB	-7.6	-11.1	5.9	-10.1	-11	-8.7	-5.4	-3.4*	-8.4	-7.8	-8.2	-7.2	-9.7	-8.6	-11.4*	-11.8	-11.7	-11.9	-23.6	-24.1	-22.7															
LEVANT	-1.6	-3.5	12.2	-5.8	-5.1	-8.4	-5.8	-4.7	-10.1	-5.8	-4.3	-12	-4.6	-5.1	-2.4*	-10.7	-10	-13.5	-23.5	-23.6	-23.1															
ANATOLIA	5.7	3.7	10	0.5*	3.1*	-4.1	-0.1*	2*	-3.9*	2.6	5.3	-2.4	0.2*	2.2	-4.1	-2.4	-0.2	-6.5	-8.7	-4.4	-16.4															
CENTRAL MED	2.8	1.3	6.3	0.1*	1.8	-2.4	0.8*	2.1*	-1.2*	3.8	6.3	-0.8	-0.5*	1.5*	-4*	-3.3	-0.7	-7.6	-9.7	-4.7	-18.1															
IBERIA	4.9	10	-4.1	-0.1	2.3*	-3.6	0.3*	2.6	-3.2*	5	10.1	-2.8*	-1.5*	0.5*	-4.9*	-6	-2.2	-12.4	-15.1	-7.5	-27.7															
FRANCE	-1.8	0.7	-4.4	2.5	4.3	0.3*	2.4	4.2	0.2*	4.1	5.2	2.8	1.6	3.1	-0.3*	0.1	4.4	-5.2	-2.5*	6.7	-13.9															
ALPS	-0.5	2.2	-2.8	3.5	6.3	0.7*	4.6	6	3.2	6.2	7.1	5.3	2.9	5	0.9*	3.1	6.6	-0.4*	1.5	9.2	-6.3*															
BALKANS	2.5	6.7	-0.4	5.3	8.8	2.4	4.5	6	3.3	8.3	11.8	5.4	3.9	6.6	1.5	4.7	8.1	1.6*	4.9	14.4	-3.7*															
CHANGE IN SURFACE SOLAR RADIATION (W/M², 90% INTERVAL BASED ON CMIP5 SIMULATIONS, LAND+SEA)																																				
	ANN			ANN			ANN			ANN			ANN			ANN			ANN																	
MED	0.3 ; 5.2			0.8 ; 8.6			0.8 ; 10.7			0.5 ; 12.6			0.6 ; 7.7			1.0 ; 10.5			0.6 ; 12.4																	
CHANGE IN CLOUD COVER (% , 90% INTERVAL BASED ON CMIP5 SIMULATIONS, LAND+SEA)																																				
MED	-2.1 ; 0.2			-3.6 ; 0.2			-4.4 ; 0.5			-4.3 ; 0.5			-3.3 ; 0.0			-4.9 ; -0.2			-7.4 ; -0.8																	
CHANGE IN AOD (-, 90% INTERVAL BASED ON CMIP5 SIMULATIONS, LAND+SEA)																																				
MED	-0.06 ; -0.02			-0.12 ; -0.04			-0.15 ; -0.04			-0.18 ; -0.05			-0.10 ; -0.03			-0.12 ; -0.04			-0.14 ; -0.04																	
CHANGE IN SURFACE TEMPERATURE (°C)																																				
	CMEMS (2000-2017)			TOTAL RANGE BASED ON MED-CORDEX RCSM																																
MEDSEA	0.74 (+0.23 ; +0.87)			X			X			X			0.59 ; 1.18			1.30 ; 2.07			2.86 ; 4.10																	
CHANGE IN SEA LEVEL (CM, BASED ON BLENDED MULTIPLE DATABASE, SEE THE TEXT, CM, VERY LIKELY RANGE)																																				
MEDSEA										[+42, +82]									[+70, +110]																	

Table 2.1 | Climate change as a function of time period and Representative Concentration Pathway for the Land sub-regions in Fig. 2.1 and the whole Mediterranean Sea area. All changes are with respect to the (1980-1999) reference period. For temperature and precipitation, recent changes are based on the Climate Research Unit CRU-TS, future changes are based on the Euro-Cordex regional model simulations. For surface solar radiation, cloud cover and aerosol optical depth values are based on CMIP5 global simulations. For sea surface temperature, recent changes are based on CMEMS observations (2000-2017), future changes on the Med-Cordex regional simulations. For sea level rise, future changes are based on blended multiple databases (see text).

global ocean for two reasons. It is more alkaline, thus giving it greater chemical capacity to take up anthropogenic CO₂, and deep waters are ventilated on shorter timescales (Schneider et al. 2010), thus allowing rapid penetration of CO₂ in its interior.

2.2.9.1 Observed change in acidity

Concerning the past trends of anthropogenic carbon absorption by the Mediterranean Sea, the presence of natural CO₂ prevents to determine it from direct measurements in the water column. Estimations of anthropogenic CO₂ from data-based approaches are limited and with large uncertainties that provides concentrations that disagree by more than a factor of two in the Mediterranean Sea (Schneider et al. 2010; Touratier and Goyet 2011). These large differences further result in even opposing estimates for the net CO₂ transport across the Strait of Gibraltar.

In this context, the modeling approach using high-resolution regional model provided some insights on the information resulting from the data based-estimates and quantification of processes responsible of anthropogenic CO₂ storage and acidification of the Mediterranean Sea (Palmiéri et al. 2015). 25% of total anthropogenic carbon inventory in the Mediterranean Sea is due to net exchange at the Strait of Gibraltar, while the remaining 75% is from the air-sea flux. It confirms that the 10% higher mean total alkalinity of the Mediterranean Sea is responsible for a 10%

increase in anthropogenic carbon inventory. The higher alkalinity acts to neutralize acidification and simulated average surface pH change is thus similar for the Mediterranean Sea and the global ocean (-0.08 units), with deep waters exhibiting a larger anthropogenic change in pH than typical global ocean deep waters because ventilation times are faster (between -0.005 and -0.06 units) (Palmiéri et al. 2015).

2.2.9.2 Future change in acidity

The published literature concerning specifically the future acidification of the Mediterranean Sea is quite limited leading to low confidence in the assessment. Based on thermodynamic equations of the CO₂/carbonate system chemical equilibrium in seawater, Goyet et al. (2016) calculated the variation of pH (ΔpH) as a function of theoretical anthropogenic CO₂ concentrations. Under the most optimistic SRES scenario, the results indicate that in 2100, pH may decrease down to 0.245 in the western basin and down to 0.242 in the eastern basin (compared to the pre-industrial pH). Whereas for the most pessimistic SRES results for 2100 project a pH decrease down to 0.462 and 0.457, for the western and for the eastern basins, respectively (Goyet et al. 2016). However, these estimates do not consider that the warming of seawater will decrease exchanges across ocean-atmosphere interface and penetration of anthropogenic Jungcarbon, thus they tend to overestimate future acidification.

2.3 Pollution

2.3.1 Introduction

The 2030 Agenda for Sustainable Development pledges to "ensure that all human beings can enjoy prosperous and fulfilling lives and that economic, social and technological progress occurs in harmony with nature" (UN 2015). Pollution puts at risk the possibility of achieving these outcomes and hence health and well-being. Pollution touches all parts of the planet. It is affecting our health through the food we eat, the water we drink and the air we breathe. Approximately 19 million premature deaths are estimated to occur annually as a result of the way we use natural resources and impact the environment to support global production and consumption. By definition, "pollutant" shall mean any substance that is introduced into the environment that has undesired

effects, or adversely affects the usefulness of a natural resource (air, soil, water and ecosystems). Pollutants can take many forms: (i) physical, substances that are not necessarily involved in chemical or biological reactions, e.g., dust; (ii) chemical, substances that are involved in chemical reactions, e.g., pesticides; and (iii) biological, e.g., bacteria.

2.3.2 Physical pollutants

Particulate matter (PM) levels have been monitored during the past decades, mainly because of their effects on health and climate. Aerosols into the atmosphere arise from a variety of anthropogenic activities (transport, industry, biomass burning, etc.) as well as natural sources (volcanic eruptions, sea salt, soil dust suspension, natural forest fires,

etc.) (Seinfeld and Pandis 2006). Both sources result in direct emission of PM (primary PM) and emission of gaseous aerosol precursors (leading to secondary PM). A number of epidemiological studies have examined the impact of PM on human health, expressed as increased mortality and morbidity varying according to the physical (size, shape, etc.) and chemical (composition) characteristics of PM (Van Dingenen et al. 2004).

The PM impact on climate is primarily a cooling effect due to increased scattering to space as the atmospheric aerosol burden increases. The overall cooling by aerosols might be equivalent to a radiative forcing of up to 2.5 W m^{-2} , counterbalancing global warming by greenhouse gases (Gillett et al. 2013; Knutson et al. 2013). More important than this direct effect may be the indirect effect that aerosols have on climate, acting as cloud condensation nuclei (CCN) (Levin et al. 2003; Gerasopoulos et al. 2006). Moreover, the particles have a large effect in reducing visibility as well as play a significant role in the deterioration of monuments and buildings (Gerasopoulos et al. 2006). Several studies conducted over the Mediterranean Basin revealed a distinct spatial inhomogeneity (Gerasopoulos et al. 2006), with PM levels increasing from north to south and west to east of the basin (Querol et al. 2009), and distinct sources. PM analysis through the years allowed the identification and classification of PM episodes as follows: (i) local urban PM pollution events (mostly in the cold season), (ii) regional PM pollution episodes (warm season) and (iii) dust outbreaks (Rodríguez et al. 2003).

2.3.2.1 Particulate matter (PM) levels and sources

Several factors favor the occurrence of high PM concentrations in the Mediterranean Basin. First, the abrupt topography, coupled with the characteristic synoptic scale patterns, results in low mean wind speeds that hinder the air mass renovations and favor the accumulation of PM in the surrounds of emission regions – leading to the so-called Atlantic/Northern clean air advects events (Rodríguez et al. 2007). Second, the low precipitation in the Mediterranean Basin favors the long residence time of PM in the atmosphere, leading to higher background PM levels (Rodríguez et al. 2007; Querol et al. 2009). The joint influence of low precipitation rates and traffic-forced resuspension of road (which is strongly enhanced by the dust accumulation in streets and roads), construction and demolition dust promoted the local urban episodes in the Mediterranean Basin

(Rodríguez et al. 2007; Talbi et al. 2018). This factor, combined with the high percentage of water coverage of the area, especially in the East region of the basin, has a great contribution of the sea-salt aerosols to the PM levels and composition (Im 2013).

PM observations from monitoring networks, in the period 2007-2009, were analyzed in order to characterize particulate pollution and its health effects across Mediterranean countries (Karanasiou et al. 2014). It was concluded that the average concentrations for PM across the Mediterranean Basin are within the range of annual means typical of European sites and according to the monitoring site characteristics (traffic and urban background sites) (Querol et al. 2004; Putaud et al. 2010). The regional patterns mentioned in previous studies were highlighted, with higher PM concentrations in Italian and Greek cities, and lower levels in the Western Mediterranean (Barcelona, Marseille, Madrid, Huelva). PM10 levels at the traffic sites showed a quite similar variation. In Turin, as in the other cities of the Po valley (Bologna, Milan, Parma, Modena and Reggio Emilia), the combination of stagnant air conditions with high emissions and high population density is the main cause of very strong pollution episodes (Cyrys et al. 2012). Similarly, the air pollution problems in Athens and Thessaloniki are the result of the high population density and the accumulation of air pollutants over the city, due to topography (basin surrounded by mountains), narrow and deep street canyons and adverse meteorological conditions (Karanasiou et al. 2009; Kassomenos et al. 2011). Thermal inversions, followed by accumulation of air pollutants in the lower layers of the atmosphere are also very common in different locations like Athens (Karanasiou et al. 2014) or Beirut (Saliba et al. 2006), increasing the evening concentrations of ambient PM10.

In most countries of the southern Mediterranean, air pollution is not sufficiently monitored (Naidja et al. 2018). Emission inventories are less precise than that available in the northern Mediterranean since they are generally based on surveys and questionnaires. Because of that, local scientific articles were relatively scarce and hard to find. However, most of the available studies show that PM concentrations in this Mediterranean region are much higher than the limit values given in WHO guidelines (Naidja et al. 2018). Emissions from road traffic, resuspension of road dust, especially on unpaved roads, and natural contributions have been found to be an important source of fine particles and play a key role on the

observed levels and exceedances (Mahmoud et al. 2008; Abderrahim et al. 2016; Naidja et al. 2018). Cairo (Egypt) is an example of a city where road traffic emissions are hugely important in PM10 concentration, and, according to WHO is ranked in the 33rd position on the list of the most polluted cities by PM10 (Mahmoud et al. 2008; Lowenthal et al. 2014; Naidja et al. 2018).

Since it is expected that the majority of the Mediterranean population will continue to live in cities, especially in the eastern and southern part of the basin, with a tendency to growth, higher anthropogenic pressure in a context of climate change will occur (Rafael et al. 2015; Naidja et al. 2018). Most PM exceedances were registered in regional background sites (Escudero et al. 2007), with more than 70% of them being attributed to dust outbreaks (Escudero et al. 2007; Mitsakou et al. 2008). Compared with the central and northern Europe, the occurrence of higher PM concentrations associated with dust outbreaks is higher in the Mediterranean Basin (Rodríguez et al. 2007), and are more frequent and more intense in the central and eastern than those in the western Mediterranean Basin. These episodes have been studied on an 11-yr period (2001–2011) (Pey et al. 2013). Dust outbreaks are very frequent in the southern Mediterranean, where they occur more than 30% of the days, while in northern Mediterranean this value is below 20%. The central Mediterranean appears as a transitional area, with a decreasing south to north gradient of dust outbreaks, with slightly higher frequency of dust episodes in its south, when compared to west and east sides of the basin, for similar latitudinal positions (Pey et al. 2013).

Regarding intensity characteristics and seasonality patterns, significantly high contributions are common in autumn-spring in the eastern Mediterrane-

an, with occurrence of many severe episodes (daily dust averages over $100 \mu\text{g m}^{-3}$ in PM10) throughout the year. However, in the western Mediterranean a clear summer prevalence is noticed, with low occurrence of severe episodes; and no seasonal trend is detected in the central region, with moderate-intensity episodes (Pey et al. 2013). The contribution of dust outbreaks to PM concentrations reveals a downward trend in the period between 2006 and 2011, a period in which there was also a decrease of the NAO index for the summer period. Therefore, it can be concluded that a sharp change in the atmospheric circulation have affected the number of dust episodes and, consequently, the annual dust inflows to PM10 (Fig. 2.15) observed in the Mediterranean Basin (Pey et al. 2013).

The low PM2.5/PM10 ratio (approximately 0.25) in the eastern Mediterranean region also indicates that the particle size distribution has a large contribution of coarse particles which are either affected by a background level of naturally occurring dust (dust outbreaks from the Saharan Desert and sea salt particles from the Mediterranean Sea itself) or that the region is characterized by high levels of primary coarse PM emissions (Koçak et al. 2007b, 2007a). Even though the PM2.5/PM10 ratio showed seasonal variations, the values remained lower than 0.5 in most cases (Koçak et al. 2007a; Asaf et al. 2008), a value that is least two times lower than those of the western Mediterranean (Saliba and Massoud 2010).

2.3.2.2 Particulate matter (PM) chemical profiles

Regarding the chemical composition of PM, different species can be found such as carbonaceous compounds, inorganic ions and metals (Galindo et al. 2018). Although they are present at extremely low levels, some components such as trace metals are relevant in air quality studies because of



Figure 2.15 | PM10 concentration above the annual limit value of $40 \mu\text{g}\cdot\text{m}^{-3}$ (based on EU Directive 2008/50/CE).

their toxicity and environmental persistence (Roig et al. 2013). Recent clinical and toxicological studies demonstrate the link between exposure to airborne metal through inhalation and pulmonary and cardiovascular effects, genotoxic and carcinogenic outcomes and increased daily mortality (Gottipolu et al. 2008; Lippmann and Chen 2009; Tchounwou et al. 2012).

Cooling metal concentrations are considered as good tracers of specific pollution sources, both natural and anthropogenic (Arhami et al. 2017; Diapouli et al. 2017). The main natural sources include wind-blown dust and sea-spray (Chen et al. 2008; Engelbrecht and Jayanty 2013) including elements such as calcium, aluminum, iron, potassium, sodium and magnesium. Desert dust contribute to PM composition and have a high influence on climate in the North as well as in the South of the Mediterranean (Kchih et al. 2015; Kaskaoutis et al. 2019). Specific meteorological circulations and natural sources like the Mediterranean Sea and the proximity of Sahara create specific patterns of aerosol concentrations that could influence not only the particulate concentrations through Europe but also the global climate due to the transport of dust from the Sahara (Ganor et al. 2010).

Regarding anthropogenic activities, exhaust and non-exhaust vehicle emissions, coal combustion and a variety of industrial processes, like metal works and smelters, are the major sources of heavy metals such as zinc, copper, nickel or chromium (Thorpe and Harrison 2008; Pant and Harrison 2013). In the last decades, emissions of some heavy metals in Mediterranean Basin have dropped significantly, in particular from industrial facilities due to improvement of abatement techniques (Dayan et al. 2017). In the case of lead, a drastic reduction in ambient concentrations has been observed since the introduction of unleaded gasoline (Cho et al. 2011; Salvador et al. 2012). The influence of traffic and dust outbreak intrusions on PM levels and metal content have been studied (Galindo et al. 2018), showing that the PM coarse fraction was affected more by variations in traffic intensity than the submicron fraction: the highest decreases during the weekends due to the reduction in traffic induced resuspension. That dust outbreaks had a greater impact on the levels of other metals such as titanium and lead, significantly affecting their seasonal variability. High concentrations of vanadium and nickel compared with the values found at larger urban areas were observed. This could be attributed to a significant contribution from soils, dust outbreaks (Galindo et al. 2018) and even ship emissions (Monteiro et al. 2018a; Russo et al. 2018).

Another issue related to PM composition is its radionuclide content. Radionuclides in the atmosphere rapidly attach on submicron-sized aerosols, and their variability in ground-level air is driven by the behavior of aerosols (Povinec et al. 2012; Hirose and Povinec 2015). Atmospheric radionuclides are deposited from the air onto the land and sea surface by wet and dry deposition. In this way, the terrestrial and marine environments are labeled by natural and anthropogenic radionuclides that can be used as tracers of environmental processes (Pham et al. 2017). Radionuclide content can pose a health hazard following an accident involving nuclear material (Baeza et al. 2016). However, the occurrence of anthropogenic radionuclides in aerosols is also due to erosion and resuspension processes, as well as the emission and transport of particulate matter due to biomass burning as consequence of wild fires (Strode et al. 2012; Evangelou et al. 2014), and dust transport due to dust outbreaks (Hernández et al. 2005). Due to these processes, the anthropogenic radionuclide concentration in near surface atmosphere is variable. Naturally occurring radionuclides are also present in airborne particles as they are also present in soil particles able to be eroded, re-suspended or transported by the processes previously described, and also due to the radon exhalation from soil, which is especially significant to lead-210 and polonium-210 (Baeza et al. 2016).

2.3.2.3 Plastics (macro/micro/nano)

We live in the plastic age, since synthetic polymers are present in most aspects of human life both in developing and industrialized countries. The worldwide production for plastics increased annually by 10% since the 1950s, reaching 300 Mt in 2015 (Geyer et al. 2017). As of 2015 approx. 6,300 Mt of plastic waste had been generated, around 9% of which had been recycled, 12% was incinerated, and 79% was accumulated in landfills or the natural environment (Geyer et al. 2017). Synthetic thermoplastics constitute the most abundant and still growing component of anthropogenic debris entering the Earth's oceans (Ivar do Sul and Costa 2014). Up to 80%, or sometimes more, of the waste that accumulates on land, shorelines, the ocean surface or seabed is plastic (Barnes et al. 2009). The smallest form of plastic litter is called micro-plastic (<5 mm) and can represent up to 335,000 items km⁻² or 5 kg km⁻² in marine waters, and up to 25 kg km⁻² in coastal sediments (Koutsodendris et al. 2008; Ryan et al. 2009). Plastic debris, their dissolved derivatives, as well as, the adsorbed organic pollutants (Hirai et al. 2011) pose a direct risk to human and marine ecosystem health (Galloway 2015; Koelmans et al. 2017). As a rule, widely used plastics do not rapidly degrade

naturally when released into the environment, it can take 50 or more years for plastic to fully decompose (Müller et al. 2001).

In the Mediterranean Sea, the average density of plastic (1 item per 4 m²), as well as its frequency of occurrence (100% of the sites sampled), are comparable to the accumulation zones described for the five subtropical ocean gyres (Cincinelli et al. 2019), increasing the impact for marine biota with hotspots for the risk of plastic ingestion across multiple taxa especially in the coastal zone (Compa et al. 2019). Plastic debris in the Mediterranean surface waters (Fig. 2.16) was dominated by millimeter-sized fragments, but showed a higher proportion of large plastic objects than that present in oceanic gyres, reflecting the closer connection with pollution sources (Cózar et al. 2015). Multi-annual simulations of advected surface passive debris depict the Tyrrhenian Sea, the northwestern Mediterranean sub-basin and the Gulf of Sirte as possible retention areas (Mansui et al. 2015). No permanent structure able to retain floating items in the long-term were found, as the basin circulation variability brings sufficient anomalies to alter the distribution (Mansui et al. 2015).

Beyond the concern with “traditional” PM effects, an emergent research issue worldwide has been

focused in the occurrence of microplastics in the atmospheric compartment (MP; plastic particles with a longest dimension < 5 mm). MP may undergo photo-oxidative degradation in the environment, along with wind shear and/or abrasion against other ambient particulates, eventually fragmenting into fine particles (Gasperi et al. 2018). The risk of inhaling fibrous MP following widespread contamination within different environmental compartments deserves special attention owing to both the scale of their worldwide production and their potential to fragment into smaller, more bioavailable fibers. Human exposure to MP could also occur through ingestion, for example fibrous MP can settle on the floor; children, owing to crawling and frequent hand-to-mouth contact, ingest daily settled dust (Gasperi et al. 2018). Two studies have demonstrated the presence of MP in the atmosphere (Dris et al. 2016, 2017), thereby suggesting potential human exposure (none of these studies has been conducted in the Mediterranean Basin).

2.3.3 Chemical pollutants

2.3.3.1 Nutrients

Nutrients (mainly nitrogen, N, and phosphorous, P) constitute an important factor controlling marine




Figure 2.16 | Averaged 2013–2017 map of plastic debris concentration (g km⁻²) at the sea surface. Some geographical names used in the text are given: the (1) Gulf of Arzew, (2) Gulf of Tunis, (3) Gulf of Sidra, (4) Abu Qir Bay, (5) Cilician Sea, (6) Izmir, the (7) Thermaic Gulf, (8) Saronic Gulf, (9) Buna-Bojana, (10) NW Adriatic, (11) Taranto Gulf, (12) Gulf of Naples, (13) Gulf of Genoa, (14) Gulf of Lion, (15) Catalan Sea, and (16) Malaga Bay (Liubartseva et al. 2018).

primary producers, as they control phytoplankton growth, biomass and species composition (Sakka Hlaili et al. 2006). According to the nutrient concentrations, marine waters are characterized as oligotrophic (low nutrient concentrations), mesotrophic (nutrient enriched water), or eutrophic (nutrient rich water). The Mediterranean Sea is characterized by oligotrophic off-shore waters, with decreasing levels of nutrients eastwards from Gibraltar to the Levantine Sea (Ignatiades et al. 2009; Tanhua et al. 2013). The Eastern Mediterranean Sea is the most oligotrophic region, with very low nitrate concentrations ($< 0.5 \mu\text{M}$) and phosphorous ($< 0.2 \mu\text{M}$) (Pujo-Pay et al. 2011). Pronounced phosphorous limitation, with N/P ratio > 30 , is observed for the south of Levantine Sea and Ionian Sea (Kress et al. 2003; Pujo-Pay et al. 2011).

However, Mediterranean coastal areas, which are highly populated, are experiencing increasing N and P loading from anthropogenic activities, such as urban effluents, industrial discharges, agricultural runoffs, aquaculture activities and riverine inputs from a drainage area of $1.5 \times 10^6 \text{ km}^2$ (UNEP/MAP 2017). The overall inputs of N and P in these areas are about 1.5-4.5 and 0.1-0.4 Mt yr $^{-1}$, respectively. The main sources of N in Mediterranean are urban wastewater treatment (45%) and livestock farming (24%). Fertilizer use can also bring nitrogen and inputs can exceed 10^6 kg yr^{-1} (Fig. 2.17). Aquaculture contributed also to the emission on N (10%). For P, the main emitters are manufacture of fertilizers (40%), farming of animals (39%) and urban wastewater treatment (13%) (UNEP/MAP 2012b).

Some coastal regions are known as hotspots of nutrient inputs. In the North of Mediterranean Sea, the Lagoon of Venice and the Gulf of Lion sustained high nitrate levels, $18 \mu\text{M}$ and $9 \mu\text{M}$, respectively (Acri et al. 2004; Severin et al. 2014). Nitrate rich waters characterize also the Eastern Adriatic Sea ($4 \mu\text{M}$) (Skejic et al. 2017) and the Western Tyrrhenian Sea ($6.5 \mu\text{M}$) (Astraldi et al. 2002). In the southern Mediterranean, The Gulf of Gabès is a main region known for P enrichment ($1-11.2 \mu\text{M}$), since Tunisia is an important producer country of P fertilizers. Nitrate ($6-6.5 \mu\text{M}$) and ammonia ($\sim 4 \mu\text{M}$) showed also pronounced levels in this Gulf (Drira et al. 2016). High nitrate concentrations were often measured in other Tunisian coastal systems, such as the Lagoon of Bizerte (NO_3^- : $1-6.3$, NH_4^+ : $20-30 \mu\text{M}$) (Sahraoui et al. 2012) and the North Lake of Tunis (NO_3^- : $7.5-198 \mu\text{M}$) (Armi et al. 2011). In the Algerian-Provençal Basin and the Gibraltar Strait, enrichment of water with nitrate has been reported ($9.5-10 \mu\text{M}$) (Béthoux et al. 1998; Gómez et al. 2000).

Nutrient enrichment of Mediterranean Sea may result in a high increase in phytoplankton growth and biomass, leading to the eutrophication. The impacts of eutrophication include hypoxia or anoxia and may provoke harmful algal blooms, some of them toxic. Harmful algal blooms (HABs) cause human illness and mortality and have socio-economic impacts related to toxicity of harvested fish and shellfish, loss of aesthetic value of coastal ecosystems, and reduced water quality impacting tourism (Section 2.3.4).




Figure 2.17 | Fertilizer use and nitrogen release in Mediterranean Sea (UNEP-GRID Arendal 2013).

2.3.3.2 Gaseous pollutants: nitrogen dioxide, sulphur dioxide, ozone

The Mediterranean Basin is one of the regions in the world where high concentrations of gaseous air pollutants (nitrogen dioxide – NO₂, sulphur dioxide – SO₂, and ozone – O₃) have been reported frequently (Dayan et al. 2017). The elevated concentrations observed are attributed to the combination of diverse emission sources affecting the Mediterranean Basin including industry, traffic and shipping emissions (Karanasiou et al. 2014). The Mediterranean climate is characterized by arid conditions as well as many hours of sunshine and specific atmospheric recirculation patterns that significantly enhance air pollution levels (Karanasiou et al. 2014; Querol et al. 2018).

Commonly, NO₂ concentrations in North Mediterranean countries are higher than those observed in northern Europe (Cyrus et al. 2012). This fact was attributed to the transport sector, and to the higher conversion of nitric oxide (NO) to NO₂ caused by high temperatures and O₃ concentrations (Schembri et al. 2012; Karanasiou et al. 2014). The spatial pattern of N deposition varies across the Mediterranean Basin (Fig. 2.18). In Iberia, dry deposition is an important component of the total atmospheric N input to natural habitats (García-Gómez et al. 2018; Oliveira et al. 2020).

Ships are among the major emitters of air pollutants such as SO₂ and NO_x, their contribution to the emissions from the transport sector (Schembri et al. 2012) and to the air pollution in the Mediterranean Basin (Monteiro et al. 2018b; Russo et al. 2018) is growing. Several studies have also shown that NO₂ exceedances (yearly and hourly) in cities of the Mediterranean Basin are caused by road traffic emissions (Borrego et al. 2012; Belhout et al. 2018) (Fig. 2.19).

The large variety of Volatile Organic Compounds (VOCs), NO_x emissions and the climate conditions of Mediterranean Basin influences O₃ formation and destruction (Sahu and Saxena 2015; Sahu et al. 2016). These factors result in higher O₃ concentrations and frequent tropospheric O₃ episodes recorded across the Mediterranean Basin, with different frequencies in the East and West (Sicard et al. 2013).

The western Mediterranean Basin is characterized by frequent sea breezes, driven by sea-land thermal contrast. These sea-land breezes play an important role for the O₃ concentrations since they transport air masses, including O₃ precursor gases, from urban agglomerations located in coastal areas, towards inland suburban and rural areas (Millán et al. 2000).




Figure 2.18 | Modelled nitrogen deposition for the Mediterranean region based on the European Monitoring and Evaluation Programme (EMEP) model at 0.1x0.1° longitude-latitude resolution (EMEP MSC-W chemical transport model version rv4.7)⁶. Modelled N deposition is based on 2013 emissions data. (A) Total N deposition (oxidized+reduced; dry+wet), (B) percentage of dry deposition, (C) percentage of wet deposition, (D) percentage of oxidized deposition and (e) percentage of reduced deposition (Ochoa-Hueso et al. 2017).

High O₃ episodes in this region are linked to the combination of one or several of these mechanisms: (i) local/regional photochemical production and surface transport from coastal to inland regions;

⁶ www.emep.int



Figure 2.19 | Nitrogen dioxide (NO₂) concentrations above the annual limit value of 40 µg·m⁻³ (based on EU Directive 2008/50/CE).

(ii) O₃ transport from higher-altitude atmospheric layers, due to air mass re-circulation in the previous days; and (iii) long-range transport of O₃ and its precursor gases (Querol et al. 2018). In the eastern Mediterranean Basin, the O₃ episodes depends on the relative strength of the high-pressure system covering the eastern Mediterranean and Balkan area: (i) strong pressure gradient with northerly winds, creating good ventilation in the Athens Basin (Kallos et al. 2014); (ii) weak pressure gradient with local/regional O₃ events prevail; and (iii) stratospheric O₃ contributions to increase surface O₃ concentrations during specific meteorological scenarios (Zanis et al. 2014; Kalabokas et al. 2015). Tropospheric O₃ concentrations observed in the summer over this region (Fig. 2.20) are among the highest over the Northern Hemisphere (Dayan et al. 2017).

2.3.3.3 Trace metallic elements

Metal trace elements (MTE, or heavy metals in the old designation) whose main ones are cadmium (Cd), chromium (Cr), copper (Cu), lead (Pb), nickel (Ni),

zinc (Zn) and mercury (Hg) are common elements in the earth's crust (Navarro-Pedreño et al. 2008). They are also generated by human activities (Hassanien and Abdel-Latif 2008; Tovar-Sánchez et al. 2016; Merhaby et al. 2018). Urban and industrial wastewaters, atmospheric deposition and run-off from metal contaminated sites constitute the major sources of toxic metals (UNEP/MAP 2012a).

High MTE levels have been found in various soils (vegetated soils, dikes, waste rock and slag) at mining sites in Morocco. These mining activities in addition to agricultural and pastoral practices constitute a way of entry of the MTEs into the food chain and thus increase the risk of contamination of the population. Several plant species are adapted to these high levels of MTE and thus represent an important potential for the development of mining site rehabilitation strategies (Smouni et al. 2010). In agricultural soils of the Argolida Basin (Peloponnese, Greece), the MTE concentrations are high, following a decreasing order: Fe > Mn > Ni > Zn ~ Cr > Cu > Co ~ Pb > As > Cd (Kelepertzis 2014).



Figure 2.20 | Number of days (more than 25) above ozone (O₃) limit value of 120 µg·m⁻³ (based on EU Directive 2008/50/CE).

	Al	Cr	Cu	Fe	Hg	Mn	Ni	Pb	Ti	Zn
Min	36,866	75.2	42.4	26,313	0.04	552	47.9	44.7	2,343	86.8
Max	72,020	102.8	52.3	36,098	0.41	2,826	60.7	74.8	3,876	129.0
Mean	58,564	85.9	47.4	31,566	0.12	893	53.3	57.8	3,065	102.3

Table 2.2 | Metal concentrations ($\mu\text{g g}^{-1}$ dry weight) in marine sediment of Taranto Gulf (Ionian Sea, southern Italy) (Buccolieri et al. 2006).

In marine ecosystems, the hotspots of lead, mercury and cadmium were essentially located on the north-central and southeastern shores of the Mediterranean Basin (Fig. 2.21) (UNEP/MAP 2012b). Polluted surface samples on the Barcelona city continental shelf taken in 1987 reached enrichment factors of up to 490 for Hg, about 40 for Pb and Cd, and about 17 for Zn, Cr and Cu. In 2008, the data showed a decline with enrichment factors between 20 and 30 for Hg and Cd and between 5 and 12 for Zn, Cr, Pb and Cu (Palanques et al. 2017). In the Taranto Gulf (Ionian Sea, Southern Italy) (Table 2.2), MTE distribution is principally influenced by industrial and urban wastes. River discharges and prevailing anticlockwise marine currents are further factors influencing metal accumulation in sediments (Buccolieri et al. 2006). In surface sediments in Lebanon (eastern Mediterranean Sea), trace metals (Cd, Pb, Zn, and Cr) contamination at Beirut Port was classified as “the most highly polluted” and as “moderately polluted” at Tripoli Port (Merhaby et al. 2018).

Freshwater ecosystems are also affected by MTE pollution. In the Ichkeul Lake Basin (northeastern Tunisia) MTE showed concentrations in the sediment samples following the order: Fe > Mn > Zn > Pb > Ni > Cr > Cu > Cd (Touaylia et al. 2016). The concentrations of Fe, Cd, Ni and Cr in the bottom sediments of the Lower Litani River Basin (Lebanon) were higher in the dry season (Nehme et al. 2014). In river sediments from a semi-arid Mediterranean Basin (Algeria), MTEs were grouped by their level of contamination: high (Pb, Cd, Zn, Cu) and low (Al, Fe, Cr, Co, Ni). Sources of this contamination were essentially industrial, agricultural and domestic waste, as well as very specific ones (gasoline station) and diffuse pollution from atmospheric deposition (gasoline, ores, aerosols) (Benabdela et al. 2018).

MTE are known for their toxicity, persistence, and bioaccumulation in human and animal tissues, and biomagnify (concentrate at successively higher levels in tissues) in food chains (UNEP/MAP 2012b). In the northwestern Mediterranean Sea, the concentrations of 21 trace elements showed great variability in three species; fish

(sea bass, *Dicentrarchus labrax*), mussels (*Mytilus galloprovincialis*) and oysters (*Crassostrea gigas*). The essential elements (Cu, Mn and Zn) were highest in oysters, but Fe, Cr, Ni, Se, Co and Mo levels were highest in mussels. Fish had the lowest concentrations for all trace elements, which were at least one order of magnitude lower than in




Figure 2.21 | Mean concentrations of principal trace metals in coastal sediments of the Mediterranean Basin (UNEP/MAP 2012b).

bivalves. The maximum values set by European regulations for Hg, Cd and Pb were never exceeded (Squadrone et al. 2016). The levels of As, Hg and Pb measured in some commercially key species from Sicilian coasts suggest relatively reduced pollution levels for fish resources in this part of the Mediterranean region (Traina et al. 2019).

2.3.3.4 Organic pollutants

Polycyclic Aromatic Hydrocarbons (PAHs)

The contamination of the marine environment by polycyclic aromatic hydrocarbons (PAHs) has received considerable attention since the early 1980s after occurrence of tragic marine oil spills like the Amoco Cadiz oil spill [March 1979, Brest, France; 223×10^3 t crude oil]. PAHs consist of two or more fused benzene rings. PAH distribution is controlled by multiple and inter-dependent parameters. Some of these parameters are linked to intrinsic physicochemical properties of these compounds, mainly hydrophobicity which controls their partition between dissolved and particulate phases. Others are related to the hydrological and biogeochemical characteristics of the environment including water agitation and turbidity, sediment granulometry and particulate or dissolved organic matter content. They are issued from unburned petroleum or oil-derived products (petrogenic PAHs) or from the incomplete combustion of fossil fuels and biomass (pyrogenic PAHs). Petrogenic PAHs are characterized by lighter compounds (phenanthrene, naphthalene, fluorene) and their alkylated derivatives (methyl, dimethyl etc.) and have high affinity for the dissolved phase. Pyrogenic PAHs are characterized by unsubstituted higher molecular weight compounds that exhibit more affinity for suspended particles. Distribution of petrogenic-like and pyrogenic-like compounds help identification of PAH sources in the marine environment. Eight PAHs belong to the list of priority substances: Anthracene, Benzo (a) Pyrene, Benzo (b) Fluoranthene, Benzo (ghi) Perylene, Benzo (k) Fluoranthene, Indeno (123cd) Pyrene, Fluoranthene and Naphthalene. The top 5 are listed as priority hazardous substances due to their potential toxic, mutagenic and carcinogenic effects on organisms and human health (Hussein et al. 2016).

The diagnosis of the Mediterranean with regard to these substances is governed by international and national environmental agencies (i.e., the US Environmental Protection Agency (US-EPA), the European Environmental Commission, the United Nations Environment Program (UNEP), The National Observatory of the Environment and Sustainable Development, ONEDD (Algeria), National Environ-

mental Protection Agency, ANPE (Tunisia), Egyptian Environmental Affairs Agency, EEAA (Egypt) that recommend the monitoring of 16-19 PAHs with special emphasis on PAHs micropollutants in marine matrices.

The overall data set suggests that both petrogenic and pyrogenic PAHs contribute to the PAH pool in Mediterranean coastal waters. Pyrogenic inputs increase in urbanized areas due to large atmospheric inputs and deposition in winter (Guigue et al. 2014; Barhoumi et al. 2018). Total PAH inputs from the Rhône river represent 50 t yr^{-1} whereas inputs from sewage treatment plant are much lower ($<1 \text{ t yr}^{-1}$) (Witkowski et al. 2017). Due to their hydrophobicity, PAHs are considered to be preferentially associated with particles in coastal marine waters and sediments (Adhikari et al. 2015). Dissolved PAHs concentrations in the water column may be 4-fold higher than in suspended particles (Guigue et al. 2011). Dissolved PAH concentrations are in the range $0.158\text{--}3.655 \text{ ng L}^{-1}$ ($\Sigma 18$ PAHs) (Berrojalbiz et al. 2011) in Mediterranean open sea waters whereas higher concentrations characterize coastal urbanized areas, up to 560 ng L^{-1} ($\Sigma 32$ PAHs) in Marseille-Gulf of Fos (France) (Guigue et al. 2011), $12\text{--}267 \text{ ng L}^{-1}$ ($\Sigma 17$ PAHs) in Venice lagoon (Italy) (Manodori et al. 2006), $13\text{--}120 \text{ ng L}^{-1}$ ($\Sigma 7$ PAHs) in Alexandria coastal waters (Egypt) (El Nemr and Abd-Allah 2003). In coastal sediments, concentrations range from $10\text{--}200 \text{ ng g}^{-1}$ d.w. (dry weight) (Asia et al. 2009) in Marseille Bay, France (Gogou et al. 2000), Cretan Sea, Greece (Zaghden et al. 2005), Sfax, Tunisia (Cannarsa et al. 2014), Liguria, Italy (Merhaby et al. 2015), Tripoli harbor, Lebanon (Emara et al. 2008) and Eastern harbor, Egypt (Emara et al. 2008).

PAH concentration gradients are pronounced along coast-open sea transects. PAH concentrations rapidly decrease from the vicinity of rivers, estuaries and small effluents towards coastal and offshore waters. The contamination of estuaries of large rivers and that of harbors in the vicinity of big industrial and urban centers indicates a poor quality environment where a potential risk to the local population may occur (Barhoumi et al. 2018). Otherwise in most Mediterranean coastal waters, data reflect contamination levels from slightly polluted to polluted. Industrial areas near the cities of Sfax and Gabès (Tunisia) were reported moderate-to-highly impacted by hydrocarbons compared to other Mediterranean coastal environments (Fourati et al. 2018b, 2018a). They remain globally higher than those reported in the northern Gulf of Mexico and much lower than those recorded in Chinese coastal environments (Zhou and Maskooui 2003; Fourati et al. 2018b).

In some cases, PAH concentrations are influenced by physical circulation processes that can lead to deposits of contaminants an order of magnitude higher offshore than those near the source of pollutants. Episodic processes of pollutants redistribution may also significantly affect the pollution status of marine areas. For example, the physical accumulation of PAH at air/sea interface ($\times 200\text{-}1,000$) during microlayer formation in absence of wind (Wurl and Obbard 2004), followed by PAH scattering during microlayer disruption by wind blow recovery, can locally enhance PAHs concentrations and impact the biota. Similarly, PAH sediment remobilization during resuspension events may greatly modify their potential harmful effects on marine biota (Guigue et al. 2017). In Toulon bay (France), the resuspension of highly contaminated surface sediments (concentration of $\Sigma 34$ PAHs = 38.2×10^3 ng g⁻¹) led to a 10-fold increase of dissolved $\Sigma 34$ PAH concentrations in the water above. The remobilization in seawater was higher for 4-6 ring PAHs, especially benzo(g,h,i) perylene, whose concentration exceeded the authorized limit values of the European Water Framework Directive (Guigue et al. 2017). It is important to monitor pollutants not only at active industrial facility sites but also in disused industrial areas close to the sea border where remnant pollution can produce chronic adverse effects on marine biota.

Species feeding on particles and phytoplankton may bioaccumulate and/or bioamplify PAH concentrations in their body tissues. Measuring accumulation of PAHs in mussel bivalves from the *Mytilus edulis* complex has become a European Commission control strategy of marine waters quality in the Mediterranean (Olenycz et al. 2015; Sire and Amouroux 2016). Different metabolites may be measured in fish and shells and considered as markers of exposure to PAH. Research on the relationships with emergent contaminants is at its beginning and cocktail effects have not been much studied yet. Microplastics have a high potential to adsorb these hydrophobic contaminants and to transfer them throughout the food web to the deep ocean for longer sequestration time.

Pesticides

Pesticide Active Ingredients (PAI) can be considered as a contaminant as well as a pollutant, in the compartments where they are detected. Mainly originated by agricultural activities, water pollution by PAI is a concern for continental water resources (rivers, lakes and aquifers) and coastal and marine environment of the Mediterranean Sea. Studies

carried out in southern Europe showed the high leaching of herbicides in Mediterranean weather conditions (Louchart et al. 2001), allowing the contamination of groundwater resources. In the other side, by runoff process, surface waters would be contaminated by PAI and their metabolites or degradation products. A large number of pesticide active ingredients (PAI) (over 1,300) are presently used or were used until non-approval or non-renewal in Europe (European Food Safety Authority 2011) and even modern screening methods limit the number of PAI analyzed in one sample (< 450) (Rousis et al. 2017). In water bodies, the maximum allowable concentrations are 2 µg L⁻¹ for each PAI and 5 µg L⁻¹ for all quantified PAIs. For drinkable water these limits are 0.1 µg L⁻¹ and 0.5 µg L⁻¹ respectively.

Concentrations of these molecules in water bodies (surface and underground waters) were qualified and quantified in nearly all the countries around the Mediterranean Sea. But a recent review at world scale (Stehle and Schulz 2015) stated the difference of availability of referenced data sets for insecticide concentrations in water: notably in the North and the North-East Africa poor information was available.

In European countries, particularly in France (Dubois et al. 2010; Lopez et al. 2015), Italy (Onorati et al. 2006; Meffe and de Bustamante 2014), Spain (Balaguer et al. 2018) and Greece (Lekkas et al. 2004; Konstantinou et al. 2006) implementation of the EU Water Framework Directive (WFD) produced large public data sets for pesticide concentrations in surface waters and aquifers by state administrations. According to the statistical office of the European Union, Spain and Italy are the countries with most use of pesticide. As a result, in those countries, pesticides are one of the most frequently detected classes of micro-pollutants in water⁷. It is not possible to give in this report an exhaustive account of all PAI mentioned in the literature as encountered in Mediterranean waters. Thus, we decided to focus on the most frequently detected PAI and give some examples of maximum concentrations measured.

In water bodies most frequently mentioned PAI-insecticides already in use are chlorpyrifos (18.8 µg L⁻¹) (Ccanccapa et al. 2015), dimethoate (0.640 µg L⁻¹) (Campo et al. 2013), malathion (0.048 µg L⁻¹) (Yurtkuran and Saygi 2013), imidacloprid (0.350+/- 0.433 µg L⁻¹) (Herrero-Hernández et al. 2013) and diazinon (14.5 µg L⁻¹) (Youssef et al. 2015). Prohibited PAI-insecticides mentioned are DDT (dichlorodiphenyltrichloroethane) and its metabolites (0.40 to 3.22 µg L⁻¹) (Dahshan et al. 2016), HCB (hexa-

⁷ <https://www.eea.europa.eu/airs/2018/environment-and-health/pesticides-sales>

chlorobenzene) ($1.1 \mu\text{g L}^{-1}$) (Youssef et al. 2015) and endosulfan ($0.247 \mu\text{g L}^{-1}$) (El Bakouri et al. 2008).

The most frequently mentioned PAI-herbicides in use are simazin ($3.18 \mu\text{g L}^{-1}$) (Konstantinou et al. 2006), terbutylazin ($0.0219 \mu\text{g L}^{-1}$) (Ricart et al. 2010), linuron ($13.13 \mu\text{g L}^{-1}$), 2,4-D ($20 \mu\text{g L}^{-1}$) and glyphosate with its metabolite AMPA ($167 \mu\text{g L}^{-1}$) (Meffe and de Bustamante 2014). Most prohibited PAI-herbicides mentioned in studies are atrazine-desethyl ($0.158 \mu\text{g L}^{-1}$) (Campo et al. 2013), metolachlor ($1.120 \mu\text{g L}^{-1}$) (Konstantinou et al. 2006), DEA (diethyl-atrazine) ($1.98 \mu\text{g L}^{-1}$) (Hildebrandt et al. 2008), diuron ($0.0169 \mu\text{g L}^{-1}$) (Robles-Molina et al. 2014), DIA (deisopropyl-atrazine) ($8 \mu\text{g L}^{-1}$) (Shomar et al. 2006), alachlor ($0.213 \mu\text{g L}^{-1}$) (Stamatis et al. 2013), isoproturon ($7 \mu\text{g L}^{-1}$) (Ricart et al. 2010) and molinate ($0.026 \mu\text{g L}^{-1}$) (Gómez-Gutiérrez et al. 2006).

Concentrations of PAI-fungicides are less mentioned than other PAI. Metalaxyl ($0.49 \mu\text{g L}^{-1}$) (Hildebrandt et al. 2008) and carbendazim ($1.81 \mu\text{g L}^{-1}$) (Licciardello et al. 2011) have been reported.

For the future, in most southern Mediterranean countries, prohibitions of particular PAIs for agricultural uses are applied within a few years delay from European Commission decisions. Many countries developed programs to reduce pesticide use that have uncertain effects like in France (Guichard et al. 2017). New proposed PAI are characterized by shorter standard half-lives and lower dose requirements, nevertheless there is no assurance that these "new" PAI once used widely in many different contexts will have virtuous environmental spreading behaviors.

Climate changes may have conversational effects on PAI water contamination. Firstly, some authors predict an increase use of pesticides to compensate increases of abundance and seasonal activity of bioaggressors (Boxall et al. 2009), although temperature enhancement has been declared to decrease PAI efficiency by conditional resistance of bioaggressors towards herbicides, insecticides and fungicides (Matzrafi 2019). Under these conditions, doses applied should be increased to guarantee the same protection with unseen consequences in environment spreading. Daily temperature fluctuation may increase PAI toxicity like for chlorpyrifos (Verheyen and Stoks 2019), but temperature enhancement should boost degradation of most known PAI probably shortening their half-life.

Change in precipitation patterns with increased occurrence of extreme precipitation events should also modify agricultural patterns. For instance,

rained barley Mediterranean production yields would decrease (Verheyen and Stoks 2019) fueling some changes in the crop systems. Implementation of more intensive systems (Malek and Verburg 2018) can imply an increase of treatments. Increasing weight of extreme events would increase erosion (Raclot et al. 2018) and then promote displacement from land to sea for the PAIs easily adsorbed on soil particles like clay and organic matter.

2.3.3.5 Emerging contaminants

In the Mediterranean Basin, 63% of coastal settlements with more than 2,000 inhabitants operate a wastewater treatment plant, while 37% do not. Secondary treatment is mostly used (67%) in Mediterranean treatment plants, while 18% of the plants have only primary treatment (Chatha et al. 2017). As a consequence of this technical, social and environmental issue, different types of chemical substances are released into the environment (Gros et al. 2010; Ratola et al. 2012; Moreno-González et al. 2015; Paluselli et al. 2018b, 2018a). Among these substances, emerging contaminants (ECs) are a category that has received special attention over the last 25 years. ECs are defined as "contaminants of emerging concern that are naturally occurring, manufactured or man-made chemicals or materials which have now been discovered or are suspected present in various environmental compartments and whose toxicity or persistence are likely to significantly alter the metabolism of a living being" (Sauvé and Desrosiers 2014).

There are different classifications of these contaminants due to their usage or origin and effects. For example, (1) antibiotics, (2) antimicrobials, (3) detergent metabolites, (4) disinfectants, (5) disinfection byproducts, (6) estrogenic compounds, (7) fire or flame retardants, (8) fragrances, (9) insect repellants, (10) PAHs (polycyclic aromatic hydrocarbons), (11) personal care products, and (12) pesticides or insecticides (13) pharmaceuticals, (14) plasticizers, (15) reproductive hormones, (16) solvents, (17) steroids and (18) surfactants (Singh and Kumar 2017).

ECs that are very soluble in water (tetracycline, sulfamethoxazole, carbamazepine, and erythromycin, etc.) receive more attention than others because of their impact on the environment (Klaper and Welch 2011). Potential ECs sources and pathways of ground and surface water pollution are shown in Fig. 2.22. Typically, the route of these compounds towards a water body begins with the excretion of the metabolites and parent compounds and their disposal to the wastewater treatment plants (Barrios-Estrada et al. 2018).




Figure 2.22 | Potential sources and pathways for grounds and surface water pollution (Barrios-Estrada et al. 2018).

The endocrine disruptors (EDs) are a subdivision of ECs with chemicals that interfere with the body's endocrine system and produce adverse developmental, reproductive, neurological, and immune effects in humans, abnormal growth patterns and neurodevelopmental delays in children (Moreno-González et al. 2015). The majority of the EDs come from products used to fight undesirable wildlife and agricultural threats (Moreno-González et al. 2015), for example, pesticides, fungicides and rodenticides, synthetic products used in plastic industry (bisphenols or phthalates) and a variety of buildings materials, isolation materials (polychlorinated biphenyl and metals). A list of common sources of EDs is shown in Fig. 2.23.

The presence of ECs in Mediterranean Basin is well documented. Phthalic Acid Esters (PAEs), including dimethyl phthalate (DMP), diethyl phthalate (DEP), di-isobutyl phthalate (DiBP), di-n-butyl phthalate (DnBP), benzylbutyl phthalate (BzBP) and diethylhexyl phthalate (DEHP), with total concentrations ranging from 130 to 1,330 ng L⁻¹ were found in Marseille Bay (northwestern Mediterranean Sea) (Paluselli et al. 2018b). High concentrations of PAEs were also observed in deep waters offshore (310.2 ng L⁻¹) as well as in the Rhône River (615.1 ng L⁻¹) (Paluselli et al. 2018a).

A total of 20 pharmaceuticals in sea water and 14 in sediments were found at concentrations from

low ng L⁻¹ up to 168 ng L⁻¹ (azithromycin) in sea water and from low ng g⁻¹ up to 50.3 ng g⁻¹ (xylazine) in sediments of Mar Menor lagoon located in the South East of Spain (Moreno-González et al. 2015). Pharmaceutically active compounds (PhACs) were detected in the Evrotas River (Southern Greece) waters. The diuretics and the analgesics/anti-inflammatory class were the most abundant, followed by antihypertensives, psychiatric drugs, β-blocking agents and antibiotics and the concentration levels ranged from 0.31 ng L⁻¹ up to 51 ng L⁻¹ (Mandaric et al. 2019). Antibiotics were detected in more than 90% of the water samples collected from a Mediterranean river (Llobregat, Spain) and the concentration levels ranging from 0.3 ng L⁻¹ (flumequine) to 907.6 ng L⁻¹ (sulfamethoxazole) (Proia et al. 2013). Triclosan (an antimicrobial) was reported to be a contaminant of the Llobregat and Ebro rivers (Spain) and the concentrations in some samples were higher than 150 ng L⁻¹. These concentrations should be considered significant considering the toxicity of these compounds and their expected ability to be a precursor of other highly toxic compounds such as dioxins (Kantiani et al. 2008). Azithromycin (antibiotic) was measured at 16,633 ng L⁻¹ in a tributary of El Albujón (Spain) (Moreno-González et al. 2015) and acetaminophen was detected at 3,000 ng L⁻¹ off Thessaloniki, Greece (Nödler et al. 2014). Some drug classes, such as analgesics, antibiotics and betablockers, were still quantified at levels between 0.3 (metoprolol) and hundreds of ng L⁻¹ (azithromycin) in seawater. In




Figure 2.23 | Common sources of endocrine disruptors in the environment (Barrios-Estrada et al. 2018).

particular, six substances (e.g., azithromycin, amoxicillin, venlafaxine, salicylic acid, acetaminophen and ibuprofen) were measured at median concentrations higher than 20 ng L^{-1} . Pharmaceuticals were also found in coastal and oceanic waters adjacent to the Strait of Gibraltar (Biel-Maeso et al. 2018).

The impacts of exposure to some ECs have caused concern for both people and wildlife. Many of these substances may cause disorders of the nervous, hormonal and reproductive system, thus posing adverse health outcomes (Rezg et al. 2014; Bilal et al. 2018, 2019; Hernandez-Vargas et al. 2018; Ullah et al. 2018; Rasheed et al. 2019). These synthetic chemicals exhibit geno- or cytotoxic activity and can cause obesity, diabetes, cardiovascular and reproductive disorder or even leads to cancer (Tiwari et al. 2012). Etteieb et al. (2016) have shown that the components responsible of cytotoxicity in water samples from Medjerda river in Tunisia were mainly cyclopentasiloxane, decamethyl (D5), cyclohexasiloxane, dodecamethyl (D6), D-limonene, and ergoline-8-methanol, 8,9-didehydro-6-methyl. Some ECs, such as 17- α -estradiol, bisphenol A and phthalates were reported to alter marine community structure (Essid et al. 2013; M'Rabet et al. 2019).

2.3.4 Biological pollutants

Numerous viruses or bacteria of human or animal origin can spread in the environment and infect peo-

ple via water, air and food, mostly through ingestion and occasionally through skin contact. These viruses and bacteria are released into the environment by various routes including water run-offs and aerosols. Furthermore, they can infect humans exposed to contaminated surface waters and ground water used for agriculture irrigation with severe consequences for human health (de Giglio et al. 2017). In most semi-arid areas, groundwater and surface waters constitutes an important and strategic resource, particularly as water stress increases and water resources of good quality become scarce (El Ayni et al. 2013). Molecular epidemiology and regular surveillance are necessary to elucidate the public health hazards associated with exposure to environmental viruses and bacteria (Cabral 2010; Rodriguez-Lazaro et al. 2012) especially in the South region where the water contamination combined to water scarcity strongly have large socio-economic impacts (UN-Water 2014). Climate projections for the Mediterranean climate areas estimate general warming and changes in precipitation distribution. Mediterranean coastal rivers are subject to flash floods during extreme events that transport the majority of the annual loads of bacteria and other contaminants (Chu et al. 2011). The frequency of extreme summer precipitation events increased over large regions of the Mediterranean (Giorgi and Lionello 2008) (Sections 2.2.5 and 3.1.3.3), increasing so the supply of fecal bacteria and viruses to the coastal zone. In a global context, wastewater management will be the key to preventing environmental dispersion of human




Figure 2.24 | Harmful algal blooms recorded before 1980 (A) and between 1980 and 2015 (B) along the Mediterranean coast (Cecchi et al. 2016).

fecal pathogens in future climate change scenarios (Rusiñol et al. 2015).

Harmful Algal Blooms (HABs) are sporadic phenomena triggered by massive proliferations of phytoplankton species reaching high cell concentrations (Sellner et al. 2003) that are suspected to have greater occurrences due to global warming (Hallegraeff 2010). HABs have environmental impacts (red-tide, mucilage production, anoxia) (Rodger et al. 2011; D'Silva et al. 2012) and represent serious economic threat for fisheries, aquaculture and tourism (Hoagland et al. 2002). They may also harm human health, since 40% of blooming microalgae are able to produce toxins responsible of different human intoxications (Santi Delia et al. 2015). HABs can occur in both freshwater and marine environments. In the Mediterranean Basin, marine ecosystems have received more attention although occurrence of HAB in freshwater lakes are reported (Cook et al. 2004; Romo et al. 2013) and are suspected to increase with climate change in the region (Romo et al. 2013) with consequences for potable water supply (Lévesque et al. 2014). Before 1980, HABs were rarely documented in the Mediterranean Sea. Since then, adverse events and several toxic episodes have been reported in different coastal regions (Fig. 2.24) (Cecchi et al. 2016; Garrido et al. 2016). Nowadays, harmful and toxic phytoplankton species become dominant in many coastal environments of the Mediterranean Sea.

Most toxic events in the Mediterranean Sea are mainly related to the dinoflagellates *Alexandrium* (known to induce Paralytic Shellfish Poisoning - PSP) and *Dinophysis* (producers of toxins causing the Diarrheic Shellfish Poisoning - DSP) (Vila et al. 2001; Penna et al. 2007). *A. minutum* is the most observed dinoflagellate, with recurrent blooms (10^3 - 10^8 cells l $^{-1}$) in the Ebro Delta and the Gulf of Gabès, inducing significant fish mortality (Abdennadher et al. 2012; Garcés and Camp 2012). Blooms of *A. catenella* (10^4 cells l $^{-1}$) have been observed both in the northern (Thau Lagoon) and the southern Med-

iterranean Sea (Bizerte Lagoon, Tunisia), causing sometimes shellfish contamination (Laabir et al. 2013). The epi-benthic dinoflagellate *Prorocentrum lima* (producer of DSP toxins) has been detected on most Mediterranean coasts (Garcés and Camp 2012), sometimes with high densities ($>10^7$ cells l $^{-1}$) (Aissaoui et al. 2014; Moncer et al. 2017). Blooms of *Karenia selliformis* (10^3 - 10^5 cells l $^{-1}$), associated with intensive fish mortality, were reported for many years in the Gulf of Gabès (Feki et al. 2013). Recently, human health problems are caused by blooms of *Ostreopsis cf. ovata* (10^3 - 10^4 cells l $^{-1}$) in Italy, Spain, Algeria and France (Ciminiello et al. 2014). New records of *Gambierdiscus* and *Fukuyoa* in the West and East of the Mediterranean Sea increase the risk of Ciguatera intoxication (Laza-Martinez et al. 2016).

Toxic diatoms *Pseudo-nitzschia* (producers of toxins causing the Amnesic Shellfish Poisoning - ASP) showed also blooms (10^3 - 10^7 cells l $^{-1}$) in Mediterranean coasts of Spain, France, Greece, Italy and Tunisia, and contamination of mussels by ASP were reported (Sakka Hlaili et al. 2016). Mussel contamination by ASP toxin were reported in Spanish and Tunisian waters and have been linked to the blooms of *Pseudo-nitzschia* spp. and *Nitzschia bizertensis*, respectively (Giménez Papiol et al. 2013; Bouchouicha-Smida et al. 2015).

Toxic and harmful algal blooms continue to increase in magnitude, frequency and geographical distribution around the world and over the Mediterranean Sea (Hallegraeff 2010). The chronic eutrophication and climatic change including global warming are reported as significant factors involved in this global increase of HABs and toxic events (Anderson et al. 2012; Sakka Hlaili et al. 2016).

2.4 Land and sea use changes

2.4.1 Land use changes

2.4.1.1 Past trends and recent dynamics

Mediterranean landscapes

The climatic conditions of Mediterranean Basin have played a key role for drawing its landscapes, which result from the interaction between human activity, a complex topography, and an extreme varied soil and climate (Pinto-Correia and Vos 2004). Altitudinal gradients and the distance from the sea are the main factors differentiating the landscapes of the Mediterranean Basin (Pinto-Correia and Vos 2004), and the main ecosystems contributing to outline these landscapes include the Mediterranean shores with a high number of habitats such as areas of dunes, rocks and wet areas. Moving away from the coast, Mediterranean maquis (a typical dense and closed shrub vegetation, mainly constituted by sclerophyllous species) and forests evolve, which includes evergreen shrubs or trees fully adapted to xeric conditions. In many cases, the maquis derives from the evergreen Mediterranean forest that develops in less water limiting growing conditions. Mediterranean steppic prairies represent the typical degradation of maquis that advances desertification. Finally, the Mediterranean deciduous forest extends where climates gradually shifts from typical Mediterranean to inland.

The modifications occurred in the Mediterranean landscapes are the result of land use practices that increased in intensity since the Neolithic, when in the Middle east domestication of plant and animals took the place of hunting and gathering (Blondel 2006). Human pressure on the natural ecosystem greatly varied depending on societal evolution developed over the area. In any case, wood exploitation has been key in localized periods of empires expansion. Forest destruction was the first step consequence of human pressure on the natural habitat as the effect of increasing agricultural activity and livestock and wood exploitation in the Mediterranean Sea trade cycle in order to cover the needs of the maritime empire (Barkaoui 2003). There is a long history of urbanization, reflected in countless archeological and historical settlements over the region (Diappi 2015).

As a result, the Mediterranean landscape is a complex mosaic of alternating semi-natural hab-

itats. Grasslands and pastures represent one of the most spread land use in European Mediterranean areas in the plain and low hilly areas (Cosen-tino et al. 2014) and they can provide forage for grazing animals or hay for conservation. The use of terraces in hilly areas permitted the cultivation on slopes of olive groves, vineyard and sowing crops while at the same their cultivation represented a means to reduce soil erosion, prevent run-off and increase water saving (Blondel 2006). The Dehesa-Montado System (typical of Spain and Portugal) is characterized by low density trees (represented by evergreen Mediterranean oaks) combined with crop production or pastoral activities, mainly represented by animal grazing. This system integrates the three main rural activities (forest/cork product harvesting, livestock husbandry, and agriculture) within a single landscape that consists of grass and trees. This system that combines extensive grazing of natural pastures, cereal cultivation and harvest of wood products, has shown remarkable stability, biodiversity, and sustained productivity over 800 years or longer as the result of the maintenance of botanically rich mosaic-like herbaceous plant layers (Joffre and Rambal 1993). In addition to agricultural production, these complex systems contributed, at the same time, to several ecosystem services such as preservation of the environment and its natural resources securing the sustainability of the system (Blondel 2006; Hao et al. 2017).

Despite small-scale traditional farming systems has been practiced for long time and it is still adopted in many parts of the Mediterranean region, huge changes in agricultural practices have taken place during the last 50-100 years across several Mediterranean areas as driven by increasing profitability of new agricultural systems (Debolini et al. 2018). The greatest changes involved uprooting of ancient and small-scale vineyards, orchards and olive groves, which turned into industrial scale fruit or olive plantations. Similarly, mixed rotational farming systems were simplified and replaced usually by intensive monocultures, requiring high inputs (i.e., high fertilization rate and water requirement) (Debolini et al. 2018). Intensive and large-scale farming systems also required the creation of new infrastructure, such as basins to cope with water shortage, which contributed to change the natural landscape. All these changes caused unsustainable pressure on the surrounding environment, resulting in either loss of wildlife-rich habitats and socio-economic

viability over large parts of the region due to land-abandonment of small-scale farmers.

Recent changes in Mediterranean landscapes

The rate of change of landscapes in the Mediterranean Basin has increased since the second half of the 20th century. Many regions in Mediterranean Europe experienced the abandonment of marginal lands, especially in arid and mountain areas (Lasaña et al. 2017) (*Section 3.2.3.1*), and the following development by shrubs and tree species. Many studies show the abandonment of typical features like farming terraces, olive orchards, and upland grasslands leading to non-managed reforestation; for example, in 29% of the Iberian Peninsula for the 1989-2004 period (Hill et al. 2008); all-over Italy during 1990-2013, small forest patches cover increased a 27.4% (Sallustio et al. 2018), while 24% of pastures were turned into forests in areas of Tuscany from 1954 to 2005 (Amici et al. 2017); in Italian mountains, 16.3% of agricultural areas disappeared during the 1990's (Conti and Fagarazzi 2015); similarly in pre-alpine France during 1956-1991 (Taillefumier and Piégay 2003); in areas from the Eastern Mediterranean region of France, 14.2% of crops and 78.2% of pastures were converted into forests from 1958 to 2010 (Abadie et al. 2018); a 35.6% of vineyard areas in Serbia was transformed into meadows and pastures during 1985-2013 (Perović et al. 2018); Mediterranean islands as Elba, where up to 52% of agricultural areas were abandoned during the 1954-2000 period (Carta et al. 2018), and Lesvos, showing recent (2001-2011) slight decreases in agricultural land (Van der Sluis et al. 2016).

The remaining agricultural systems have generally become more intensive, with a shift towards livestock production and an increase of industrial inputs (fertilizers and pesticides), as it has been reported for Spain, especially since the 1960's to 2008 (Guzmán et al. 2018), and punctual areas of Greece and Italy, from 2001 to 2011 (Van der Sluis et al. 2016) (*Section 3.1.2.1*).

Conversely, scarce and mainly pre-21st century studies from North Africa show extreme land degradation due to overexploitation (Le Houérou 1995), principally by grazing pressure but also by forest conversion to agriculture and fuel-wood recollection. For the total North Africa and Middle East regions, the rate of deforestation increased from the 1980's to the 1990's by 160%, the fastest increase worldwide (Hansen and DeFries 2004). For example, in Morocco, it has been registered forest regression and degradation from 1962 to

1992 (Rejdali 2004), while in Rogassa (Algeria) a long-term experiment (1975-1993) demonstrated the main role of overgrazing in such degradation (Slimani and Aidoud 2004). Recent studies based in new observations and remote sensing show similar desertification trends in most parts of the Maghreb (Hirche et al. 2018). Grazing has been also the principal factor of forest degradation in northern Mediterranean islands as Crete during the 1977-1996 period (Hostert et al. 2003). Moreover, increases over 10% in livestock density have been observed between 2001 and 2011 in Portofino (Italy) and Lesvos (Greece) (Van der Sluis et al. 2016), and up to 40.1% in Nisyros (Greece) from 1991 to 2001 (Petanidou et al. 2008), showing that land abandonment does not always result in forest encroachment.

Changes in Mediterranean landscapes have been particularly intense in metropolitan areas and their surroundings. These landscapes are growing very rapidly all-over the Mediterranean (17% between 1990 and 2000) (Underwood et al. 2009) but especially in coastal areas. Urbanization mainly occurred at the cost of arable land (e.g., in Barcelona) (Basnou et al. 2013) and forested areas, as generally reported in the Mediterranean (Gerard et al. 2010). However, in some periurban areas, agricultural land has also increased following the growth of cities, as in Murcia (Spain) between 1995 and 2007 (García-Ayllón 2018). Examples could be found also in eastern Mediterranean, such as in Erdemli (Turkey), where the total length of the roads increased 23.6% between 2004 and 2015 following the growth of periurban agriculture (Alphan 2018).

2.4.1.2 Principal impacts of land use changes

Forestry and other natural resources

New forests after land abandonment could alter biodiversity patterns in the Mediterranean (Fabbio et al. 2003), as has been demonstrated for the range of certain bird species (Gil-Tena et al. 2010). New forests are established by tree species whose dynamics have been favoured by recent land use changes, even more than climate (Améztegui et al. 2010). Meanwhile, other species as oaks are less benefited by such changes (Acácio et al. 2017), although general patterns show in fact successional dynamics towards *Quercus* dominance at the expense of *Pinaceae* (Alfaro-Reyna et al. 2018). Fire regimes are also altered, as fuel continuity is increased facilitating fire spread, which in turn could result in more landscape

homogeneity (Loepfe et al. 2010). Also, forest continuity could increase forest insect pest spread (Hódar and Zamora 2004). The effect of increased forest cover in the diminution of water resources is more conflictive although the pattern is clearer in catchments with records of large and rapid forest expansion (Gallart et al. 2011) (*Section 3.1.1.3*). During the first stages of land abandonment there is also a great risk of land degradation due to soil and nutrient loss (Thornes 2009). Nevertheless, the contribution of these new forests to carbon storage is certain (Vilà-Cabrera et al. 2018).

Despite the natural forest recovery after land abandonment, forests in the Mediterranean Basin are still interested by many different threatening factors, such as wildfires, overgrazing, incorrect management and extreme climate and meteorological events (drought, windstorms) and this in turn could lead eventually to desertification (Vilà-Cabrera et al. 2018). Wildfires become more and more frequent due to drier summers coupled with wood expansion due to land abandonment (Pausas and Fernández-Muñoz 2012). Although evidence indicates that fires are decreasing due to increased efforts in fire suppression all-over Mediterranean Europe in more recent periods (Turco et al. 2016), there is large potential risk of mega-fires in the near future (Loepfe et al. 2011). Changes of traditional silvicultural schemes able to produce high productive goals to a more extensive management or forest abandonment has led to deep modifications of composition and structure.

Another major factor affecting degradation of Mediterranean forests is grazing by domestic animals that utilize understory especially in period of reduced forage availability in pastures (such as summer or, in certain cases, winter). The traditional grazing in forest formation should not be banned as when the stocking rate is adequate, this result in a proper sustainable forest management that can reduce potential fuel biomass and can preserve this tradition landscape (Kairis et al. 2015) while, on the contrary, overgrazing can produce erosion, reduction in soil cover, losses of nutrients and in this way is it can be considered one of the most important factors of desertification in Mediterranean areas (Papanastasis and Kazaklis 1998). Droughts, heat waves or windstorms can have a negative effect on many forests in the Mediterranean Basin producing reductions in forest growth and of forest declines (Vayreda et al. 2012) that prelude to land degradation and, in turn, desertification. All these factors produce stresses to forests that can be exacerbated in the next decades by climate change (Valladares et al. 2014).

Land degradation is the principal consequence of plant cover loss (by uncontrolled forestry, overgrazing, fires, etc.), which could lead to desertification in combination with increasing aridity and extreme climatic events (Thornes 2009). In mountain environments, forest and understory cover loss, principally by overgrazing, have been clearly associated to higher erosion rates (Cheggour et al. 2012), as well as recent increases in forest cover were linked to the opposite (Barreiro-Lostres et al. 2017). Soil erosion is a serious problem throughout the Tunisian Dorsal (the easternmost part of the Atlas mountain range). It has been estimated that 7% of the area is badly damaged by erosion and 70% of the area is moderately damaged (DG/ACTA 1993). This degradation is an accumulated effect of agricultural strategies adopted over the Tunisian semi-arid areas during the last three millennia (Jebari 2009). If we only consider short-term effects, the degradation can be partly attributed to the building of large dams during the 1960s and 1970s. This was done without giving sufficient attention to proper management of upper catchment areas (Jebari et al. 2010). However, the contribution of the specific bioclimatic conditions of the Mediterranean climate in this degradation should not be underestimated. The soils are better characterized by the degradation of rock material rather than their organic matter content (Cerdan et al. 2004; Cudennec et al. 2007). Consequently, they are not well developed and often shallow. The human influence is crucial on catchment scale in terms of landscape degradation that affects the hydrological regime (Jebari et al. 2010). In the last hundred years, continuous changes were undertaken by the introduction of new crop cover (in fact, the most important change leading to better water management in the southern part of the Mediterranean may come from improving water efficiency in agricultural irrigation) (Berndtsson et al. 2016), deforestation, urbanization, river network modification, dam buildings and embankment. Fortunately, better water resource planning, reservoir maintenance, shortage, flood management, and hydro-agricultural infrastructure design are currently promoted (Verkerk et al. 2017) (*Section 4.5.1*).

Urbanization is considered a major driving force of biodiversity loss and biological homogenization (Grimm et al. 2008), causing landscape fragmentation, dramatic loss of open habitats and of the land use gradient, replacing adjacent land uses such as agricultural and more natural vegetation. Urbanization is also one of the main drivers of introduction of non-indigenous species, generating high propagule pressure, and frequent and intense

disturbance with complex consequences for biodiversity (Basnou et al. 2015; Clotet et al. 2016). The strong human pressure has also contributed to increase water shortages, pollution, forest fires, and the abandonment of ancient pastoral regimes. Recent studies also demonstrate the negative consequences of new artificial areas in coastal dunes habitats, both affecting carbon stocks (Carranza et al. 2018), and generating more complex effects when stabilizing the natural changing dynamics of dune systems (Manzano et al. 2019).

Crops and livestock

Food production in the Mediterranean region is changing rapidly, due to multiple local and global social and environmental changes. The increased number of urban and displaced people increase demand for food in urban areas, with limited agricultural production and with great water restrictions (FAO 2017a). Scarce resources, such as fisheries, are being exhausted (see next section). Evidence for the limited capacity to cope with these challenges can be documented in recent history. For example, water reserves were not able to cope with extensive droughts in the last two decades in Spain, Morocco and Tunisia, causing many irrigation dependent agricultural systems to cease production (Faurès et al. 2002; Garrido et al. 2006; FAO 2015).

Livestock production, mainly located in semi-arid and arid lands has shifted from extensive modes to systems heavily dependent on feed grain (32% of total food imports), inducing high poverty rates and rural exodus and rendering production sensitive to climatic shifts elsewhere (Sections 3.1.2.1 and 4.5.1).

Besides soil erosion, the major land degradation processes in the Mediterranean Basin are soil sealing, compaction mainly due to agricultural intensification, salinization, and contamination due to industrial activities. Soil organic carbon stocks tend to decrease when transforming grasslands, forest or other native ecosystems to croplands and to increase when restoring native vegetation on former croplands or by restoring organic soils to their native condition. Permanent and traditional woody cultivation such olive tree and grapevine may compensate this trend due to their positive contribution of their carbon uptake.

Human society will have to rely on an increase of output per unit area in agriculture and forestry. Intensively used agricultural systems are often N-saturated and the augmented use of fertiliz-

er increases the leaching of N into aquifers and aquatic ecosystems and thus carries costs to environmental services such as water quality. In summer irrigated crops conditions are propitious for high N₂O losses. Emission factors for N₂O, distinguishing the effects of water management, crop type, and fertilizer management. Mediterranean agricultural soils produce large CH₄ emissions in flooded crops (e.g., rice) through methanogenesis, representing 6% of all CH₄ production from agricultural sources (Section 3.2.3.2).

During the past two decades, rural areas were reshaped by technological improvements in resources exploitation, the accelerating abandonment of traditional rural life and an increase in the mobility of individuals (Pinilla et al. 2008; Domon 2011). This pathway has led in many developed countries to a particular land-cover change pattern that consists in low plains and coastal areas that are being increasingly utilised for human activities due to their higher potential for agricultural productivity, while mountain or marginal areas are being abandoned because no more economically viable for production (Statuto et al. 2016; Nori 2018). Considering the increasing demand of food for an ever-growing population, leading to an increased productivity and intensification efforts in producing areas (Phelps and Kaplan 2017), this pattern has noticeable effects on the patchwork alternating semi-natural habitats that characterized the Mediterranean environment and the relevant natural ecosystem services. Management practices of grassland in hilly Mediterranean areas were progressively reduced because of reduction of animal grazing and abandonment. This produced remarkable effects on floristic simplification, loss of biodiversity, reduction of habitat for wildlife or to the survivor itself of the resource due to shrubs encroaching (Papanastasis 2004; Argenti et al. 2011). Perennial cultivation, such as olive tree grove, originally planted in marginal areas on terraces and representing a quite stable ecosystem managed with few chemical inputs, were replaced by intensive modern plantations managed under an intensive and highly mechanized system. This intensification resulted into a progressive abandonment of marginal areas because of their low economic viability leading to a shift of land use to pasture for sheep and goats, since the land is not suitable for any other kind of cultivation (Loumou and Giourga 2003).

Conversely, the intensification process towards highly mechanized and high-density plantation is boosting farmers' income but it is also causing some

environmental drawbacks. The combined effect of a more intensive management and cultivation extension is rising the issue of contamination by excess in the use of synthetic fertilizers and other agrochemicals to increase land productivity (Beaufoy and Pienkowski 2000; Beaufoy 2001). Intensification results into a degradation of habitats and landscapes and the exploitation of scarce water resources, thus putting the naturally scarce resources of olive growing areas to an edge. Drip irrigation, associated to an improved efficiency of irrigation, often has in fact no effect on efficiency but increases global water consumption. The use of this technique decreases the need for human power, allowing the increase of cultivation area and fostering multiple cropping (Kuper et al. 2017; Molle and Tanouti 2017). Soil erosion is also a growing issue due to the widening of cultivated area, as in tree crops a significant fraction of the soil is vulnerable to the action of rainfall and runoff. The intensification of agricultural land use has therefore raised the question of the long-term sustainability of agroecosystems (Liebig et al. 2004).

Different land uses as a single driver of change

Demography, technology, socio-economic factors and climate change have gradually transformed Mediterranean landscapes. Land uses are in fact the combination between the use of natural and food resources, constructions, road networks, etc., and land use changes result in the balance of their consequences. Natural habitats as affected by human activities are consequently reduced in size and continuity finally resulting into a loss of connectivity, i.e., the capability of the landscape to help or to prevent movements of organisms across habitat components (Taylor et al. 1993). A reduced or lack of connectivity may have consequences on biodiversity through losses of ecological fluxes between habitat patches and therefore trends in connectivity across different landscapes should be evaluated to consider proper actions to counteract potentially negative impact of human pressure on animal or vegetal biodiversity (Hernández et al. 2015). Over the Mediterranean Basin, two main trends, reforestation following the abandonment of agricultural areas in hilly and mountainous areas and the relevant expansion of agricultural and artificial areas in the coastal areas, plains and valleys, had different impacts on connectivity. In particular, reforestation after agricultural abandonment was correlated to a slight improvement in the connectivity of the European forest in the period 1990-2000 (Saura et al. 2011). At the same time, the spread of monoculture in

the plain resulted in a de-fragmentation process. Both these trends produced a simplification and homogenization of the landscape, in terms of number, dimensions and typology of the patches that shape the agro-ecological territory.

2.4.1.3 Future projections

Land use change is expected to have different consequences on the productivity of several ecosystems and the carbon balance. The expected warmer and dryer conditions on southern Mediterranean Basin will likely shift crop cultivation to North, where water deficit is projected to be less harsh (Ceglar et al. 2019). Whilst the less adaptable crop systems will likely suffer changed pedological and climatic conditions, the most resilient crops, such as olive tree and grapevine may have the potential to resist to this trend due to their high adaptability to cope with high temperatures and water scarcity. Moriondo et al. (2013) and Hannah et al. (2013) predicted a gradual northward shift of this cultivation in the medium term (Tanasijevic et al. 2014). Increasing temperature joint with the expected reduce rainfall rate may lead detrimental consequences for those cultivations such as corn, rice and spring wheat requiring wetter conditions. Some crops may be replaced with more resilient crops such as barley, sorghum and hay, which, however, may not well fit with the market demand and the production chain of the Mediterranean area. For most of the main Mediterranean vegetable and cereals, a decrease in production is expected in the absence of specific high input agronomic strategies such as fertilization and irrigation (Bregaglio et al. 2017; Ruiz-Ramos et al. 2018; Brilli et al. 2019) whose requirement is expected to increase (Tanasijevic et al. 2014; Saadi et al. 2015).

Climate change impacts are expected to also affect managed forests, leading to shift in typical forest communities to higher altitudes (Gitay et al. 2001). These impacts will likely affect the whole woody spinnet, influencing timber extraction and plantations, and management practices. These latter practices (e.g., fuelwood collection, forest grazing, and road expansion) can degrade the forest ecosystem conditions, particularly when applied over new forest area. Accordingly, depredated soil and forestry systems may indirectly favor the introduction of pests and pathogens, changing fire-fuel loads, changing patterns and frequency of ignition sources, and changing local meteorological conditions (Nepstad et al. 1999). Grassland and pastures will likely experience a further decrease in extension due to a progressive

rural abandonment and emigration to urban areas, often associated with low-income level in mountain areas and lack of job opportunity (Sturaro et al. 2013), and the impacts of climate change. This latter will particularly impact natural pastures, which are acknowledged to be very sensitive and vulnerable to climate conditions.

The predicted climate warming is also expected to lead changes of grasslands structure and composition, increasing the soil-water competition with trees that will be found to place at higher altitude as effect of higher temperatures. In temperate climate, warming may lengthen the forage growing season but decrease forage quality, with important variations due to rainfall changes (Craine et al. 2010; Hatfield et al. 2011; Izaurrealde et al. 2011), whilst Mediterranean pastures will likely show production decrease due to prolonged drought conditions. Modern agricultural practice can partly overcome expected production due to changed cultivation areas, but plant adaptation would require more time to be able to adapt to the new climate. Also, higher input needed to cope with changed agronomic conditions can lead to wrong perspective of the crop, forest and grassland production trend as well as extensive environmental damage. More specifically, a larger use of fertilizer or other high inputs may result in short-term increases in food production for long-term losses in ecosystem services, such as water quality degradation (Zalidis et al. 2002; Malagó et al. 2019), soil erosion, reduced fertility, or overgrazing (Wood et al. 2000). All these changes are expected to affect the carbon balance.

Land use is an important control of carbon storage; therefore, ecosystem shifts and harsher climatic conditions may lead to different forms of stress (i.e., water and nutrient stress, pedological stress, climatic stress, abiotic, etc.) which, in turn, reduce the potentiality of the different ecosystems in terms of carbon storage (i.e., biomass reduction, less growth and development). All these changes can have negative consequences in the perspective of CO₂ mitigation capacity (Foley et al. 2005), since it is expected a decrease of total carbon sequestration capacity from agro and forestry systems, increased carbon fluxes from soils due to quicker decomposition process and lower carbon mineralized in soil.

2.4.2 Sea use changes

Fisheries (over) exploitation is the main driver of marine population decline and has led to the bad

state of most highly commercial stocks and the low abundance of top predators. Climate change and variability may be responsible for catch fluctuations of some stocks (especially the small pelagic fishes), for distribution shifts but also for altering catch composition in favour of warm-water species. Recent theory predicts fish size decreases in response to increased sea surface temperature and low oxygen supply. Excessive exploitation will certainly lead to even lower stock biomasses, especially for top predators. Further sea warming will very likely lead to a higher percentage of warm-water species in the catch and smaller fish sizes.

2.4.2.1 Trends in fisheries exploitation

In the Mediterranean Sea, which together with the Black Sea, constitutes FAO Major Fishing Area 37, fishing has been practiced since antiquity. Today, Mediterranean fisheries are diverse among areas and the fishing vessels and techniques vary geographically as a result of different environmental, oceanographic, biological, climatic, cultural and socio-economic conditions prevailing in each area (Papaconstantinou and Farrugio 2000), with a strong contrast between the northern and southern coastlines. The high number of islands, ports and shelters across the Mediterranean and the contrast between north and south renders the enforcement of fisheries regulations and management very difficult. Mediterranean fisheries are highly multispecies in nature targeting over 200 fish and invertebrate species (Dimarchopoulou et al. 2017) and are operated through a large number of small sized and low tonnage fishing vessels with no large industrial fleets (Stergiou et al. 2016). The number of small-scale coastal vessels operating in Mediterranean EU waters is about 86% of the total (around 72600 vessels) with the remaining 9% being trawlers and 5% being purse seiners (Colloca et al. 2017). Although the number of all types of Mediterranean EU fishing vessels declined since 1991, the actual fishing effort has been increasing due to new technologies and higher capacity vessels (Colloca et al. 2017).

Small pelagic fisheries operate all year round but in many Mediterranean Sea subareas they show a strong seasonality that is reflected upon their catches and is derived from fishing regulations and consumer habits. According to the monthly distribution of landings and fishing effort, the main fishing season in most areas is concentrated in spring and summer months (Lloret et al. 2004b).

2.4.2.2 Current status of marine fisheries resources

In the Mediterranean Sea, recent publications based on scientific surveys, stock assessments and catch data, generally agree that the majority of Mediterranean fisheries stocks are declining in biomass as a result of their overexploitation (Colloca et al. 2013, 2017; Vasilakopoulos et al. 2014; Tsikliras et al. 2015). Local reports also confirm the bad status of Mediterranean fisheries, e.g., in Greek seas (Tsikliras et al. 2013b) and in the Ligurian Sea (Abella et al. 2010), often attributed to inadequate management practices (Tsikliras 2014; Cardinale et al. 2017). The long-lasting overexploitation of the Mediterranean Sea has been driving the decline in biomass of most commercial fish and invertebrate stocks across the basin and the near depletion of several of them (Froese and Kesner-Reyes 2002; Vasilakopoulos et al. 2014; Osio et al. 2015; Tsikliras et al. 2015; Stergiou et al. 2016; Colloca et al. 2017; Froese et al. 2018). The overall stock status is rather uniform across the Mediterranean ecoregions with low stock biomass being the common characteristic. However, the stock specific biomass levels vary among ecoregions (Froese et al. 2018).

The catch history of Mediterranean Sea stocks unmasked the overexploitation of many stocks since the 1950s, when about 40% of them were declining in biomass (Froese and Kesner-Reyes 2002). Recent literature reveals that fisheries overexploitation occurs across the entire area (Tsikliras et al. 2013a) and locally, e.g., in Greek Seas (Tsikliras et al. 2013b). Several Mediterranean stocks have been reported overfished based on data from landings (Tsikliras et al. 2013b, 2013a), scientific surveys (Stergiou and Tsikliras 2011), or stock assessments (Colloca et al. 2013). Other studies confirm that almost all species targeted by the fishing fleets are being overexploited in the Mediterranean Sea (Cardinale et al. 2017; Fernandes et al. 2017).

Based on the catch-based method, the cumulative percentage of collapsed and overexploited stocks appeared to exceed 60% across the Mediterranean Sea in 2010 (Tsikliras et al. 2013a) with the exploitation pattern differing among the Mediterranean subareas (Tsikliras et al. 2015). The western Mediterranean has been reported to be in a better state with less overexploited and collapsed stocks and more developing ones compared to the central and eastern parts of the sea (Stergiou et al. 2016). Similarly, based on various fisheries indicators, the western and central Mediterranean are in

better condition compared to the eastern part of the sea (Tsikliras et al. 2015). According to the official stocks assessments that were then available, the percentage of overexploited stocks exceeds 90% in most areas (Colloca et al. 2013) and even reaching 95% in some (Osio et al. 2015). The model approach of Osio et al. (2015) estimates that 98% of the unassessed demersal fish species are potentially overexploited in most areas. Cardinale et al. (2017) reported that the stocks of all target species that have been assessed are overexploited with the average ratio of F/F_{MSY} (actual fishing mortality to the level that would provide maximum sustainable yield) ranging from 1.7 (giant red shrimp *Aristaeomorpha foliacea*) to 8.1 (hake *Merluccius merluccius*). Steadily increasing exploitation rates and deteriorating gear selectivity have been recently reported as two conditions that lead to shrinking fish stocks (Vasilakopoulos et al. 2014). The most recent assessment of 169 Mediterranean stocks showed that 126 of them (75%) were subject to ongoing overfishing (Froese et al. 2018).

2.4.2.3 The future of marine resources

The Gill-Oxygen Limitation Theory (GOLT) predicts a reduction in the size of fish due to their inability to compensate, via their gill surface, for the increased metabolic rate that results from higher temperatures. Fish individuals that survive are expected to shrink in size (Cheung et al. 2013a). The Mediterranean Sea is among the semi-enclosed areas where local species extinctions and range shifts were predicted to be most common (Cheung et al. 2009). The GOLT theory may also explain the poleward shift of marine organisms (Cheung et al. 2013b) and their expansion to deeper waters (Perry et al. 2005) both of which occur in the Mediterranean Sea (Tsikliras and Stergiou 2014) and may have an impact on Mediterranean fisheries in terms of catch and revenue (Cheung et al. 2010).

Besides fish distribution shifts and declines in local fish stocks, scientific projections suggest that marine resources and biodiversity will suffer increasing stress if temperatures are not held below 2°C above preindustrial levels (Gattuso et al. 2015). Sea warming and deoxygenation combined with fishing pressure and other stresses could affect growth, and distribution of fish populations, resulting in changes in the potential yield of exploited marine species and economic losses (Sumaila et al. 2011) as fisheries are expected to decline (Cheung et al. 2010). Reaching the goals of the UNFCCC Paris Agreement would benefit ocean life and economies by protecting millions of metric

tons of high valued catch with 75% of maritime countries benefiting from this protection (Sumaila et al. 2019).

Fisheries in the Mediterranean will not be sustainable in the future unless the marine exploited populations are fished less, i.e., if they are allowed to recover and rebuild their biomass (Pauly and Zeller 2016) through the reduction of the

fishing pressure that is applied upon them (Froese et al. 2018). Ecosystem-based approach has also an important role that will ensure that both higher and the lower trophic levels are rebuilding (Pikitch et al. 2004) and fully marine protected areas are a key management tool to accomplish rebuilding of the biomass of marine populations, ensure ecosystem health and resilience against sea warming (Roberts et al. 2017).

2.5 Non-indigenous species

The human-aided introduction of non-indigenous species into new biogeographic regions – has been one of the main increasing global drivers of ecological change for over a millennium (Elton 1958). Non-indigenous species homogenize biodiversity across the globe, resulting in shifted and sometimes more simple ecological communities (Mooney and Hobbs 2000; Rilov and Crooks 2009). They can displace native species out of their natural habitats through competition, consumption, or parasitism. This displacement results in affected communities with reduced native species diversity, altered species composition as well as in major changes in ecosystem functioning (Vilà et al. 2011; Cameron et al. 2016). Moreover, these changes alter supporting, provisioning, regulating ecosystem services that people depend upon, impacting human well-being (Katsanevakis et al. 2014b; Vilà and Hulme 2017).

Rates of introduction and impacts have accelerated dramatically in the past few decades due to various human activities and related pathways of non-indigenous species introduction (Carlton 1989; Crooks and Suarez 2006). In the marine environment, the growth of seaborne trade with its huge fleets facilitates the dispersal of organisms attached to the hulls of ships and inside ballast water (Crooks and Suarez 2006; Katsanevakis et al. 2013). Aquaculture, live marine seafood and bait, and aquarium trade have also become important vectors for the introduction of marine non-indigenous species, as well as the artificial connections of water bodies with very different biotas (Rilov and Crooks 2009), and major catastrophes like tsunamis (Carlton et al. 2017). In terrestrial ecosystems, intentional introductions prevail over unintentional. For non-indigenous plants, ornamental and horticultural introductions escaped from cultivation account for the highest number (Lambdon et al. 2008) and are increasing steadily (Van Kleunen et al. 2018). Terrestrial non-

indigenous vertebrates follow similar patterns as plants. However, most terrestrial non-indigenous invertebrates have been introduced accidentally; many are major pests in forestry and agriculture (Roques et al. 2010).

2.5.1 Non-indigenous species in the Mediterranean Sea

The first introduction of marine species into the Mediterranean Sea dates back to the late 18th century (Poli 1791). Since then, due to maritime shipping expansion and after three centuries of accumulating human pressures, this basin has become a hotspot of introduction of non-indigenous species (Rilov and Galil 2009; Coll et al. 2010). Non-indigenous species in the Mediterranean Sea mostly arrive from the Indo-Pacific region either directly (by swimming or drifting) or indirectly as foulers or as hitchhikers inside ballast water in a process called Lessepsian migration (Por 1978). But there are many other vectors that deliver non-indigenous species into the Mediterranean and have varying importance depending on the region (Rilov and Galil 2009). Today, the total




Figure 2.25 | Number of new non-indigenous species per 6 years in the Mediterranean since 1988 (Zenetos 2019).

known number of non-indigenous species is still debated, but the last count puts it close to a thousand species (Zenetos et al. 2017). For some of the species, the impact on local biodiversity, and possibly also ecosystem functions and services, seems obvious, but in most cases, the impact is unknown because of lack of research (Katsanevakis et al. 2014b). The Mediterranean Sea is warming rapidly (Nykjaer 2009; Sisma-Ventura et al. 2014) (Section 2.2.4). Therefore, it is quite possible that the establishment of thermophilic non-indigenous species is strongly facilitated by climate change (Stachowicz et al. 2002).

2.5.1.1 Spatiotemporal trends, sources and vectors of introduction

The most recent assessment of marine non-indigenous species in the Mediterranean (November 2018) counts 957 species, including Foraminifera (Zenetos 2019). Among the introduced species during the last 30 years (491 taxa), invertebrates dominate with >58% (287 species) represented mostly by molluscs and decapods. Primary producers follow with approximately 114 species among which macroalgae, especially rhodophytes, prevail. Vertebrates (mostly fishes) follow with 90 species.

The trend in introduction of species in the Mediterranean (Fig. 2.25), which culminated in the 2000–2005 period with more than 20 new species per year (122 in total), appears to be overall decreasing after 2005 (Fig. 2.26). As opposed to invertebrates and primary producers, vertebrates continue increasing, with 34 species detected in the 2012–2017 period vs 26 species in the period 2006–2011. The overall decreasing rate in non-indigenous species is evident across the Mediterranean Marine Strategy Framework Directive (MSFD) areas, except for the Central Mediterranean where an increase is observed, attributed to vertebrates (Fig. 2.26). 17 new fish species were detected in the central Mediterranean in the period 2012–2017 vs. 8 fish species in the period 2006–2011. These species are either spreading from the eastern Mediterranean to the central region or are newly introduced species in the area. Vertebrates (fish only) are dominated by Lessepsian immigrants but over the last decade the number of fish species related to aquarium trade which have been intentionally released to the wild (classified as escapees from confinement) is increasing (Zenetos et al. 2016; Marcelli et al. 2017; Deidun 2018).

Regarding the spatial distribution of non-indigenous species, the number of Lessepsian species is very high on the eastern Mediterranean coastline, reaching 129 species per 100 km², and declines toward the north and west (Katsanevakis et al. 2014a). The distribution of species introduced by shipping is strikingly different, with several hot-spot areas occurring throughout the Mediterranean Basin. Two main hotspots for aquaculture-introduced species have been identified (the Thau and Venice lagoons). Certain taxonomic groups were mostly introduced through specific pathways-fish through sea corridors, macrophytes by aquaculture, and invertebrates through sea corridors and by shipping (Katsanevakis et al. 2014a). Hence, the local taxonomic identity of the non-indigenous species is greatly dependent on the dominant maritime activities/interventions and the related pathways of introduction. The composition of non-indigenous species assemblages differs among Mediterranean ecoregions; such differences are greater for Lessepsian and aquaculture-introduced species.

2.5.1.2 Non-indigenous species as drivers of biodiversity and ecosystem change

The introduction of non-indigenous species in the Mediterranean Sea have caused modifications in biodiversity patterns. One of the best documented and most profound impacts of the introduction of non-indigenous species on native Mediterranean ecosystems is the deforestation of algal forests and the creation of extent barrens (i.e., areas with bare rock and encrusting calcified algae; Fig. 2.27) by the overgrazing activity of two non-indigenous herbivore rabbitfishes: *Siganus luridus* and *S. rivulatus*. These species have become dominant in the ichthyofauna of shallow rocky habitats in the eastern Mediterranean, and have caused ecosystem-wide changes by creating and maintaining areas denuded of canopy algae (an important habitat for many coastal fishes) (Cheminée et al. 2013). This form of "deforestation" is associated with a dramatic reduction in biodiversity, biomass, and algal growth, and effects that move up the food chain to the local fisheries (Sala et al. 2011; Vergés et al. 2014b).

In the Levantine Sea, the catch of commercial fisheries is now dominated by non-indigenous species (Edelist et al. 2013; Katsanevakis et al. 2018), reflecting the decline of native biota and its replacement by thermophilic non-indigenous species (Arndt et al. 2018). In Turkish coastal waters in the Levantine Sea, non-indigenous




Figure 2.26 | Change in the number of new non-indigenous species per 6 years in the Mediterranean MSFD regions (Adriatic, Western Mediterranean, Central Mediterranean, Eastern Mediterranean Seas) since 1988. Notice that the scale is different for the eastern Mediterranean (Zenetos 2019).

fish species biomass exceeds 50% of the total fish biomass and 90% of the total herbivore fish biomass (Giakoumi et al. 2019). Fishers in the same area (Kaş, Turkey) perceived the introduction of non-indigenous species as the most important reason for the current fish stock depletion.

Yet, it is uncertain whether the decline of native biota in the Levantine Sea is driven mainly by biotic interactions with non-indigenous species or climate change, or both, as the Lessepsian species are thermophilic and their establishment is assisted by climate change (see discussion below). A recent meta-analysis suggests that the decline of native fish can be mostly attributed to ocean warming and not to negative interactions with non-indigenous species (Givan et al. 2017a). Multi-species collapses of native species are also at least partially attributed to climate change (Rilov 2016), with the collapse of the sea urchin *Paracentrotus lividus* experimentally demonstrated to be related to the fast ocean warming (Yeruham et al. 2015).

Analysis focused on shallow reef fish shows that non-indigenous species are very diverse ecologically, and they considerably increase the total com-

munity trait diversity of the Mediterranean (Givan et al. 2017b). Furthermore, trait similarity between non-indigenous and indigenous Mediterranean species was lower than expected, indicating that non-indigenous fish tend to occupy relatively vacant niches within the Mediterranean. Temporally, non-indigenous fish species display increased trait similarity to native Mediterranean species, suggesting that forecasting future establishment may be challenging. Givan et al. (2017b) conclude that the Mediterranean, at least in fish, is transforming into an extension of the Red Sea in terms of trait and species composition. Such biological trait analysis is also required for other taxonomic groups.

There is a serious research gap in assessing and quantifying the impacts of non-indigenous species on marine ecosystems in the Mediterranean Sea. Impact assessment is mostly based on expert judgment or correlational studies, while manipulative or natural experiments are largely lacking for assessing the impacts of most non-indigenous species in the region (Katsanevakis et al. 2014b). Disentangling the role of the introduction of non-indigenous species and climate change or other local or global stressors to derive cause-



Figure 2.27 | Ecosystem shift from algal forests to barrens due to the overgrazing activity of non-indigenous herbivore rabbitfish. Algal forests host high fish, invertebrate, and algae biodiversity whereas barrens are associated with low levels of biodiversity across all taxonomic groups. *Rabbit fish photo: Murat Draman.*

effect pathways is inherently difficult and would probably necessitate a combination of experimental and modelling approaches.

Efforts are made to assess non-indigenous species impacts with existing knowledge. Based on a conservative additive model, which downgrades reported impacts of low inferential strength, an index of the Cumulative IMPacts of invasive ALien species (CIMPAL) on marine habitats in the Mediterranean has been developed and estimated (Katsanevakis et al. 2016). The estimation of CIMPAL was based on assessments of impacts for every combination of 60 non-indigenous species and 13 habitats, and their distributions in the Mediterranean (i.e., presence in 10x10 km cells). It showed strong spatial heterogeneity in impacts. Spatial patterns varied depending on the pathway of introduction of the non-indigenous species in the Mediterranean Sea. Species introduced by shipping gave the highest impact scores and impacted a much larger area than those introduced by aquaculture and through sea corridors. Overall, non-indigenous macroalgae had the highest impact among all taxonomic groups, when estimated as the sum of impact scores across the entire Mediterranean Sea, i.e., accounting not only for the severity of the impacts but also their spatial extent. The most impactful non-indigenous species was *Caulerpa cylindracea*, which has become dominant over large areas of shallow waters in the western Mediterranean and compete with native species (Piazzi et al. 2016). Negative impacts of *C. cylindracea* have been documented on al-

gal, sea grass, and sponge communities (Ceccherelli et al. 2002; Piazzi et al. 2005; Piazzi and Ceccherelli 2006; Baldacconi and Corriero 2009).

2.5.1.3 Further introductions, monitoring and managing non-indigenous species

The introduction of species in the Mediterranean Sea is a continuous process and it is very likely that it will continue for years to come. In most of the cases, these species fail to thrive but, evidently, some become numerically and ecologically dominant in their new environment, generating new and sometimes severe impacts of the introductions. As a consequence, in the past two decades research interests in non-indigenous species have increased, mostly stimulated by evidence on their ecological and socio-economic impacts in the Mediterranean region. This has also raised the urgency of innovative approaches to forecast, track and manage these species (Corrales et al. 2018).

One of the most recent and potentially damaging non-indigenous species for this basin is the common lionfish (*Pterois miles*), which increasingly appears in many parts of the eastern Mediterranean in the last few years (Bariche et al. 2013). Due to its rapid increase in abundance (Kletou et al. 2016) and fast geographical expansion (Azzurro et al. 2017), this harmful species has become emblematic for raising concern on Mediterranean non-indigenous species introductions but it also well illustrates a process of developing monitoring capabilities and

management strategies within the Mediterranean region.

Documenting the spread of this non-indigenous species can greatly benefit by the participation of resource users, a partnership which can support monitoring objectives (Azzurro and Bariche 2017) and be used to reduce, at least locally, the abundance of non-indigenous species (Kleitou et al. 2019). More generally, participatory approaches benefit the scientific consensus, which is a key element for both documenting and responding to these introductions (Scyphers et al. 2015). This is also reflected in some of the guiding principles on the management of non-indigenous species adopted by key regional and international bodies/legislative frameworks, concerning non-indigenous species, such as those provided by the EU (Regulation 1143/2014); UNEP-MAP and by FAO-GFCM, which are converging towards finding common strategies to face Mediterranean species introductions. In this regard, the Integrated Monitoring and Assessment Programme and related Assessment Criteria (IMAP) adopted through Decision IG.22/7 by the 19th Ordinary Meeting (COP 19, Athens, Greece, 9-12 February 2016) of the Contracting Parties to the Barcelona Convention, stress the need of comprehensive monitoring and coordinated transnational actions to face the common issue of Mediterranean non-indigenous species introductions. A recent expert assessment of management options of marine non-indigenous species prioritized 11 management actions for controlling 12 model species according to their dispersion capacity, distribution, and taxonomic identity (Giakoumi et al. 2019). The actions were assessed using five criteria (effectiveness, feasibility, acceptability, impacts on native communities, and cost), combined in an "applicability" metric. Raising public awareness and encouraging the commercial use of non-indigenous species gained the highest priority, and biological control was considered the least applicable (Giakoumi et al. 2019).

To predict future change in the distribution of native species and the spread of non-indigenous species, species distribution models (SDM) are regularly used. In these models, climate matching is calculated between the area of origin (donor) and the area of potential spread (recipient). However, a recent study that matched the native range of Red Sea fish and their new range in the Mediterranean Sea showed poor matching, and thus indicated that SDMs may underestimate the potential spread of non-indigenous species (Parravicini et

al. 2015). The authors call for caution in employing such models for forecasting the introduction of non-indigenous species and their response to environmental change, as uncertainty is large. Better knowledge of the fundamental niche of species (their physiological performance under different environmental conditions) can potentially improve the prediction of spread of non-indigenous species in the Mediterranean. Furthermore, an analysis combining ship movements with port environmental conditions and biogeography can be used to quantify the probability of new primary introductions through ballast water (Seebens et al. 2013).

2.5.2 Terrestrial non-indigenous species and pests

2.5.2.1 Spatial patterns and temporal trends

Degree of introduction across Mediterranean-type ecosystems and geographical areas

The information available on terrestrial non-native species in the Mediterranean Basin countries is not comprehensive, and the number of non-indigenous species is underestimated due to the incompleteness of collected data and the monitoring bias towards some taxonomic groups. For example, Abellán et al. (2016) reported more than 370 non-indigenous birds for Spain and Portugal, more than twice the number listed by DAISIE (2009) for the whole Europe.

Most non-indigenous species in the Mediterranean Basin are plants, followed by invertebrates (Fig. 2.28). Natural habitats in the Mediterranean Basin host more than 400 non-indigenous plant species (Arianoutsou et al. 2013). The taxonomic similarity of the non-native flora among Mediterranean Basin countries is very low. For example, less than 30 species are common across 4 Mediterranean European countries (Arianoutsou et al. 2010), and only 10 species are common in 8 major Mediterranean islands (Lloret et al. 2004a). However, non-native plants share similar traits, as shown by the high proportion of perennial herbs with a long flowering period, which are pollinated and dispersed by the wind (Lloret et al. 2004a, 2005).

The major part of non-indigenous invertebrate species are arthropods, especially insects (Roques 2010), with a low representation of nematodes and flatworms (Naves et al. 2016; Justine et al. 2018). Phytophagous pest species are largely dominating among non-indigenous species all




Figure 2.28 | Number and proportion of terrestrial non-indigenous plant, invertebrate and vertebrate species per country (EASIN)⁸.

over the Mediterranean Basin, accounting for more than a half of the invertebrate species and about one-third are associated with woody plants (Matošević and Pajač Živković 2013; Roques 2015; Avtzis et al. 2017). Among them, Hemipterans, mostly scales and aphids, constitute the dominant group, accounting for 40-75% of the non-indigenous species in any part of the Mediterranean Basin (Roll et al. 2007; Inghilesi et al. 2013; Matošević and Pajač Živković 2013; Seljak 2013; Avtzis et al. 2017). This over-representation of Hemiptera in non-indigenous species seems related to their small size and easier of transportation with infested imported plants.

A complete plant pathogens database is not available. However, a European database of non-indigenous forest and woody pathogens indicate a list of 123 plant pathogens. In Mediterranean countries, Ascomycota is the most numerous group, while Oomycota and Basidiomycota represent 21% and 9% of the total, respectively (Santini et al. 2013). With regard to vertebrates, most non-indigenous species are birds, followed by mammals, while the number of reptiles and amphibians is low.

The impact of non-indigenous species varies largely among countries. Countries with higher Human Development Indexes and imports host a large density of non-native plants (Vilà and Pujadas 2001). Between countries and within countries, the density of non-native plants is related to the length of terrestrial transport networks. Areas with extensive road and rail networks, high anthropogenic disturbance, low altitude, short distance to the coastline and dry, hot climate show higher richness of non-indigenous plant species (Gassó et al. 2012). Most affected landscapes are those highly urbanized and with high population densities (Sobrino et al. 2002; González-Moreno et al. 2013). Accordingly, the most affected ecosystems by non-indigenous plants are human-modified such as ruderal, waysides or agricultural fields (Vilà et al. 2007; Hulme et al. 2008; Arianoutsou et al. 2010).

As for other non-indigenous taxa, most pests are introduced in managed habitats, such as agricultural lands and parks and gardens, forests being less affected (Matošević and Pajač Živković 2013). Among the infested plant species, non-native ornamental plants (palms, legume trees),

⁸ <https://easin.jrc.ec.europa.eu>

Citrus and *Eucalyptus* are slightly more colonized than native species (Roques 2015). Vertebrates also tend to occupy anthropic habitats and, to a lower extent, woodlands (at least for birds) (Kark et al. 2009). The degree of introduction of non-indigenous invertebrate species also varies among countries, Italy and France showing much more established non-indigenous species than any other European country (Roques 2010). The same trend is reported also for plant pathogens (Santini et al. 2013).

Mediterranean islands and islets host a large number of non-indigenous species, mainly plants (Brundu 2013). For example, an analysis of 37 small Italian islands showed that they are affected by 203 non-native plants, with a remarkable increase of acacias and succulents in the last decades (Celesti-Grapow et al. 2016). The main determinants of non-indigenous plant species richness in small islands are tourist development and the percentage of artificial land-cover. However, at the local scale, para-oceanic island ecosystems such as the Balearic Islands have a relatively lower number of non-native plants than their mainland counterparts (Vilà et al. 2010). Yet, some species can be introduced into more ecosystem types in islands than in the mainland (Gimeno et al. 2006). In general, there are large differences in the taxonomic composition of non-indigenous insect assemblages between islands and continental countries, e.g., France and Corsica, Italy and Sicily (Liebhold et al. 2016), and Greece and Crete (Avtzis et al. 2017).

Temporal trends of non-indigenous species and pests

The rate at which humans have moved species beyond their native ranges has tremendously increased over the last 150–200 years (di Castri 1991; Reichard and Hamilton 1997), and more so in the last decades (Genovesi et al. 2009; Cardador et al. 2019). Although all taxonomic groups have shown a general rise during this period consistent with the exponential increase in trade and travel (Jeanmonod et al. 2011; Seebens et al. 2017; Cardador et al. 2019), little is known about how temporal dynamics of non-indigenous species varies among taxa. Where time series are available, the number of non-indigenous species established in Europe has increased exponentially in terrestrial ecosystems (Jeschke and Strayer 2005; Hulme et al. 2008; Lambdon et al. 2008; Santini et al. 2013). Abellán et al. (2016) analyzed data on bird introductions in Spain and Portugal since 1912 and found that most of them (99.9%) were recorded

from 1955 onwards, with a sharp increase after the 1980s that mirrors the number of non-native birds imported into these countries. Cage birds (mainly Passeriformes and Psittaciformes) constitute the bulk of the species introduced during the last 40 years through escapes of individuals kept in captivity as pets. Although the information is less detailed, and reptiles and amphibians have smaller numbers of recorded non-indigenous species than birds, both groups have also increased their numbers during the 20th century in parallel with the rise in human immigration into Europe (Jeschke and Strayer 2006) and the international trade (Jenkins 1999).

The rate of establishment of non-indigenous insect species has also increased during the last decades (Roques 2010, 2015; Matošević and Pajač Živković 2013; Avtzis et al. 2017). A fast and quite linear increase, with about 10 new species per year, was noticed in Italy since World War II (Inghilesi et al. 2013), and an even higher rate of increase was noted in Croatia since 2007 (Matošević and Pajač Živković 2013). The species newly established during the last three decades tend to spread all over the Mediterranean Basin significantly faster than those that arrived between 1900–1990s (Roques et al. 2016). Such a rapid spread was especially impressive in some species, often relying on multiple introductions in different countries being used as bridgeheads (Rugman-Jones et al. 2013; Kerdelhué et al. 2014; Garnas et al. 2016; Roques et al. 2016; Bras et al. 2019; Lesieur et al. 2019).

Non-indigenous plant pathogenic species have increased exponentially in the last four decades (Santini et al. 2013). Since then, new non-indigenous plant pathogenic species have been introduced mainly from North America, and recently from Asia. Hybrid pathogens also appeared. Countries with a wider range of environments, higher human disturbances or international trade host more non-indigenous species. Rainfall influences the diffusion rates. Environmental conditions of the new and original ranges and systematic and ecological attributes affect pathogen success (Santini et al. 2013).

For plants, the success of introduction in terms of their area of occupancy is larger in species introduced a few centuries ago than species introduced in the 20th century (Lambdon and Hulme 2006), while for birds, establishment success is positively related to time since first introduction (Abellán et al. 2017).

Pathways of introduction (intentional and accidental) of non-indigenous species and pests

The majority of non-indigenous plants have been introduced into the Mediterranean Basin intentionally, as ornamentals that have escaped from gardens associated with anthropic developments and housing (e.g., touristic urbanizations) but also to embellish infrastructures (Hulme et al. 2008). Furthermore, many non-native trees (e.g., *Acacia*, *Pinus*, *Eucalyptus*) have been planted at large scales as forestry species and also in restoration programs for dune-stabilization, riverine water flux control, soil fertilization or afforestation of agricultural abandoned land. Many plant species have also been introduced unintentionally (accidentally) as "hitchhikers" or seed contaminants.

The main pathway of introduction for vertebrates, for example birds, are accidental escapes from private collections (Abellán et al. 2016). International wildlife trade is one of the main (if not the main) sources of current vertebrate non-indigenous species. When the EU banned the imports of wild birds, there was a rapid trade shift from wild-caught birds to captive-bred birds (which have lower potential to establish populations than wild-caught birds) (Carrete and Tella 2008, 2015; Cabezas et al. 2013) and a sharp decrease in the number of new introduced avian species in the wild (Cardador et al. 2019). However, this positive effect of the EU ban on wild-caught birds coincides with a significant increase in the trade in reptiles (Cardador et al. 2019).

For invertebrates, the vast majority of species introductions have been accidental (Hulme 2009; Roques 2015). A few introductions have been intentional, mostly for biological control between 1950 and 1999 (Rasplus et al. 2010), but such species always represent less than 15% of the total number of non-indigenous species per Mediterranean country, except in Israel (17.4%) (Roll et al. 2007). Since the majority of invertebrates established in the Mediterranean Basin are phytophagous, the major pathway of unintentional introductions appears to be via international trade in live plants (Rabitsch 2010; Inghilesi et al. 2013; Eschen et al. 2015; Roques 2015). Seed trade has also provided a few species and pests (Auger-Rozenberg and Boivin 2016) as well as firewood, logs and fallen timber (Meurisse et al. 2019). The trade of vegetable and fruit commodities also constitute an important pathway for non-indigenous pests (Desneux et al. 2010; Abbes et al. 2012; Cini et al. 2014). Hitchhiking is another significant pathway of pests as stowaways using wood packaging material

(Rassati et al. 2015; Javal et al. 2019; Lesieur et al. 2019), transport infrastructure and vehicles (Javal et al. 2019; Kirichenko et al. 2019) or used tires such as for mosquitoes (Rabitsch 2010). Several other examples are associated with beekeeping (Mutinelli et al. 2014).

For plant pathogens, all the introductions occurred unintentionally. The exact pathway of introduction is almost unknown for most species. However, the most probable is the trade of living plants (57%) or wood (10%). Less than 10% of the introductions occurred through any of the other pathways (Santini et al. 2013). Introductions of even harmless fungi in a new environment give them the opportunity of mating with local or introduced related species giving rise to hybrid progenies. The hybridization process may result in an increase of pathogenicity in one of the species or in the emergence of a completely new plant disease, both of which may threaten the original host plant, and new and naïve host species (Ghelardini et al. 2017).

2.5.2.2 Non-indigenous species as drivers of biodiversity and ecosystem change

The introduction of non-native plants can decrease local flora and fauna diversity and change the community composition and functional structure of affected ecosystems (Vilà et al. 2006; Zahn et al. 2009; Rascher et al. 2011). Native plants that are most vulnerable to such introductions are those with small population sizes (Lapiedra et al. 2015). At least 12 endemic or critically endangered plant species from the "Top 50 Mediterranean Island Plant" list are threatened by non-native plants (de Montmollin and Strahm 2005). For birds, the impact of non-indigenous species is higher on island species, and in those species with small distribution ranges (Clavero et al. 2009). Destabilized ecosystems, including systems used for food and agricultural production, tend to be more vulnerable to the spread of non-indigenous species (e.g., Marvier et al. 2004; Chytrý et al. 2008). However, there is little evidence to support the hypothesis that highly diverse ecosystems are inherently more resistant to non-indigenous species than less-diverse systems (e.g., Keller et al. 2011).

Changes in ecosystem functioning after the introduction of non-indigenous plants are highly context-dependent and include alterations in decomposition rates, light and water soil availability, and changes in soil carbon and nitrogen pools (Vilà et al. 2006; Castro-Díez et al. 2009; Rascher et al.




Figure 2.29 | Number of non-indigenous woody plant pathogens per country (Santini et al. 2013).

2011). Plant introductions can disrupt the positive relationship between native species diversity and multifunctionality (Constán-Nava et al. 2015) and may trigger regime shifts by changing plant succession (Stinca et al. 2015).

The impacts of vertebrates can be dramatic through competition for resources, predation and as vectors of diseases (Genovesi et al. 2009). For example, the introduction of non-native ungulates is a major threat to endangered plants, especially in islands (Pisanu et al. 2012). Some native amphibians have collapsed after the introduction of non-native anurans (Lillo et al. 2011). Non-indigenous parakeets can also interact with native species and have an impact on native populations and communities, largely in the form of harassment, displacement from nest sites and food competition (Hernández-Brito et al. 2014, 2018; Menchetti and Mori 2014; Menchetti et al. 2016; Covas et al. 2017). NIS arthropods can also negatively impact native biodiversity and ecosystem processes by destroying host plant populations, causing disturbances in native genetic resources or, indirectly, affecting the affected communities because of new species assemblages (Kenis et al. 2009; Kenis and Branco 2010; Auger-Rozenberg and Boivin 2016).

Some non-indigenous species can damage productive sectors, most notably agriculture and forestry, with important economic consequences. Non-native insects impact major crops in many countries of the Mediterranean region (Abbes et al. 2012; Abdallah et al. 2012; Mutke et al. 2016). Some non-indigenous bird species such as Monk and ring-necked parakeets can also cause important damages to crops (Senar et al. 2016; Turbé et al. 2017).

Forests as well as urban trees in the Mediterranean Basin can be severely economically impacted by non-indigenous pathogens (Fig. 2.29) (Santini et al. 2013; Ghelardini et al. 2017) and arthropods (Kenis and Branco 2010) some of which can be considered pests (Santini et al. 2013; Rassati et al. 2015; Auger-Rozenberg and Boivin 2016; Branco et al. 2016; Mendel et al. 2016) and can also transmit pathogenic fungi (Montecchio et al. 2014). For example, after World War II, chestnut blight epidemic in the mountains of southern Europe aggravated food shortages for local human populations and increased migration to urban areas (Adua 1999). Besides, canker stain disease of the plane tree was considered a nasty non-indigenous species of urban trees, until it was introduced into Greece, where the Oriental plane is endemic. The disease is presently destroying natural river wood ecosystems in Greece (Ghelardini et al. 2017; Tsopelas et al. 2017) and it is also spreading in neighbouring countries such as Albania (Tsopelas et al. 2017) and Turkey (Lehtijärvi et al. 2018). Currently, the pinewood nematode is massively killing pines, changing the landscape in Portugal (Naves et al. 2016).

Human health can also be affected by non-indigenous species. Of major concern are non-native plants that are allergenic; their advanced flowering phenology enlarges the period of airborne prevalence of allergens (Belmonte and Vilà 2004). Non-indigenous birds of the order Psittaciformes (parrots) are potential reservoirs of *Chlamydophila psittaci*, the etiological agent of human psittacosis, and can transmit other diseases to humans and wildlife (Menchetti and Mori 2014; Turbé et al. 2017). Some non-native invertebrates, mainly insects, can cause distress, and allergic reactions

(e.g., the Asian hornet, *Vespa velutina*) or be vectors of infectious diseases, (e.g., the tiger mosquito, *Aedes albopictus*) (Lounibos 2002; Jucker and Lupi 2011; Monceau et al. 2014; Goubert et al. 2016; Roques et al. 2018; Lioy et al. 2019).

Some impacts of introduced non-indigenous species have attracted worldwide attention, e.g., on cultural heritage in Palermo (Manachini et al. 2013), on the survival of the endangered date palm, *Phoenix theophrasti* in Crete (Avtzis et al. 2017), and the severe attacks of cypress canker disease in southern Tuscany (Italy). Since researchers and policymakers rarely address the connection between non-indigenous species and damage to cultural heritage directly, the cost of these losses is often neglected or underestimated. The Mediterranean Basin has a long history of civilization and it is rich in cultural heritage that can be threatened by non-indigenous species. For example, in southern Tuscany (Italy), severe attacks of the cypress canker disease (caused by the North American fungal pathogen *Seiridium cardinale*) are threatening the survival of trees flanking a monumental avenue (Danti and Della Rocca 2017).

2.5.2.3 Further introductions, spread and impacts of non-indigenous species and pests

Existing tools for predicting the risk of introduction and research needs

Horizon scanning, prioritization and Pest Risk Analysis (PRA) are essential tools for focusing limited resources to predict the species which can have a high rate of spread, inflict high impacts, and can be cost-effectively managed. PRA are defined by the International Plant Protection Convention as "the process of evaluating biological or other scientific and economic evidence to determine whether a pest should be regulated and the strength of any phytosanitary measures to be taken against it" (FAO 2017b). An important step in the PRA scheme is the "Pest management section" which assesses phytosanitary measures for relevant pathways and their effectiveness in preventing the entry, establishment and spread of non-indigenous species.

Since 2006, European and Mediterranean Plant Protection Organization (EPPO) has formed expert working groups (EWG) to conduct PRA comprised of experts on the pest and cropping systems, mapping and modelling experts, along with experts on EPPO's PRA scheme, risk managers and EPPO PRA Core Members, all which acts to ensure

consistency. EPPO is an international organization responsible for cooperation and harmonization in plant protection within the European and Mediterranean region. One of EPPO's main aims is to provide assistance and guidance to member governments on the administrative, legislative and operational measures necessary to prevent the introduction and spread of non-native plant pests (Smith 1979; Roy et al. 2011). Since 1999, EPPO has maintained an Alert List of plant pests and non-indigenous plants which acts as an early warning for pests, which can present a risk to the EPPO region. EPPO has also developed a prioritization tool for non-indigenous plants that classifies species into one of three lists: minor concern, observation list or list of non-indigenous plant species (EPPO 2012a, 2012b). Those species included in the list of non-indigenous plants are assessed for a PRA, where a higher priority is given to those species with a limited distribution in the EPPO region (EPPO 2012b).

In addition to the EPPO PRA tools, there are also a number of other PRA protocols (Roy et al. 2018) that can be applied to the Mediterranean Basin. To better improve PRA and the risk assessment process, a greater level of transparency and consistency between protocols would be beneficial (Vanderhoeven et al. 2017; González-Moreno et al. 2019). EU Mediterranean member states need to follow the EU Regulation that includes restrictions on keeping, importing, selling, breeding and growing non-indigenous species (European Union 2014).

A quite novel empirical approach to identify potential pests is the use of sentinel plantations of Mediterranean trees, e.g., cork oak, evergreen oak and cypress, in other continents as a priori identification of non-indigenous insect and pathogens capable of colonizing such plants. For example, such plantations in China provided a list of 39 potential non-indigenous insects of which five could be highly damaging (Roques 2015) and several pathogens (Vetraino et al. 2015). The development at potential ports of entry of trapping programs using lures presenting a generic attractiveness for some insect groups is expected to allow early detection of emerging non-indigenous species, even when not listed on quarantine lists (Rassati et al. 2014, 2015; Fan et al. 2019).

Non-indigenous species likely to be introduced into the Mediterranean in the next 20-50 years

As temperature increases, current major non-indigenous species are predicted to shift north-

wards at an average pace of 37–55 km decade⁻¹, leaving a window of opportunity for new non-indigenous species better adapted to xeric conditions (Gallardo et al. 2017). Regarding non-native plants, gardening practices and ornamental trade will have a major impact on the selection of these future non-indigenous species. The use of non-native drought-tolerant species for gardening and landscaping (i.e., xeriscape) is at its earliest stages in the Mediterranean, but it has already raised concerns in California because of its potential risk as a source of new non-indigenous species (Bradley et al. 2012). Global species niche modelling indicates that xeric shrublands in Mediterranean areas are among the most susceptible ecosystems to introduction by Cactaceae plant species from arid American areas (Novoa et al. 2015). Although some Cactaceae are already largely distributed across the Mediterranean (e.g., *Opuntia ficus-indica*), it is very likely that close relatives with currently restricted distribution or absence in the Mediterranean, such as *Cylindropuntia* spp., would thrive in the next decades aided by new gardening practices (Essl and Kobler 2009). Besides gardening, other relevant terrestrial plant species could be easily introduced and established as contaminants in soil, seeds or containers. For instance, *Parthenium hysterophorus* is a species not currently in the Mediterranean that has been highlighted as of high risk for the region because of its large potential negative impact on agriculture and human health (Kriticos et al. 2015).

While plants have been proportionally the main new non-indigenous species in Europe up until the 19th century, the trend has shifted towards an increasing number of introduced invertebrates and vertebrates in the 20th century (Hulme 2009). This is a pattern that is very likely to continue in the near future by increasing air and maritime cargo, where these taxa can be easily transported as stowaways. The establishment of non-indigenous

invertebrates of tropical origin affecting woody ornamental plants has increased (Eschen et al. 2015), meaning that many ornamental plants, especially palms, fig trees and exotic legumes, are at risk for further introduction as well as *Citrus* and *Eucalyptus* trees (Floris et al. 2018). The recent establishment of ambrosia beetles of tropical origin directly threatening plants of the Mediterranean maquis (Faccoli et al. 2016; Francardi et al. 2017) suggests that such process is going to be amplified with global warming. A list of fruit flies likely to be introduced has been recently proposed (Suffert et al. 2018). Special attention should be paid to major agricultural pests currently not present in the Mediterranean but with the potential to be introduced and cause a major impact. The EPPO A1 quarantine list considers up to 128 species of insects, mites, nematodes and gastropods, currently absent from the EPPO region, recommended for quarantine measurements. For instance, Lepidoptera species such as *Spodoptera* or *Helicoverpa* spp. are polyphagous species that could easily thrive in the Mediterranean if they become established. *Spodoptera frugiperda*, a pest native to the Americas, has quickly spread in Africa causing large yield loss. A recent modelling exercise has identified small pockets of suitable habitats in the Mediterranean area but the potential for permanent populations is still uncertain (Early et al. 2018).

Regarding terrestrial vertebrates, several species have been recently highlighted in a horizon scanning exercise for European non-indigenous species, including the Mediterranean (Roy et al. 2019). Of special relevance is the common myna, *Acridotheres tristis*, a non-indigenous species with very restricted populations in the region, and *Lampropeltis* spp., a family of snakes mainly native to North America and adapted to arid conditions. These species are traded as pets and can easily be introduced due to accidental escapes.

2.6 Interaction among drivers

2.6.1 Drivers impacting other drivers

The potential for interactions among drivers is a key issue for analyzing their impacts on environment and human societies, and for developing effective conservation policies (Brook et al. 2008). Climate change, pollution, land and sea use change, and non-indigenous species are of-

ten studied and managed in isolation, although it is becoming increasingly clear that a single driver perspective is inadequate when ecosystems are threatened by multiple, co-occurring drivers (Halpern et al. 2008a, 2008b). Conceptually, there are three broad categories of interaction types describing the outcome of multiple stressors, the effects can be additive/cumulative (all the dif-

ferent stresses derived from the implied drivers show up), synergistic (increased stress) or antagonistic (decreased stress) (Folt et al. 1999; Crain et al. 2008). Also, and particularly for the Mediterranean, how different drivers interact could result in alteration, intensification, and even in generation of new impacts (Doblas-Miranda et al. 2017).

In order to facilitate the multi-stressor approach, as a key recognized concept, this section offers two different approximations. First, we describe potential pair interactions within the individual driver classes described before, and second, we provide a few examples of characteristic disturbances of Mediterranean ecosystem that are the result of the combination among multiple drivers.

2.6.2 Pairs of interacting drivers

2.6.2.1 Climate change effects on pollution

Generally, increases in temperature enhance the toxicity of contaminants and increase concentrations of tropospheric O₃ regionally, but will also likely increase rates of chemical degradation (Lelieveld et al. 2014). In general, climate change coupled with air pollutant exposures may have potentially serious adverse consequences for human health in urban and polluted regions (Noyes et al. 2009).

The increase in the intensity and frequency of storm events linked to climate change can lead to more severe episodes of chemical contamination of water bodies and surrounding watersheds (Noyes et al. 2009). Climate change may also increase the occurrence and the global expansion of harmful algal blooms (Paerl and Paul 2012) (Sections 2.3.3 and 2.3.4).

2.6.2.2 Pollution effects on climate change

Many air pollutants that are harmful to human health and ecosystems also contribute to climate change by affecting the amount of incoming sunlight that is reflected or absorbed by the atmosphere, with some pollutants warming and others cooling the Earth. These so-called short-lived climate-forcing pollutants include methane, black carbon, ground-level O₃, and sulfate aerosols. They have significant impacts on the climate; black carbon and methane in particular are among the top contributors to global warming after CO₂ (Shindell et al. 2009; Stohl et al. 2015). Over the Mediterranean Basin the increase and decrease of anthropogenic aerosols during the second half of the 20th century have had an important role in the dimming-brightening phases, because of their direct action on the incoming solar radiation (Section 2.2.3.1).

Pollution by heavy metals or organic compounds can also affect ecosystem functioning by inhibiting CO₂ fixation performed by photosynthetic organisms, thereby increasing global warming (Rochelle-Newall et al. 2008; Magnusson et al. 2010; Ben Othman et al. 2012) (Sections 2.2.3 and 2.3.2).

2.6.2.3 Impact of climate on land and sea use

Effects of climate change on land use

Recent accelerated climate change has exacerbated existing environmental problems in the Mediterranean Basin caused by the combination of changes in land use, increasing pollution and biodiversity decline (Cramer et al. 2018). Sea-level rise, combined with land subsidence, may significantly reduce the area available for agriculture. The effects of sea level rise in North Africa, especially

Impacting (column) – Impacted (row)	Climate change	Pollution	Land and sea use changes	Non-indigenous species
Climate change		2.6.2.2	2.6.2.4	?
Pollution	2.6.2.1		2.6.2.7	?
Land and water use changes	2.6.2.3	2.6.2.6		2.6.2.10
Non-indigenous species	2.6.2.5	2.6.2.8	2.6.2.9	

Table 2.3 | Main interactions among drivers

on the coast of the Delta region of Egypt, would impose additional constraints to the agricultural land (*Section 3.2.2.1*), and also the salinization of coastal aquifers (*Section 3.1.2.2*).

Similarly, with 42% of the population living in coastal areas (Mediterranean Wetlands Observatory 2018), important direct effects of climate change on coastal settlements include dry-land loss due to erosion and submergence, damage of extreme events (such as wind storms, storm surges, floods, heat extremes, and droughts) on built environments, effects on health (food- and water-borne disease), effects on energy use, effects on water availability and resources, and loss of cultural heritage (Hunt and Watkiss 2011) (*Section 2.2.8.2*). Coastal industries, their supporting infrastructure including transport (ports, roads, rail, airports), power and water supply, storm water, and sewerage are highly sensitive to a range of extreme weather and climate events including temporary and permanent flooding arising from extreme precipitation, high winds, storm surges, and sea level rise (Horton et al. 2010; Handmer et al. 2012; Hanson and Nicholls 2012; Aerts et al. 2013). The tourism development experienced a comparable pattern, requiring host facilities and corresponding services. In Algeria, for example, construction projects have been carried out among the coastal paleo-dunes despite the existing Littoral Law 02-2002 (coastal protection) and the Law 01-3-2003 related to the Impact Expertise (Senouci and Taibi 2019). A similar situation exists in the industrial sector (e.g., desalinisation plant and electricity power station built on the beach).

Increases in temperature and decreases in precipitation could alter fire regimes affecting forest cover and could increase the intensity and frequency of drought resulting, in combination with other factors, in desertification (*Sections 2.4.1.2* and *2.6.3*). Future changes in climate could decrease food production (*Section 2.4.1.2*) and may alter the use of land all over the Mediterranean (*Section 2.4.1.3*). Future changes in the quantity and intensity of rain could affect the water cycles and increase the risk of floods (*Sections 2.2.6* and *2.6.3*).

Climate change and variability drives dynamics of marine species

Climate change and variability has led to concomitant changes in Mediterranean marine ecosystems and resources, with various implications on species diversity and composition, where species with limited locomotive capacity or confined in fragmented habitats seem more likely to be affected (Lejeusne

and Chevaldonné 2006; Ledoux et al. 2015). Examples of this changing environment, among others, are the mass mortality events of gorgonians and other sessile metazoans in northwestern Mediterranean (Garabou et al. 2009; Rivetti et al. 2014) and the continuous decline of *Posidonia* meadows (Marbà and Duarte 2010), the increase in the frequency of red tides and of gelatinous carnivore outbreaks (Conversi et al. 2010), the “tropicalization” of marine fauna in favour of the more thermophilic ones (Bianchi 2007), and the increase spread of microbial pathogens associated with water temperature rise (Danovaro et al. 2009) (*Section 4.1.1.1*).

At the end of 1980s and especially during the mid-1990s the Mediterranean Sea underwent regime shifts (Conversi et al. 2010; Alheit et al. 2019) that inflicted major atmospheric, hydrological and ecosystem changes, also affecting marine resources, mainly fisheries. There have been various studies linking ocean-atmospheric processes such as the Atlantic Multidecadal Oscillation (AMO), the North Atlantic Oscillation (NAO) and the Western Mediterranean Oscillation (WeMO) indices to alterations on the distribution and biomass of pelagic fish, as well as their catch composition (Alheit et al. 2019). Pelagic fish populations, more than other fish species, act as sentinels of these environmental changes. For example, during mid-1990s in the Mediterranean the highly correlated sea surface temperature and AMO index show a sharp increase (Marullo et al. 2011; Macías et al. 2013), whereas the dynamics of many fish species - mainly pelagic - show a conspicuous change around that time.

It is not yet clear how these changes impact pelagic fish population dynamics, combined with the pressures imposed by anthropogenic activity. Fifty-nine taxonomic groups (species or groups of species) showed an abrupt change in their landings in the mid-late 1990s (Tzanatos et al. 2014) with approximately 64% of these changes being correlated with sea surface temperature, mostly inversely correlated. The landings of some species (European sardine *Sardina pilchardus*, squids *Loligo* spp., Norway lobster *Nephrops norvegicus*, and hake *Merluccius merluccius*) decreased conspicuously in the mid-1990s, whereas those of other species (European anchovy *Engraulis encrasicolus* and greater amberjack *Seriola dumerili*) increased (Tzanatos et al. 2014). A study of the fisheries landings of 30 fish and invertebrate taxonomic groups revealed regime shifts at the mid-1990s, concurrent with the sea temperature increase in the eastern and western basins (Vasilakopoulos et al. 2017). The late 1990s

was determined as the turning point for the northward expansion of warm-water species in the Mediterranean (Azzurro et al. 2011). This was confirmed by Raitsos et al. (2010) who showed a clear increase of non-indigenous species entering into the eastern Mediterranean Sea in 1998. Earlier, Pinnegar et al. (2003) reported that the diversity of the western Mediterranean finfish landings increased dramatically after 1995 as a result of new species entering the catch.

Round sardinella (*Sardinella aurita*), a warm-water small pelagic fish species distributed along the southern Mediterranean coastline has been reported to have expanded its distribution to the northern Aegean (Tsikliras 2008), the northern Adriatic (Sinović et al. 2004), the Gulf of Lions (Francour et al. 1994), and the northwestern Mediterranean (Sabatés et al. 2006, 2009). A significant positive relationship between round sardinella landings and sea surface temperature anomalies has been reported for the western (Sabatés et al. 2006) and eastern Mediterranean (Tsikliras 2008). The northward distributional shift coincides with the beginning of positive temperature anomalies in the mid-1990s (Tsikliras 2008; Sabatés et al. 2009; Stergiou et al. 2016). Similarly, concomitant with the sea surface temperature change in the western Mediterranean, the landings of bluefish (*Pomatomus saltatrix*) quadrupled due to a northward expansion of the species (Sabatés et al. 2012) and, at the same time, anchovies returned to high biomass, as a result of increasing sea surface temperatures in the Adriatic Sea (Vilibić et al. 2016).

The effect of the AMO and NAO signals across the Mediterranean Sea sub-regions (western, central and eastern) on the small (European sardine *Sardina pilchardus*, European anchovy *Engraulis encrasicolus*, round sardinella *Sardinella aurita* and European sprat *Sprattus sprattus*) and medium (Atlantic mackerel *Scomber scombrus*, Atlantic chub mackerel *Scomber japonicus*, Atlantic horse mackerel *Trachurus trachurus*, Mediterranean horse mackerel *Trachurus mediterraneus*) pelagic fishes have been recently studied in the western, central and eastern Mediterranean Sea (Tsikliras et al. 2019). The pelagic fishes of the central and eastern Mediterranean respond most strongly to AMO variability and those of the central and western Mediterranean also respond to the NAO, while the effect of the NAO on pelagic fishes of the eastern Mediterranean was not significant (Tsikliras et al. 2019). Generally, various indicators revealed that the time of the pelagic fish response to the AMO and NAO signals varied among the

Mediterranean sub-regions (Alheit et al. 2014; Tsikliras et al. 2019).

Finally, the mean temperature of the catch, an indicator that assesses the effect of global warming on the exploited marine communities (Cheung et al. 2013b), has been increasing across the Mediterranean showing that the ratio of thermophilous (warm-water) to psychrophilous (cold-water) marine species has been changing in favour of the former. This is indicative of either an increase in the relative proportion of thermophilous species in the catches or a decrease in the relative proportion of the psychrophilous ones (Tsikliras and Stergiou 2014).

2.6.2.4 Effects of land use on climate change

Changes in crop use (Tribouillois et al. 2018), especially in forest cover, affect the balance between sink and release of CO₂ and the emissions of biogenic volatile organic compounds (BVOCs) in the atmosphere (Doblas-Miranda et al. 2017) (Sections 2.4.1.2 and 3.1.2.1).

Modification of surface albedo by land use changes also entail a highly potential impact on climate change (Benas and Chrysoulakis 2015). Changes in forest or dehesa/montado cover due to reforestation could reduce albedo (Rotenberg and Yakir 2011; Godinho et al. 2016), while fires increase radiations returns to the atmosphere (Sánchez et al. 2015), with contrasting effects on local climate. Agricultural cover may decrease or increase albedo (Giannakopoulou and Toumi 2012; Carrer et al. 2018), while urban sprawl definitely increases the radiation absorption and therefore local temperature (Salvati et al. 2019) (Section 3.1.3.1).

Change of land use and irrigation practices increase evapotranspiration and have a net cooling effect in some areas of the Mediterranean region (Zampieri and Lionello 2011; Thiery et al. 2017; Gormley-Gallagher et al. 2020).

2.6.2.5 Links between trends in non-indigenous species and climate change

Impact of climate change on marine non-indigenous species

The introduction of non-indigenous species and global warming interact in complex ways (Stachowicz et al. 2002), and are linked also in the Mediterranean Sea (Occhipinti-Ambrogi 2007).

This connection strongly depends on the species and the mode of its introduction, establishment and colonization. Overall, there is a strong trend of "tropicalization" of temperate areas through the movement of warm-loving (thermophilic) species toward the poles in areas of rapid ocean warming, and with increasingly strong impacts on local communities (Vergés et al. 2014a, 2016). These are not considered as introductions of non-indigenous species per se. But ocean warming may facilitate the establishment and spread of thermophilic non-indigenous species. The success of establishment of an introduced species depends on how suitable the ocean climatic parameters are in the region of introduction. Because successful non-indigenous species are typically generalists with broader climatic tolerances, they are usually considered able to cope better with climate change than native ones (Walther et al. 2009).

There is limited evidence for effects of climate change on the introduction of non-indigenous species. Theoretically, at the trailing "warm" edge of species distributions, the populations of sensitive cold-affinity species should reduce (and eventually extirpate) and that of warm affinity species (including thermophilic species) should increase (Bates et al. 2014). In the Mediterranean Sea, tropicalization evidently occurs (Vergés et al. 2014a), mainly in the Levant, by Lessepsian introductions, and ocean warming was suggested to facilitate the successful establishment of non-indigenous species (Raitsos et al. 2010). Ocean warming probably also helps to spread both native thermophilic species and successful Lessepsian species westward along the basin's temperature gradient, and also northward into the Aegean and Adriatic seas, or even the Ligurian Sea, but there very few direct empirical studies to demonstrate that. Recent analysis of fish trawl data from the southeast Mediterranean (Israel) does strongly suggest that non-indigenous species are indeed promoted by warming while natives are declining (Givan et al. 2017a).

Some studies suggest that habitats degraded by global warming are more likely affected by non-indigenous species than nearly-pristine habitats, envisaging explicitly or not, a cause and effect link between climate warming and the success of introductions (Stachowicz et al. 2002; Bianchi 2007; Galil 2007; Occhipinti-Ambrogi 2007). However, field observations do not support this idea, but reveal instead conflicting results that have provoked intense debate (Boudouresque and Verlaque 2010). For example, in the Mediterranean

Sea, well-structured and conserved habitats (such as coralligenous or *Cystoseira* forests) are able to mitigate and delay the proliferation and spread of the non-indigenous alga *Caulerpa cylindracea*, probably because the complexity of substrata (enhanced by gorgonians or canopy algae presence) is a key factor limiting its colonization and spread (Ceccherelli et al. 2002; Bulleri and Benedetti-Cecchi 2008; Verdura et al. 2019). In contrast, mass mortality of structural native species and subsequent increase of turf-forming species due to an extreme climatic event indirectly promoted the introduction of *C. cylindracea* in a coralligenous habitat (Verdura et al. 2019). However, in other non-indigenous algae such is *Lophocladia lallemandii*, introduction is favoured by more complex and rich communities (Cebríán et al. 2018), and thus simplification derived from climate change effects is expected not to enhance the capacity of *Lophocladia* to establish itself, but prevents its spread.

Using natural laboratories to test the thermal performance curves and sensitivity to acidification of key native and non-indigenous species, as well as the impact of climate change related environmental alteration on species interactions and communities and their ecosystem functions, are critical for better understanding and forecasting of the interactions between climate change and the introduction of non-indigenous species (Rilov et al. 2019a). For example, heat polluted areas and CO₂ vents (Hall-Spencer et al. 2008), as well as laboratory experiments in near-natural mesocosm systems (Wahl et al. 2015). Such recent measurements and experiments in the southeastern Levant have shown that some non-indigenous species (foraminifera) are tolerant to extreme thermal stress (Titelboim et al. 2017), that under warming and acidification conditions most Lessepsian species perform better than native species (Guy-Haim et al. 2016; Guy-Haim 2017), and demonstrated that different thermal performance of two Red Sea foraminifera explain why one species was introduced and the other did not (Titelboim et al. 2019). Furthermore, mesocosm work showed that a *Cystoseira* community becomes more heterotrophic and more dominated by non-indigenous species (but species richness does not change), demonstrating the profound impact of the combination of climate change and the introduction of non-indigenous species on ecosystem function (Guy-Haim et al. 2016; Rilov et al. 2019b).

Impacts of climate change on terrestrial non-indigenous species

There are five non-exclusive consequences of climate change on non-indigenous species: (1) altered transport and introduction mechanisms, (2) establishment of new species, (3) altered impact of existing non-indigenous species, (4) altered distribution of existing non-indigenous species, and (5) altered effectiveness of control strategies (Hellmann et al. 2008) (Section 2.5.1.3).

The influence of climate change on terrestrial non-indigenous species highly depends on species physiological strategy and reproductive adaptations (Bale and Hayward 2010; Antunes et al. 2018). Generalized ecosystem models of plant functional groups applied to Mediterranean islands indicate that climate change might promote the introduction of broadleaved trees (e.g., *Ailanthus altissima*) more than C₄ tropical grasses (e.g., *Amaranthus retroflexus*) (Gritti et al. 2006). Many non-indigenous species from temperate and cold climates might only be able to shift their ranges northward or to expand in altitude because they will be limited by drought and high temperatures (Storkey et al. 2014; Gallardo et al. 2017). While non-indigenous species whose native ranges are drier and warmer than their introduced ranges can be at an advantage to occupy niches at southern latitudes (Gallardo et al. 2017). Therefore, some species might loose and some gain suitable areas for introduction. Regions which will get drier are predicted to lose the highest number of potential non-indigenous species.

For introduced gardening plants, the climatically suitable areas with future climate change are unequally distributed across Europe with more suitable areas in the East than in the West of the Mediterranean Basin (Dullinger et al. 2017). This will be the case for *Cortaderia* which suitable area can increase 69-116% for 2060 (Tarabon et al. 2018) or for *Nassela* that can increase up to 47% for 2018 (Watt et al. 2011).

Similarly, weeds in crops can experience range shifts, niche shifts and trait shifts with climate change that will influence the agronomic practices to reduce their interference to crop production (Peters et al. 2014). Weeds in cereals crops will also advance towards northeastern Europe and remain or contract their distribution in warm areas of the Mediterranean region (Castellanos-Frías et al. 2014).

Climate change can advance the phenology of non-indigenous plant species including their fecundity

(Chuine et al. 2012), pollen production and seed maturation (Leiblein-Wild et al. 2016). Changes in pollen production can exacerbate the problem caused by allergenic non-indigenous plants such as the American *Ambrosia* because the allergenic risk is predicted to increase under all climate scenarios tested (Rasmussen et al. 2017).

Besides the influence of climate change on the establishment and spread of non-indigenous species, a remaining question is whether their impacts on native species increase in combination with climate change. A few greenhouse experiments have explored the interaction between competition of non-indigenous species and drought on the performance of native species (García-Serrano et al. 2007; Matesanz et al. 2008; Werner et al. 2010) and have found a non-synergistic effect. The interaction of climate change and introduction of non-indigenous species is a research area that requires further experimentation for productive systems such as the effect of weeds, pests and pathogens on crops and forestry (Ramesh et al. 2017).

2.6.2.6 Impacts of pollution on land and sea use

One of the major drivers relative to greenhouse gases pollution may be the CO₂ fertilization affecting forests. The balance between faster growth due to the fertilization effect and hydric stress due to most likely warmer and drier conditions have generated a considerable debate in the Mediterranean area (Keenan et al. 2011; Peñuelas et al. 2011). However, the most recent studies mainly corroborate that the effects of CO₂ fertilization will be negligible under the predicted climate conditions for the region (Camarero et al. 2015; Nunes et al. 2015; Gea-Izquierdo et al. 2017), despite some exceptions (Koutavas 2013; Barbeta and Peñuelas 2017). The potential effects of nitrogen deposition on Mediterranean forest growth also seem to be low (Ochoa-Hueso et al. 2014).

2.6.2.7 Impacts of land and sea use change on pollution

Intensive farming increases releases of nutrients and pesticides in aquifers while higher releases of methane in the air. The effects of the increase of livestock production on greenhouse gas emissions are assessed in Section 3.2.3.2.

Urban sprawl is associated to higher traffic related emissions (Sections 2.3.3 and 2.4.1.2).

2.6.2.8 Pollution effects on non-indigenous species

Pollution can make environmental conditions less tolerable for native species, and provide space and nutrients for opportunists, including non-indigenous species (Crooks et al. 2011).

2.6.2.9 Effects of land and water use on non-indigenous species

Habitat destruction causes disturbance, which opens space for non-indigenous species (Hobbs and Huenneke 1992).

2.6.2.10 Effects of non-indigenous species on land and sea use

Outbreaks of forest non-indigenous insects could alter forest cover (*Section 2.5.2.2*).

2.6.3 More complex interactions among drivers

2.6.3.1 Floods

Floods are an illustrative example of the combination of different drivers such as climate change (extreme precipitation events), land use change (catchment changes on river forests, forest cover, etc.) and even indirect drivers (among them and principally, urban sprawl in risk areas) (*Sections 3.1.3.3 and 3.1.4.1*).

2.6.3.2 Desertification

Puigdefábregas and Mendizabal (1998) analyzed FAO data from Morocco, Algeria and Tunisia during the period 1950-1993, associating desertification to socio-economic boundary conditions and over-exploitation by showing clear increases in population (pressure) and in the use of unsustainable land use practices in the Mediterranean, principally irrigation (*Section 6.6*).

Desertification is in fact the result of two different factors in origin operating in combination, prolonged drought of climatic origin and land exploitation of human origin (Le Houérou 1996). In Mediterranean arid lands, mainly during the 20th century, short-term planning of agricultural policies and overexploitation, mainly in the form of overgrazing but also fuelwood collection and ground water exploitation, contributed to soil quality decline and massive erosion. Deteriorating conditions have a great impact on the lives of inhabitants of Mediterranean drylands and force

most of them to migrate (Mohamed and Squires 2018) (*Sections 3.2.1.4, 4.3.1 and 6.6*).

2.6.3.3 Wildfires

One relevant consequence of the Mediterranean Climate characterized by dry summers are forest fires. Those can be exacerbated by drought conditions (Turco et al. 2018) but in turn they can affect drastically the flood generation both due to the erosion and the loss of forest mass. Although some forest fires can be provoked or as a result of recklessness, they mostly depend on the state of the vegetation and the climatic and meteorological situation. Consequently, fire regimes will be affected by climate change, if not already affected (Sarris et al. 2014). The Mediterranean is a high fire-risk region, where fires are the cause of severe agricultural, economic and environmental losses and even human casualties (Moreira et al. 2011; Keeley et al. 2012; San-Miguel-Ayanz et al. 2013; Bowman et al. 2017). For instance, the fire seasons in 2017 and 2018 was severe in many regions of Southern Europe, with large wildfires associated with unusually intense droughts and heat-waves (Sánchez-Benítez et al. 2018). In Portugal, the year of 2017 was particularly tragic. An extended and extraordinarily intense fire season yielded a record total burned area of about 500,000 hectares and more than 120 fatalities (Turco et al. 2019). Instead, the summer of 2018 will be remembered by the deadliest fires ever recorded affecting Greece, when a series of wildfires close to Athens killed 99 people, the deadliest in Greece history (AghaKouchak et al. 2018).

However, although several reports, ranging from popular media through to peer-reviewed scientific literature, have led to a shared perception that fires have increased or aggravated in recent years, the quantitative evidence available indicated that fires are decreasing on recent decades in this area (Turco et al. 2016). The increased efforts in fire suppression have probably played an important role in driving the general downward trends described for most of the Mediterranean area (Moreno et al. 2014; Ruffault et al. 2015). In recent decades fire management strategies have improved thanks to new technologies and experience while climate drivers have led to an opposite trend (Amatulli et al. 2013; Batllori et al. 2013; Bedía et al. 2013; Turco et al. 2014; Dupire et al. 2017; Fréjaville and Curt 2017) (*Section 4.3.2.1*).

FULL NAME	SHORT NAME	THEMATIC FOCUS	NO. OF SCENARIOS	TIME HORIZON	SPECIFIC REGIONAL FOCUS	REFERENCE(S)
GLOBAL SCALE						
Special Report on Emissions Scenarios	SRES	Emission of greenhouse gases	4	2100	-	Nakićenović, 2000
Shared Socioeconomic Pathways	SSPS	Multidisciplinary with a focus on challenges to climate change adaptation and mitigation	5	2100	-	(O'Neill et al. 2014, 2017)
MEDITERRANEAN SCALE						
A sustainable future for the Mediterranean	-	Multidisciplinary and cross-sectoral with an emphasis on sustainable development	2	2025	-	Benoit and Comeau, 2005
Mediterranean scenarios (MedAction project)	-	Multidisciplinary and cross-sectoral with an emphasis on desertification	3	2030	Northern Mediterranean case studies in ES, GR, IT, PT	Kok et al., 2006
EuroMed-2030	-	Multidisciplinary and cross-sectoral with a focus on the Euro-Mediterranean relationship	4	2030	-	EC/DG for Research and Innovation, 2011
Tomorrow, the Mediterranean	-	Multidisciplinary, cross-sectoral, with emphasis on economic development	3	2030	-	IPMED, 2011
Scenarios for the Mediterranean Region	-	Evolution of regional dynamics and role of the private sector in shaping business and political environments	3	2030	Exclusion of AL, BA, HR, IL, ME, PS, TR	World Economic Forum, 2011
Mediterranean Coastal SSPs	-	Regional and sectoral extension of the global SSPs for Mediterranean coastal regions	5	2100	-	Reimann et al., 2018
EUROPEAN SCALE						
Integrated Visions for a Sustainable Europe	VISIONS	Sustainable development	3	2020, 2050	-	Rotmans et al., 2000
Demographic and Migratory Flows Affecting European Regions and Cities	DEMIFER	Demography and European policies	5	2050	-	Rees et al., 2012
Climate Change Integrated Assessment Methodology for Cross-Sectoral Adaptation and Vulnerability in Europe	CLIMSAVE	Multidisciplinary and cross-sectoral, with emphasis on ecosystem services and provisions	4	2050	-	Gramberger et al., 2013
Territorial Scenarios and Visions for Europe	ET2050	Territorial development and cohesion	4	2050	-	MCRIT, 2015
Demographic Scenarios for the EU	-	Demographic development with a focus on aging, migration and education	3-4	2060	EU	Lutz et al., 2019
European Shared Socioeconomic Pathways	Eur-SSPs	Regional extension of the global SSPs for the European context	4	2040, 2070, 2100	Additional case study in Iberia	Kok et al., 2019

Table 2.4 | Overview of selected socioeconomic scenarios that cover Mediterranean countries, partly based on Rohat et al. (2018) and Sanna and Le Tellier (2013). ISO country codes: AL: Albania, BA: Bosnia and Herzegovina, ES: Spain, GR: Greece, HR: Croatia, IL: Israel, IT: Italy, ME: Montenegro, PT: Portugal, PS: State of Palestine, TR: Turkey.

2.7 Mediterranean socioeconomic scenarios

Environmental-change-related impacts will be driven not only by changes in climatic conditions, but also by changes in socioeconomic conditions. Prevailing socioeconomic conditions, in particular, determine a society's resilience to climatic hazards. When assessing future risks due to climate change, it is therefore crucial to account for plausible changes in socioeconomic conditions using a range of socioeconomic scenarios (González-Moreno et al. 2013).

A large number of socioeconomic scenarios have been developed in the past decades, focusing on a multitude of disciplines, sectors, and regions. Few of these scenarios were developed specifically for the Mediterranean region and even those usually only cover some of the Mediterranean countries, with a strong bias toward northern Mediterranean countries that are members of the European Union. *Table 2.4* provides an overview of a range of

socioeconomic scenarios developed in the last two decades that cover socioeconomic developments




Figure 2.30 | Mediterranean population projections under the different Shared Socioeconomic Pathways (SSPs), A) total Mediterranean population, B) population by geographical region (Kc and Lutz 2017).

in the Mediterranean either fully or partially in terms of geographic coverage.

The most recent socioeconomic scenarios that account for socioeconomic developments in the entire Mediterranean region (as defined in this report) are the state-of-the-art global-scale Shared Socioeconomic Pathways (SSPs). The SSPs explore broad-scale societal trends in the course of the 21st century both qualitatively, in the form of scenario storylines, and quantitatively, in the form of national-level projections of key variables such as population (Kc and Lutz 2017), urbanization (Jiang and O'Neill 2017), and Gross Domestic Product (GDP) (Cuaresma 2017; Dellink et al. 2017; Leimbach et al. 2017). The Mediterranean population is projected to range from 607 million (SSP1) to 659 million (SSP3) in 2050 and from 518 million to 770 million in 2100 (Fig. 2.30a), with considerable differences across regions (Fig. 2.30b). The largest share of the population is projected to live in Egypt under all SSPs, except SSP5 where France

is the most populous country in 2100 due to very high work migration into northern Mediterranean countries (O'Neill et al. 2017).




Figure 2.31 | Mediterranean Gross Domestic Product (GDP) projections under the different Shared Socioeconomic Pathways (SSPs), A) Mediterranean average GDP per capita, B) average GDP per capita by geographical region (Dellink et al. 2017).

The regional average GDP per capita is projected to grow from 16,000 US\$ yr⁻¹ in 2010 to between 24,000 US\$ yr⁻¹ (SSP3) and 45,000 US\$ yr⁻¹ (SSP5) in 2050 and to roughly 35,000 US\$ yr⁻¹ (SSP3) to 142,000 US\$ yr⁻¹ (SSP5) in 2100 (*Fig. 2.31a*). The differences in average GDP per capita are small between (potential) EU candidate countries, countries from the Middle East and the Maghreb region. EU member states have the highest average GDP per capita under all SSPs (*Fig. 2.31b*).

In order to increase the usefulness of SSPs for impact, adaptation, and vulnerability assessments (van Ruijven et al. 2014), spatially explicit population projections that account for spatial changes in population distribution in the course of the 21st century have been produced, using the national totals as input data. These are available for all Mediterranean countries at a horizontal resolution of 7.5 arc minutes (Jones and O'Neill 2016) and 30 arc seconds (Merkens et al. 2016; Gao 2017). Further, downscaled GDP projections are available for SSPs 1-3 at a resolution of 30 arc minutes (Murakami and Yamagata 2019). All of these projections are based on the underlying global SSP assumptions.

As the global assumptions do not necessarily reflect the socioeconomic developments at the regional scale, extensions of the global-SSPs for the Mediterranean coastal zone have been developed (Reimann et al. 2018). These Mediterranean coastal SSPs account for region-specific developments as well as for changing attractiveness of coastal regions for human settlement across the SSPs, while at the same time ensuring consistency with the global SSPs (Zurek and Henrichs 2007). The Mediterranean coastal SSPs consist of qualitative narratives for each coastal SSP – SSP1 "Green Coast", SSP2 "No Wind of Change", SSP3 "Troubled Waters", SSP4 "Fragmented Coast", and SSP5 'Coast Rush' – differentiating between regional socioeconomic developments in northern versus southern and eastern parts of the region; and of spatially explicit population projections for all Mediterranean riparian countries at a resolution of 30 arc seconds (*Fig. 2.32*).




Figure 2.32 | Spatially explicit population projections produced for the Mediterranean Coastal SSPs. Selected population grids for the base year 2010 and each SSP in 2100 (Reimann et al. 2018), GPW = Gridded Population of the World (Center for International Earth Science Information Network - CIESIN - Columbia University 2016).

BOX 2.1

How much has the Mediterranean Basin warmed since the pre-industrial period?

The UNFCCC Paris Agreement of 2015 strengthens the initial goal of the Article 2 in the United Nations Framework Convention on Climate Change (UNFCCC), "to achieve ... stabilization of greenhouse gas concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system", by "holding the increase in the global average temperature to well below 2°C above pre-industrial levels and pursuing efforts to limit the temperature increase to 1.5°C above pre-industrial levels". While these temperature goals refer to the global average, it is a natural question to ask, for any region, how much warming has been observed "above pre-industrial levels".

This is only apparently a simple question. To provide such an estimate it is necessary to clarify the meaning of pre-industrial, the information available on its average temperature in the region, a definition of present average temperature and the method used for estimating it. These issues were at the core of the IPCC special report on global warming of 1.5°C (SR15, IPCC 2018) and specifically considered in its *Chapter 1* (Allen et al. 2018).

The period 1850-1900 has been identified (Allen et al. 2018) as a suitable approximation for the estimate of the pre-industrial average temperature, because it combines typical pre-industrial solar and volcanic forcing, low anthropogenic greenhouse gas concentrations, and sufficient coverage of instrumental temperature observations. The choice of the period 1850-1900 is not completely free of problems, because it is indeed already affected by increasing greenhouse gas concentrations, with a partial compensation caused by aerosols. Further, strong volcanic eruptions occurred in the period 1880-1900. In this box, we follow SR15 but also the indications of the World Meteorological Organization (WMO 2017) and compute average pre-industrial temperature for a 30-years period, the central 1861-1890 period of the 1850-1900 "pre-industrial" period.

The number and distributions of instrumental observation have changed significantly over time, and station density could become critically low in African and Asian areas of the Mediterranean Basin. Further, the collection of observations over sea is systematically more problematic than over land, notably prior to the 20th century. Rather than analyzing station data, we seek consistency with global estimates and we use the HadCRUT4 dataset (Morice et al. 2012) and the CRUTEM4 (Jones et al. 2012) data sets for the land+sea and land only analysis, respectively. These are two widely used gridded global data sets with a resolution of 5 degrees longitude/latitude since 1850 until present. Other data sets at higher spatial resolution are available (such as the recently updated version of CRU TS) (Harris et al. 2020), but they do not reach back far enough into the 19th century for the estimation of pre-industrial conditions. For this analysis, the Mediterranean Basin is defined as the

domain from 10°W to 40°E of longitude and from 30°N to 47.5°N of latitude. For the averaging, an interpolation (based on the closest neighbours) to 1° spatial resolution was undertaken.

Obviously observed temperature values are not yet available for the period 2020-2034. This prevents computing the level of warming in 2020 using a simple 30-year average. Here, we make the conservative assumption that warming in the future will continue at the same rate of the last 50 years and compute the 2020 temperature by extrapolating the linear trend of the 1970-2019 period Fig. 2.33). In 2020 the Mediterranean Basin is 1.5°C warmer than in the preindustrial with a likely uncertainty range of +0.11°C. Land areas have warmed more than the sea. If only land areas are considered, the 2020 temperature is 1.8°C warmer than pre-industrial with a likely uncertainty range of +0.12°C.




Figure 2.33 | Time series (grey bars) of the annual average temperature of the Mediterranean Basin (30°N to 47.5°N and 10°W to 40°E) considering the whole region (panel A) or only its land areas (panel B). The analysis is based on the HadCRUT4 data set for the land+sea analysis (Morice et al. 2012) and on the CRUTEM4 data set for the land analysis (Jones et al. 2012)[Jones et al, 2012]. To avoid edge effects, before calculating the averages, the data have been interpolated to 1° spatial resolution. The values are expressed as anomalies from the pre-industrial period (1860-1890). The blue horizontal lines represent the 30-yr averages by steps of 20 years. The red curve is the linear trend linear trend calculated for the 1970-2019 period, extrapolated up to 2020. The black line is the smoother time-series. The blue square represents the likely interval (probability >0.66) for the present warming.

BOX 2.2


Representative Concentration Pathways (RCPs)

Representative Concentration Pathways (RCPs) are greenhouse gas concentration pathways, developed by the IPCC in order to explore the physical outcomes of different climate policies, notably regarding the mitigation of greenhouse gas emissions.

RCP2.6: The RCP2.6 was developed by the IMAGE modeling team of the Netherlands Environmental Assessment Agency (van Vuuren et al. 2011). The emission pathway is representative for scenarios in the literature leading to very low greenhouse gas concentration levels. It is a so-called “peak” scenario: its radiative forcing level first reaches a value around 3.1 W m^{-2} mid-century, returning to 2.6 W m^{-2} by 2100. In order to reach such radiative forcing levels, greenhouse gas emissions (and indirectly emissions of air pollutants) are reduced substantially over time.

RCP4.5: It was developed by the MiniCAM modeling team at the Pacific Northwest National Laboratory’s Joint Global Change Research Institute (Clarke et al. 2014). It is a stabilization scenario where total radiative forcing is stabilized before 2100 by employment of a range of technologies and strategies for reducing greenhouse gas emissions. It is often considered as an intermediate scenario.

RCP8.5: The RCP8.5 was developed by the MESSAGE modeling team and the IIASA Integrated Assessment Framework from the International Institute for Applied Systems Analysis (IIASA), Austria (Riahi et al. 2011). The RCP8.5 is characterized by increasing greenhouse gas emissions over time representative for scenarios in the literature leading to high greenhouse gas concentration levels, reaching $+8.5 \text{ W m}^{-2}$ additional surface radiative forcing in 2100. It is often considered as a “business-as-usual” scenario.



References

- Abadie J, Dupouey J-L, Avon C, Rochel X, Tatoni T et al. 2018 Forest recovery since 1860 in a Mediterranean region: drivers and implications for land use and land cover spatial distribution. *Landscape Ecol.* 33, 289–305. doi: [10.1007/s10980-017-0601-0](https://doi.org/10.1007/s10980-017-0601-0)
- Abbes K, Harbi A, Chermiti B 2012 The tomato leafminer *Tuta absoluta* (Meyrick) in Tunisia: current status and management strategies. *EPPO Bull.* 42, 226–233. doi: [10.1111/epp.2559](https://doi.org/10.1111/epp.2559)
- Abdallah Z, Mezghani-Kmekhem M, Bouktila D, Makni H, Makni M 2012 Genetic diversity of an invasive pest (*Oryctes agamemnon* Burmeister, Coleoptera: Scarabaeidae) of date palm in Tunisia, inferred from random amplified polymorphic DNA (RAPD) markers. *African J. Agric. Res.* 7, 1170–1176. doi: [10.5897/AJAR11.1580](https://doi.org/10.5897/AJAR11.1580)
- Abdennadher M, Hamza A, Fekih W, Hannachi I, Zouari Bellaaj A et al. 2012 Factors determining the dynamics of toxic blooms of *Alexandrium minutum* during a 10-year study along the shallow southwestern Mediterranean coasts. *Estuar. Coast. Shelf Sci.* 106, 102–111. doi: [10.1016/J.ECSS.2012.04.029](https://doi.org/10.1016/J.ECSS.2012.04.029)
- Abderrahim H, Chellali MR, Hamou A 2016 Forecasting PM10 in Algiers: efficacy of multilayer perceptron networks. *Environ. Sci. Pollut. Res.* 23, 1634–1641. doi: [10.1007/s11356-015-5406-6](https://doi.org/10.1007/s11356-015-5406-6)
- Abella A, Ria M, Mancusi C 2010 Assessment of the status of the coastal groundfish assemblage exploited by the Viareggio fleet [Southern Ligurian Sea]. *Sci. Mar.* 74, 793–805. doi: [10.3989/scimar.2010.74n4793](https://doi.org/10.3989/scimar.2010.74n4793)
- Abellán P, Carrete M, Anadón JD, Cardador L, Tella JL 2016 Non-random patterns and temporal trends (1912–2012) in the transport, introduction and establishment of exotic birds in Spain and Portugal. *Divers. Distrib.* 22, 263–273. doi: [10.1111/ddi.12403](https://doi.org/10.1111/ddi.12403)
- Abellán P, Tella JL, Carrete M, Cardador L, Anadón JD 2017 Climate matching drives spread rate but not establishment success in recent unintentional bird introductions. *Proc. Natl. Acad. Sci. U. S. A.* 114, 9385–9390. doi: [10.1073/pnas.1704815114](https://doi.org/10.1073/pnas.1704815114)
- Acácio V, Dias FS, Catry FX, Rocha M, Moreira F 2017 Landscape dynamics in Mediterranean oak forests under global change: understanding the role of anthropogenic and environmental drivers across forest types. *Glob. Chang. Biol.* 23, 1199–1217. doi: [10.1111/gcb.13487](https://doi.org/10.1111/gcb.13487)
- Acri F, Aubry FB, Berton A, Bianchi F, Boldrin A et al. 2004 Plankton communities and nutrients in the Venice Lagoon. Comparison between current and old data. *J. Mar. Syst.* 51, 321–329. doi: [10.1016/j.jmarsys.2004.05.019](https://doi.org/10.1016/j.jmarsys.2004.05.019)
- Adhikari PL, Maiti K, Overton EB 2015 Vertical fluxes of polycyclic aromatic hydrocarbons in the northern Gulf of Mexico. *Mar. Chem.* 168, 60–68. doi: [10.1016/j.marchem.2014.11.001](https://doi.org/10.1016/j.marchem.2014.11.001)
- Adloff F, Jordà G, Somot S, Sevault F, Arsouze T et al. 2018 Improving sea level simulation in Mediterranean regional climate models. *Clim. Dyn.* 51. doi: [10.1007/s00382-017-3842-3](https://doi.org/10.1007/s00382-017-3842-3)
- Adloff F, Somot S, Sevault F, Jordà G, Aznar R et al. 2015 Mediterranean Sea response to climate change in an ensemble of twenty first century scenarios. *Clim. Dyn.* 45, 2775–2802. doi: [10.1007/s00382-015-2507-3](https://doi.org/10.1007/s00382-015-2507-3)
- Adua M 1999 The sweet chestnut throughout history from the Miocene to the Third Millennium. *Acta Hortic.*, 29–36. doi: [10.17660/ActaHortic.1999.494.2](https://doi.org/10.17660/ActaHortic.1999.494.2)
- Aerts JCJH, Lin N, Botzen W, Emanuel K, de Moel H 2013 Low-Probability Flood Risk Modeling for New York City. *Risk Anal.* 33, 772–788. doi: [10.1111/risa.12008](https://doi.org/10.1111/risa.12008)
- AghaKouchak A, Huning LS, Chiang F, Sadegh M, Vahedifard F et al. 2018 How do natural hazards cascade to cause disasters? *Nature* 561, 458–460. doi: [10.1038/d41586-018-06783-6](https://doi.org/10.1038/d41586-018-06783-6)
- Aissaoui A, Armi Z, Akrout F, Ben Hassine OK 2014 Environmental factors and seasonal dynamics of *Prorocentrum lima* population in coastal waters of the Gulf of Tunis, South Mediterranean. *Water Environ. Res.* 86, 2256–2270. doi: [10.2175/106143014x139750355266](https://doi.org/10.2175/106143014x139750355266)
- Alessandri A, de Felice M, Zeng N, Mariotti A, Pan Y et al. 2015 Robust assessment of the expansion and retreat of Mediterranean climate in the 21st century. *Sci. Rep.* 4, 7211. doi: [10.1038/srep07211](https://doi.org/10.1038/srep07211)
- Alexander MA, Scott JD, Friedland KD, Mills KE, Nye JA et al. 2018 Projected sea surface temperatures over the 21st century: Changes in the mean, variability and extremes for large marine ecosystem regions of Northern Oceans. *Elementa* 6, 9. doi: [10.1525/elementa.191](https://doi.org/10.1525/elementa.191)
- Alexandri G, Georgoulas AK, Meleti C, Balis D, Kourtidis KA et al. 2017 A high resolution satellite view of surface solar radiation over the climatically sensitive region of Eastern Mediterranean. *Atmos. Res.* 188, 107–121. doi: [10.1016/J.ATMOSRES.2016.12.015](https://doi.org/10.1016/J.ATMOSRES.2016.12.015)
- Alfaro-Reyna T, Retana J, Martínez-Vilalta J 2018 Is there a substitution of Pinaceae by Fagaceae in temperate forests at the global scale? *Glob. Planet. Change* 166, 41–47. doi: [10.1016/j.gloplacha.2018.04.001](https://doi.org/10.1016/j.gloplacha.2018.04.001)
- Alheit J, Gröger J, Licandro P, McQuinn IH, Pohlmann T et al. 2019 What happened in the mid-1990s? The coupled ocean-atmosphere processes behind climate-induced ecosystem changes in the Northeast Atlantic and the Mediterranean. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 159, 130–142. doi: [10.1016/j.dsr2.2018.11.011](https://doi.org/10.1016/j.dsr2.2018.11.011)
- Alheit J, Licandro P, Coombs S, García A, Giráldez A et al. 2014 Atlantic Multidecadal Oscillation (AMO) modulates dynamics of small pelagic fishes and ecosystem regime shifts in the eastern North and Central Atlantic. *J. Mar. Syst.* 131, 21–35. doi: [10.1016/j.jmarsys.2013.11.002](https://doi.org/10.1016/j.jmarsys.2013.11.002)
- Allen MR, Dube OP, Solecki W, Aragón-Durand F, Cramer W et al. 2018 Framing and Context, in *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*, eds. Masson-Delmotte V, Zhai P, Pörtner H-O, Roberts D, Skea J et al. (Cambridge, United Kingdom and New York, USA: Cambridge University Press), 49–91.

- Alpert P, Baldi M, Ilani R, Krichak S, Price C et al. 2006 Chapter 2 Relations between climate variability in the Mediterranean region and the tropics: ENSO, South Asian and African monsoons, hurricanes and Saharan dust, in *Mediterranean*, eds. Lionello P, Malanotte-Rizzoli P, Boscolo RBT-D in E and ES (Elsevier), 149–177.
doi: [10.1016/S1571-9197\(06\)80005-4](https://doi.org/10.1016/S1571-9197(06)80005-4)
- Alpert P, Neeman BU, Shay-El Y 1990a Climatological analysis of Mediterranean cyclones using ECMWF data. *Tellus A Dyn. Meteorol. Oceanogr.* 42, 65–77.
doi: [10.3402/tellusa.v42i1.11860](https://doi.org/10.3402/tellusa.v42i1.11860)
- Alpert P, Neeman BU, Shay-El Y 1990b Intermonthly Variability of Cyclone Tracks in the Mediterranean. *J. Clim.* 3, 1474–1478.
doi: [10.1175/1520-0442\[1990\]003<1474:IVOCII>2.0.CO;2](https://doi.org/10.1175/1520-0442[1990]003<1474:IVOCII>2.0.CO;2)
- Alphan H 2018 Analysis of road development and associated agricultural land use change. *Environ. Monit. Assess.* 190, 5.
doi: [10.1007/s10661-017-6379-3](https://doi.org/10.1007/s10661-017-6379-3)
- Amatulli G, Camia A, San-Miguel-Ayanz J 2013 Estimating future burned areas under changing climate in the EU-Mediterranean countries. *Sci. Total Environ.* 450–451, 209–222.
doi: [10.1016/j.scitotenv.2013.02.014](https://doi.org/10.1016/j.scitotenv.2013.02.014)
- Améztegui A, Brotons L, Coll L 2010 Land-use changes as major drivers of mountain pine (*Pinus uncinata* Ram.) expansion in the Pyrenees. *Glob. Ecol. Biogeogr.* 19, 632–641.
doi: [10.1111/j.1466-8238.2010.00550.x](https://doi.org/10.1111/j.1466-8238.2010.00550.x)
- Amici V, Maccherini S, Santi E, Torri D, Vergari F et al. 2017 Long-term patterns of change in a vanishing cultural landscape: A GIS-based assessment. *Ecol. Inform.* 37, 38–51.
doi: [10.1016/J.ECOINF.2016.11.008](https://doi.org/10.1016/J.ECOINF.2016.11.008)
- Anav A, Mariotti A 2011 Sensitivity of natural vegetation to climate change in the Euro-Mediterranean area. *Clim. Res.* 46, 277–292. doi: [10.3354/cr00993](https://doi.org/10.3354/cr00993)
- Anderson DM, Cembella AD, Hallegraeff GM 2012 Progress in understanding Harmful Algal Blooms: Paradigm shifts and new technologies for research, monitoring, and management. *Ann. Rev. Mar. Sci.* 4, 143–176.
doi: [10.1146/annurev-marine-120308-081121](https://doi.org/10.1146/annurev-marine-120308-081121)
- Antunes C, Pereira AJ, Fernandes P, Ramos M, Ascensão L et al. 2018 Understanding plant drought resistance in a Mediterranean coastal sand dune ecosystem: differences between native and exotic invasive species. *J. Plant Ecol.* 11, 26–38.
doi: [10.1093/jpe/rtx014](https://doi.org/10.1093/jpe/rtx014)
- Argenti G, Bottai L, Chiesi M, Maselli F, Staglianò N et al. 2011 Analisi e valutazione di pascoli montani attraverso l'integrazione di dati multispettrali e ausiliari. *Riv. Ital. di Telerilevamento* 43, 45–57. https://flore.unifi.it/handle/2158/406248#.XMN_mJS_pPOQ [Accessed April 26, 2019].
- Arhami M, Hosseini V, Zare Shahne M, Bigdeli M, Lai A et al. 2017 Seasonal trends, chemical speciation and source apportionment of fine PM in Tehran. *Atmos. Environ.* 153, 70–82. doi: [10.1016/j.atmosenv.2016.12.046](https://doi.org/10.1016/j.atmosenv.2016.12.046)
- Arianoutsou M, Delipetrou P, Celesti-Grapow L, Basnou C, Bazos I et al. 2010 Comparing naturalized alien plants and recipient habitats across an east-west gradient in the Mediterranean Basin. *J. Biogeogr.* 37, 1811–1823.
doi: [10.1111/j.1365-2699.2010.02324.x](https://doi.org/10.1111/j.1365-2699.2010.02324.x)
- Arianoutsou M, Delipetrou P, Vilà M, Dimitrakopoulos PG, Celesti-Grapow L et al. 2013 Comparative patterns of plant invasions in the Mediterranean biome. *PLoS One* 8, e79174.
doi: [10.1371/journal.pone.0079174](https://doi.org/10.1371/journal.pone.0079174)
- Armi Z, Milandri A, Turki S, Hajjem B 2011 *Alexandrium catenella* and *Alexandrium tamarense* in the North Lake of Tunis: Bloom characteristics and the occurrence of paralytic shellfish toxin. *African J. Aquat. Sci.* 36, 47–56.
doi: [10.2989/16085914.2011.559688](https://doi.org/10.2989/16085914.2011.559688)
- Arndt E, Givan O, Edelist D, Sonin O, Belmaker J et al. 2018 Shifts in Eastern Mediterranean fish communities: Abundance changes, trait overlap, and possible competition between native and non-native species. *Fishes* 3, 19.
doi: [10.3390/fishes3020019](https://doi.org/10.3390/fishes3020019)
- Artale V, Calmant S, Malanotte-Rizzoli P, Pisacane G, Rupolo V et al. 2006 The Atlantic and Mediterranean Sea as connected systems, in *Developments in Earth and Environmental Sciences* (Elsevier), 282–323.
doi: [10.1016/S1571-9197\(06\)80008-X](https://doi.org/10.1016/S1571-9197(06)80008-X)
- Asaf D, Perdersen D, Peleg M, Matveev V, Luria M 2008 Evaluation of background levels of air pollutants over Israel. *Atmos. Environ.* 42, 8453–8463.
doi: [10.1016/j.atmosenv.2008.08.011](https://doi.org/10.1016/j.atmosenv.2008.08.011)
- Asia L, Mazouz S, Giuliano M, Dournenq P, Mille G et al. 2009 Occurrence and distribution of hydrocarbons in surface sediments from Marseille Bay (France). *Mar. Pollut. Bull.* 58, 443–451. doi: [10.1016/j.marpolbul.2008.11.022](https://doi.org/10.1016/j.marpolbul.2008.11.022)
- Astraldi M, Conversano F, Civitarese G, Gasparini GP, D'Alcala MR et al. 2002 Water mass properties and chemical signatures in the central Mediterranean region. *J. Mar. Syst.* 33, 155–177. doi: [10.1016/s0924-7963\(02\)00057-x](https://doi.org/10.1016/s0924-7963(02)00057-x)
- Athanasiadis PJ, Wallace JM, Wetstein JJ 2010 Patterns of wintertime jet stream variability and their relation to the storm tracks. *J. Atmos. Sci.* 67, 1361–1381.
doi: [10.1175/2009JAS3270.1](https://doi.org/10.1175/2009JAS3270.1)
- Auger-Rosenberg M-A, Boivin T 2016 Invasive fruit, cone and seed insects in the Mediterranean Basin, in *Insects and Diseases of Mediterranean Forest Systems* (Cham: Springer International Publishing), 239–259.
doi: [10.1007/978-3-319-24744-1_9](https://doi.org/10.1007/978-3-319-24744-1_9)
- Avtzis DN, Coyle DR, Christopoulos V, Roques A 2017 Biological invasions, national borders, and the current state of non-native insect species in Greece and the neighbouring Balkan countries. *Bull. Insectology* 70, 161–169.
<http://easin.jrc.ec.europa.eu/> [Accessed April 26, 2019]
- Azorin-Molina C, Vicente-Serrano SM, McVicar TR, Jerez S, Sanchez-Lorenzo A et al. 2014 Homogenization and assessment of observed near-surface wind speed trends over Spain and Portugal, 1961–2011. *J. Clim.* 27, 3692–3712.
doi: [10.1175/JCLI-D-13-00652.1](https://doi.org/10.1175/JCLI-D-13-00652.1)
- Azzurro E, Bariche M 2017 Local knowledge and awareness on the incipient lionfish invasion in the eastern Mediterranean Sea. *Mar. Freshw. Res.* 68, 1950. doi: [10.1071/MF16358](https://doi.org/10.1071/MF16358)
- Azzurro E, Moschella P, Maynou F 2011 Tracking signals of change in Mediterranean fish diversity based on local ecological knowledge. *PLoS One* 6, e24885.
doi: [10.1371/journal.pone.0024885](https://doi.org/10.1371/journal.pone.0024885)

- Azzurro E, Stancanelli B, Martino V Di, Bariche M 2017 Range expansion of the common lionfish *Pterois miles* (Bennett, 1828) in the Mediterranean Sea: an unwanted new guest for Italian waters. 6, 95–98. doi: [10.3391/bir.2017.6.2.01](https://doi.org/10.3391/bir.2017.6.2.01)
- Bacer S, Christoudias T, Pozzer A 2016 Projection of North Atlantic Oscillation and its effect on tracer transport. *Atmos. Chem. Phys.* 16, 15581–15592. doi: [10.5194/acp-16-15581-2016](https://doi.org/10.5194/acp-16-15581-2016)
- Baeza A, Rodríguez-Perulero A, Guillén J 2016 Anthropogenic and naturally occurring radionuclide content in near surface air in Cáceres (Spain). *J. Environ. Radioact.* 165, 24–31. doi: [10.1016/j.jenvrad.2016.08.018](https://doi.org/10.1016/j.jenvrad.2016.08.018)
- Balaguer R, Dimastrogiovanni G, García K, González E, Lysimachou A et al. 2018 *Ríos hormonados, Amplia presencia de plaguicidas endocrinos en los ríos españoles*. Ecologista. <https://www.ecologistasenaccion.org/wp-content/uploads/adjuntos-spip/pdf/informe-rios-hormonados.pdf>
- Baldacconi R, Corriero G 2009 Effects of the spread of the alga *Caulerpa racemosa* var. *cylindracea* on the sponge assemblage from coralligenous concretions of the Apulian coast (Ionian Sea, Italy). *Mar. Ecol.* 30, 337–345. doi: [10.1111/j.1439-0485.2009.00282.x](https://doi.org/10.1111/j.1439-0485.2009.00282.x)
- Bale JS, Hayward SAL 2010 Insect overwintering in a changing climate. *J. Exp. Biol.* 213, 980–994. doi: [10.1242/jeb.037911](https://doi.org/10.1242/jeb.037911)
- Balog I, Ruti PM, Tobin I, Armenio V, Vautard R 2016 A numerical approach for planning offshore wind farms from regional to local scales over the Mediterranean. *Renew. Energy* 85, 395–405. doi: [10.1016/j.renene.2015.06.038](https://doi.org/10.1016/j.renene.2015.06.038)
- Ban N, Schmidli J, Schär C 2015 Heavy precipitation in a changing climate: Does short-term summer precipitation increase faster? *Geophys. Res. Lett.* 42, 1165–1172. doi: [@10.1002/\(ISSN\)1944-8007.2015 EDHIGHLIGHTS](https://doi.org/10.1002/2014GL062588)
- Barbeta A, Peñuelas J 2017 Increasing carbon discrimination rates and depth of water uptake favor the growth of Mediterranean evergreen trees in the ecotone with temperate deciduous forests. *Glob. Chang. Biol.* 23, 5054–5068. doi: [10.1111/gcb.13770](https://doi.org/10.1111/gcb.13770)
- Barcikowska MJ, Kapnick SB, Krishnamurti L, Russo S, Cherchi A et al. 2020 Changes in the future summer Mediterranean climate: contribution of teleconnections and local factors. *Earth Syst. Dyn.* 11, 161–181. doi: [10.5194/esd-11-161-2020](https://doi.org/10.5194/esd-11-161-2020)
- Barhoumi B, Castro-Jiménez J, Guigou C, Goutx M, Sempéré R et al. 2018 Levels and risk assessment of hydrocarbons and organochlorines in aerosols from a North African coastal city (Bizerte, Tunisia). *Environ. Pollut.* 240, 422–431. doi: [10.1016/j.envpol.2018.04.109](https://doi.org/10.1016/j.envpol.2018.04.109)
- Bariche M, Torres M, Azzurro E 2013 The presence of the invasive lionfish *Pterois miles* in the Mediterranean Sea. *Mediterr. Mar. Sci.* 14, 292–294. doi: [10.12681/mms.428](https://doi.org/10.12681/mms.428)
- Barkaoui A 2003 *La marine carthaginoise : approche des activités militaires des carthaginois sur mer depuis les origines jusqu'en 146 av. J.-C. L'Or de temps*
- Barnes DKA, Galgani F, Thompson RCC, Barlaz M 2009 Accumulation and fragmentation of plastic debris in global environments. *Philos. Trans. R. Soc. B Biol. Sci.* 364, 1985–1998. doi: [10.1098/rstb.2008.0205](https://doi.org/10.1098/rstb.2008.0205)
- Barnes EA, Polvani L, Barnes EA, Polvani L 2013 Response of the midlatitude jets, and of their variability, to increased greenhouse gases in the CMIP5 models. *J. Clim.* 26, 7117–7135. doi: [10.1175/JCLI-D-12-00536.1](https://doi.org/10.1175/JCLI-D-12-00536.1)
- Barnston AG, Livezey RE 1987 Classification, seasonality and persistence of low-frequency atmospheric circulation patterns. *Mon. Weather Rev.* 115, 1083–1126. doi: [10.1175/1520-0493\(1987\)115<1083:CSAPOL>2.0.CO;2](https://doi.org/10.1175/1520-0493(1987)115<1083:CSAPOL>2.0.CO;2)
- Barreiro-Lostres F, Moreno A, Giralt S, Nadal-Romero E, Valeiro-Garcés B 2017 Erosion in Mediterranean mountain landscapes during the last millennium: a quantitative approach based on lake sediment sequences (Iberian Range, Spain). *Catena* 149, 782–798. doi: [10.1016/j.catena.2016.05.024](https://doi.org/10.1016/j.catena.2016.05.024)
- Barrios-Estrada C, de Jesús Rostro-Alanis M, Muñoz-Gutiérrez BD, Iqbal HMN, Kannan S et al. 2018 Emergent contaminants: Endocrine disruptors and their laccase-assisted degradation – A review. *Sci. Total Environ.* 612, 1516–1531. doi: [10.1016/j.scitotenv.2017.09.013](https://doi.org/10.1016/j.scitotenv.2017.09.013)
- Bartók B, Wild M, Folini D, Lüthi D, Kotlarski S et al. 2017 Projected changes in surface solar radiation in CMIP5 global climate models and in EURO-CORDEX regional climate models for Europe. *Clim. Dyn.* 49, 2665–2683. doi: [10.1007/s00382-016-3471-2](https://doi.org/10.1007/s00382-016-3471-2)
- Bartolini G, Morabito M, Crisci A, Grifoni D, Torrigiani T et al. 2008 Recent trends in Tuscany (Italy) summer temperature and indices of extremes. *Int. J. Climatol.* 28, 1751–1760. doi: [10.1002/joc.1673](https://doi.org/10.1002/joc.1673)
- Basnou C, Álvarez E, Bagaria G, Guardiola M, Isern R et al. 2013 Spatial patterns of land use changes across a Mediterranean metropolitan landscape: implications for biodiversity management. *Environ. Manage.* 52, 971–980. doi: [10.1007/s00267-013-0150-5](https://doi.org/10.1007/s00267-013-0150-5)
- Basnou C, Iguzquiza J, Pino J 2015 Examining the role of landscape structure and dynamics in alien plant invasion from urban Mediterranean coastal habitats. *Landsc. Urban Plan.* 136, 156–164. doi: [10.1016/j.landurbplan.2014.12.001](https://doi.org/10.1016/j.landurbplan.2014.12.001)
- Bates AE, Pecl GT, Frusher S, Hobday AJ, Wernberg T et al. 2014 Defining and observing stages of climate-mediated range shifts in marine systems. *Glob. Environ. Chang.* 26, 27–38. doi: [10.1016/j.gloenvcha.2014.03.009](https://doi.org/10.1016/j.gloenvcha.2014.03.009)
- Batllori E, Parisien M-A, Krawchuk MA, Moritz MA 2013 Climate change-induced shifts in fire for Mediterranean ecosystems. *Glob. Ecol. Biogeogr.* 22, 1118–1129. doi: [10.1111/geb.12065](https://doi.org/10.1111/geb.12065)
- Beaufoy G 2001 EU policies for olive farming-unsustainable on all counts.
- Beaufoy G, Pienkowski M 2000 The environmental impact of olive oil production in the European Union: practical options for improving the environmental impact. <http://ec.europa.eu/environment/agriculture/pdf/oliveoil.pdf> [Accessed April 26, 2019].
- Bedía J, Herrera S, Martín DS, Koutsias N, Gutiérrez JM 2013 Robust projections of Fire Weather Index in the Mediterranean using statistical downscaling. *Clim. Change* 120, 229–247. doi: [10.1007/s10584-013-0787-3](https://doi.org/10.1007/s10584-013-0787-3)
- Belhout D, Kerbachi R, Relvas H, Miranda AI 2018 Air quality assessment in Algiers city. *Air Qual. Atmos. Heal.* 11, 897–906. doi: [10.1007/s11869-018-0589-x](https://doi.org/10.1007/s11869-018-0589-x)

- Belmonte J, Vilà M 2004 Atmospheric invasion of non-native pollen in the Mediterranean region. *Am. J. Bot.* 91, 1243–1250. doi: [10.3732/ajb.91.8.1243](https://doi.org/10.3732/ajb.91.8.1243)
- Belušić Vozila A, Güttsler I, Ahrens B, Obermann-Hellhund A, Telišman Prtenjak M 2019 Wind Over the Adriatic Region in CORDEX Climate Change Scenarios. *J. Geophys. Res. Atmos.* 124, 110–130. doi: [10.1029/2018JD028552](https://doi.org/10.1029/2018JD028552)
- Ben-Gai T, Bitan A, Manes A, Alpert P, Kushnir Y 2001 Temperature and surface pressure anomalies in Israel and the North Atlantic Oscillation. *Theor. Appl. Climatol.* 69, 171–177. doi: [10.1007/s00704-017-0023](https://doi.org/10.1007/s00704-017-0023)
- Ben Othman H, Leboulanger C, Le Floc'h E, Mabrouk HH, Sakka Hlaili A 2012 Toxicity of benz[a]anthracene and fluoranthene to marine phytoplankton in culture: Does cell size really matter? *J. Hazard. Mater.* 243, 204–211. doi: [10.1016/j.jhazmat.2012.10.020](https://doi.org/10.1016/j.jhazmat.2012.10.020)
- Benabdelaoui A, Taleb A, Probst JL, Belaïdi N, Probst A 2018 Anthropogenic contribution and influencing factors on metal features in fluvial sediments from a semi-arid Mediterranean river basin (Tafna River, Algeria): A multi-indices approach. *Sci. Total Environ.* 626, 899–914. doi: [10.1016/j.scitotenv.2018.01.107](https://doi.org/10.1016/j.scitotenv.2018.01.107)
- Benas N, Chrysoulakis N 2015 Estimation of the Land Surface Albedo Changes in the Broader Mediterranean Area, Based on 12 Years of Satellite Observations. *Remote Sens.* 7, 16150–16163. doi: [10.3390/rs71215816](https://doi.org/10.3390/rs71215816)
- Bengtsson L, Hodges KI, Roeckner E 2006 Storm tracks and climate change. *J. Clim.* 19.
- Beniston M, Farinotti D, Stoffel M, Andreassen LM, Coppola E et al. 2018 The European mountain cryosphere: a review of its current state, trends, and future challenges. *Cryosph.* 12, 759–794. doi: [10.5194/tc-12-759-2018](https://doi.org/10.5194/tc-12-759-2018)
- Beniston M, Stephenson DB, Christensen OB, Ferro CAT, Frei C et al. 2007 Future extreme events in European climate: an exploration of regional climate model projections. *Clim. Change* 81, 71–95. doi: [10.1007/s10584-006-9226-z](https://doi.org/10.1007/s10584-006-9226-z)
- Benoit G, Comeau A 2005 *A Sustainable Future for the Mediterranean*. London: Routledge doi: [10.4324/9781849770323](https://doi.org/10.4324/9781849770323)
- Bensoussan N, Chiggiato J, Buongiorno Nardelli B, Pisano A, Garrabou J 2019 Insights on 2017 Marine Heat Waves in the Mediterranean Sea. In: Copernicus Marine Service Ocean State Report, Issue 3. *J. Oper. Oceanogr.* 12, S26–S30. doi: [10.1080/1755876X.2019.1633075](https://doi.org/10.1080/1755876X.2019.1633075)
- Berndtsson R, Jebari S, Hashemi H, Wessels J 2016 Traditional irrigation techniques in MENA with focus on Tunisia. *Hydrol. Sci. J.* 61, 1346–1357. doi: [10.1080/0262667.2016.1165349](https://doi.org/10.1080/0262667.2016.1165349)
- Berrojalbiz N, Dachs J, Ojeda MJ, Valle MC, Castro-Jiménez J et al. 2011 Biogeochemical and physical controls on concentrations of polycyclic aromatic hydrocarbons in water and plankton of the Mediterranean and Black Seas. *Global Biogeochem. Cycles* 25, 1–14. doi: [10.1029/2010GB003775](https://doi.org/10.1029/2010GB003775)
- Berthou S, Kendon EJ, Chan SC, Ban N, Leutwyler D et al. 2018 Pan-European climate at convection-permitting scale: a model intercomparison study. *Clim. Dyn.*, 1–25. doi: [10.1007/s00382-018-4114-6](https://doi.org/10.1007/s00382-018-4114-6)
- Béthoux JP, Gentili B 1999 Functioning of the Mediterranean Sea: past and present changes related to freshwater input and climate changes. *J. Mar. Syst.* 20, 33–47. doi: [10.1016/S0924-7963\(98\)00069-4](https://doi.org/10.1016/S0924-7963(98)00069-4)
- Béthoux JP, Gentili B, Morin P, Nicolas E, Pierre C et al. 1999 The Mediterranean Sea: a miniature ocean for climatic and environmental studies and a key for the climatic functioning of the North Atlantic. *Prog. Oceanogr.* 44, 131–146. doi: [10.1016/S0079-6611\(99\)00023-3](https://doi.org/10.1016/S0079-6611(99)00023-3)
- Béthoux JP, Morin P, Chaumery C, Connan O, Gentili B et al. 1998 Nutrients in the Mediterranean Sea, mass balance and statistical analysis of concentrations with respect to environmental change. *Mar. Chem.* 63, 155–169. doi: [10.1016/S0304-4203\(98\)00059-0](https://doi.org/10.1016/S0304-4203(98)00059-0)
- Bett PE, Thornton HE, Clark RT 2013 European wind variability over 140 yr. *Adv. Sci. Res.* 10, 51–58. doi: [10.5194/asr-10-51-2013](https://doi.org/10.5194/asr-10-51-2013)
- Beuvier J, Sevault F, Herrmann M, Kontoyiannis H, Ludwig W et al. 2010 Modeling the Mediterranean Sea interannual variability during 1961–2000: Focus on the Eastern Mediterranean Transient. *JGR Ocean.* 115, C08017. doi: [10.1029/2009JC005950](https://doi.org/10.1029/2009JC005950)
- Bianchi CN 2007 Biodiversity issues for the forthcoming tropical Mediterranean Sea. In: *Biodiversity in Enclosed Seas and Artificial Marine Habitats* Developments in {Hydrobiology}, eds. Relini G, Ryland J (Springer Netherlands), 7–21. doi: [10.1007/s10750-006-0469-5](https://doi.org/10.1007/s10750-006-0469-5)
- Biel-Maeso M, Baena-Nogueras RM, Corada-Fernández C, Lara-Martín PA 2018 Occurrence, distribution and environmental risk of pharmaceutically active compounds (PhACs) in coastal and ocean waters from the Gulf of Cadiz (SW Spain). *Sci. Total Environ.* 612, 649–659. doi: [10.1016/j.scitotenv.2017.08.279](https://doi.org/10.1016/j.scitotenv.2017.08.279)
- Bilal M, Adeel M, Rasheed T, Zhao Y, Iqbal HMN 2019 Emerging contaminants of high concern and their enzyme-assisted biodegradation – A review. *Env. Int.* 124, 336–353. doi: [10.1016/j.envint.2019.01.011](https://doi.org/10.1016/j.envint.2019.01.011)
- Bilal M, Rasheed T, Sosa-Hernández JE, Raza A, Nabeel F et al. 2018 Biosorption: An interplay between marine algae and potentially toxic elements-A review. *Mar. Drugs* 16, 65. doi: [10.3390/md16020065](https://doi.org/10.3390/md16020065)
- Bilbao J, Román R, de Miguel A 2019 Temporal and Spatial Variability in Surface Air Temperature and Diurnal Temperature Range in Spain over the Period 1950–2011. *Climate* 7, 16. doi: [10.3390/cli7010016](https://doi.org/10.3390/cli7010016)
- Bladé I, Fortuny D, van Oldenborgh GJ, Liebmann B 2012 The summer North Atlantic Oscillation in CMIP3 models and related uncertainties in projected summer drying in Europe. *JGR Atmos.* 117. doi: [10.1029/2012JD017816](https://doi.org/10.1029/2012JD017816)
- Blondel J 2006 The ‘design’ of Mediterranean landscapes: a millennial story of humans and ecological systems during the historic period. *Hum. Ecol.* 34, 713–729. doi: [10.1007/s10745-006-9030-4](https://doi.org/10.1007/s10745-006-9030-4)
- Boberg F, Christensen JH 2012 Overestimation of Mediterranean summer temperature projections due to model deficiencies. *Nat. Clim. Chang.* 2, 433–436. doi: [10.1038/nclimate1454](https://doi.org/10.1038/nclimate1454)
- Boé J, Somot S, Corre L, Nabat P 2020 Large discrepancies in summer climate change over Europe as projected by glob-

- al and regional climate models: causes and consequences. *Clim. Dyn.* 54, 2981–3002. doi: [10.1007/s00382-020-05153-1](https://doi.org/10.1007/s00382-020-05153-1)
- Boé J, Terray L 2014 Land-sea contrast, soil-atmosphere and cloud-temperature interactions: interplays and roles in future summer European climate change. *Clim. Dyn.* 42, 683–699. doi: [10.1007/s00382-013-1868-8](https://doi.org/10.1007/s00382-013-1868-8)
- Borghini M, Bryden HL, Schroeder K, Sparnocchia S, Vetrano A 2014 The Mediterranean is becoming saltier. *Ocean Sci.* 10, 693–700. doi: [10.5194/os-10-693-2014](https://doi.org/10.5194/os-10-693-2014)
- Borrego C, Monteiro A, Sá E, Carvalho A, Coelho D et al. 2012 Reducing NO₂ pollution over urban areas: Air quality modelling as a fundamental management tool. *Water, Air, Soil Pollut.* 223, 5307–5320. doi: [10.1007/s11270-012-1281-7](https://doi.org/10.1007/s11270-012-1281-7)
- Bouchouicha-Smida D, Lundholm N, Sahraoui I, Lambert C, Mabrouk HH et al. 2015 Detection of domoic acid in *Mytilus galloprovincialis* and *Ostrea edulis* linked to the presence of *Nitzschia bizertensis* in Bizerte Lagoon (SW Mediterranean). *Estuar. Coast. Shelf Sci.* 165, 270–278. doi: [10.1016/j.ecss.2015.05.029](https://doi.org/10.1016/j.ecss.2015.05.029)
- Boudouresque C-F, Verlaque M 2010 Is global warming involved in the success of seaweed introductions in the Mediterranean Sea?, in *Seaweeds and their role in globally changing environments*, eds. Israel A, Einav R, Seckbach J (Springer Netherlands), 31–50. doi: [10.1007/978-90-481-8569-6_3](https://doi.org/10.1007/978-90-481-8569-6_3)
- Boutov D, Peliz Á, Miranda PMA, Soares PMM, Cardoso RM et al. 2014 Inter-annual variability and long term predictability of exchanges through the Strait of Gibraltar. *Glob. Planet. Change* 114, 23–37. doi: [10.1016/j.gloplacha.2013.12.009](https://doi.org/10.1016/j.gloplacha.2013.12.009)
- Bowman DMJS, Williamson GJ, Abatzoglou JT, Kolden CA, Cochrane MA et al. 2017 Human exposure and sensitivity to globally extreme wildfire events. *Nat. Ecol. Evol.* 1, 1–6. doi: [10.1038/s41559-016-0058](https://doi.org/10.1038/s41559-016-0058)
- Boxall ABA, Hardy A, Beulke S, Boucard T, Burgin L et al. 2009 Impacts of Climate Change on Indirect Human Exposure to Pathogens and Chemicals from Agriculture. *Environ. Health Perspect.* 117, 508–514. doi: [10.1289/ehp.0800084](https://doi.org/10.1289/ehp.0800084)
- Bradley BA, Blumenthal DM, Early R, Grosholz ED, Lawler JJ et al. 2012 Global change, global trade, and the next wave of plant invasions. *Front. Ecol. Environ.* 10, 20–28. doi: [10.1890/110145](https://doi.org/10.1890/110145)
- Branco M, Battisti A, Mendel Z 2016 Foliage Feeding Invasive Insects: Defoliators and Gall Makers, in *Insects and Diseases of Mediterranean Forest Systems*, eds. Paine T, Lieutier F (Cham: Springer International Publishing), 211–238. doi: [10.1007/978-3-319-24744-1_8](https://doi.org/10.1007/978-3-319-24744-1_8)
- Bras A, Avtzis DN, Kenis M, Li H, Vétek G et al. 2019 A complex invasion story underlies the fast spread of the invasive box tree moth (*Cydalima perspectalis*) across Europe. *J. Pest Sci.* (2004). doi: [10.1007/s10340-019-01111-x](https://doi.org/10.1007/s10340-019-01111-x)
- Bregaglio S, Hossard L, Cappelli G, Resmond R, Bocchi S et al. 2017 Identifying trends and associated uncertainties in potential rice production under climate change in Mediterranean areas. *Agric. For. Meteorol.* 237–238, 219–232. doi: [10.1016/J.AGRFORMAT.2017.02.015](https://doi.org/10.1016/J.AGRFORMAT.2017.02.015)
- Brilli L, Lugato E, Moriondo M, Gioli B, Toscano P et al. 2019 Carbon sequestration capacity and productivity responses of Mediterranean olive groves under future climates and management options. *Mitig. Adapt. Strateg. Glob. Chang.* 24, 467–491. doi: [10.1007/s11027-018-9824-x](https://doi.org/10.1007/s11027-018-9824-x)
- Brook BW, Sodhi NS, Bradshaw CJA 2008 Synergies among extinction drivers under global change. *Trends Ecol. Evol.* 23, 453–460. doi: [10.1016/J.TREE.2008.03.011](https://doi.org/10.1016/J.TREE.2008.03.011)
- Brown RD, Mote PW 2009 The response of Northern hemisphere snow cover to a changing climate. *J. Clim.* 22, 2124–2145. doi: [10.1175/2008JCLI2665.1](https://doi.org/10.1175/2008JCLI2665.1)
- Brundu G 2013 Invasive Alien Plants in Protected Areas in Mediterranean Islands: Knowledge Gaps and Main Threats, in *Plant Invasions in Protected Areas*, eds. Foxcroft LC, Pysek P, Richardson DM, Genovesi P (Dordrecht: Springer Netherlands), 395–422. doi: [10.1007/978-94-007-7750-7_18](https://doi.org/10.1007/978-94-007-7750-7_18)
- Brunetti M, Maugeri M, Monti F, Nanni T 2006 Temperature and precipitation variability in Italy in the last two centuries from homogenised instrumental time series. *Int. J. Climatol.* 26, 345–381. doi: [10.1002/joc.1251](https://doi.org/10.1002/joc.1251)
- Buccolieri A, Buccolieri G, Cardellichio N, Dell'atti A, di Leo A et al. 2006 Heavy metals in marine sediments of Taranto Gulf (Ionian Sea, Southern Italy). *Mar. Chem.* 99, 227–235. doi: [10.1016/j.marchem.2005.09.009](https://doi.org/10.1016/j.marchem.2005.09.009)
- Bulleri F, Benedetti-Cecchi L 2008 Facilitation of the introduced green alga *Caulerpa racemosa* by resident algal turfs: experimental evaluation of underlying mechanisms. *Mar. Ecol. Prog. Ser.* 364, 77–86. doi: [10.3354/meps07484](https://doi.org/10.3354/meps07484)
- Bustamante MA, Helmer EH, Schill S, Belnap J, Brown LK et al. 2018 Direct and indirect drivers of change in biodiversity and nature's contributions to people, in *IPBES (2018): The IPBES regional assessment report on biodiversity and ecosystem services for the Americas*, eds. Rice J, Seixas CS, Zaccagnini ME, Bedoya-Gaitán M, Valderrama N (Bonn, Germany: Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services), 335–509.
- Cabezas S, Carrete M, Tella JL, Marchant TA, Bortolotti GR 2013 Differences in acute stress responses between wild-caught and captive-bred birds: a physiological mechanism contributing to current avian invasions? *Biol. Invasions* 15, 521–527. doi: [10.1007/s10530-012-0304-z](https://doi.org/10.1007/s10530-012-0304-z)
- Cabral JPS 2010 Water Microbiology. Bacterial Pathogens and Water. *Int. J. Environ. Res. Public Health* 7, 3657–3703. doi: [10.3390/ijerph7103657](https://doi.org/10.3390/ijerph7103657)
- Calafat FM, Jordà G, Marcos M, Gomis D 2012 Comparison of Mediterranean sea level variability as given by three baroclinic models. *JGR Ocean.* 117. doi: [10.1029/2011JC007277](https://doi.org/10.1029/2011JC007277)
- Caloiero T, Veltri S, Caloiero P, Frustaci F 2018 Drought Analysis in Europe and in the Mediterranean Basin Using the Standardized Precipitation Index. *Water* 10, 1043. doi: [10.3390/w10081043](https://doi.org/10.3390/w10081043)
- Camarero JJ, Gazol A, Tardif JC, Conciatori F 2015 Attributing forest responses to global-change drivers: limited evidence of a CO₂-fertilization effect in Iberian pine growth. *J. Biogeogr.* 42, 2220–2233. doi: [10.1111/jbi.12590](https://doi.org/10.1111/jbi.12590)
- Camera C, Bruggeman A, Hadjinicolaou P, Michaelides S, Lange MA 2017 Evaluation of a spatial rainfall generator for generating high resolution precipitation projections over orographically complex terrain. *Stoch. Environ. Res. Risk Assess.* 31, 103–116. doi: [10.1007/s00382-016-2180-0](https://doi.org/10.1007/s00382-016-2180-0)

- Assess. 31, 757–773. doi: [10.1007/s00477-016-1239-1](https://doi.org/10.1007/s00477-016-1239-1)
- Cameron EK, Vilà M, Cabeza M 2016 Global meta-analysis of the impacts of terrestrial invertebrate invaders on species, communities and ecosystems. *Glob. Ecol. Biogeogr.* 25, 596–606. doi: [10.1111/geb.12436](https://doi.org/10.1111/geb.12436)
- Campins J, Genovés A, Picornell MA, Jansà A 2011 Climatology of Mediterranean cyclones using the ERA-40 dataset. *Int. J. Climatol.* 31, 1596–1614. doi: [10.1002/joc.2183](https://doi.org/10.1002/joc.2183)
- Campo J, Masiá A, Blasco C, Picó Y 2013 Occurrence and removal efficiency of pesticides in sewage treatment plants of four Mediterranean River Basins. *J. Hazard. Mater.* 263, 146–157. doi: [10.1016/j.jhazmat.2013.09.061](https://doi.org/10.1016/j.jhazmat.2013.09.061)
- Camuffo D, Bertolin C, Barriendos M, Domínguez-Castro F, Cocheo C et al. 2010 500-Year temperature reconstruction in the Mediterranean Basin by means of documentary data and instrumental observations. *Clim. Change* 101, 169–199. doi: [10.1007/s10584-010-9815-8](https://doi.org/10.1007/s10584-010-9815-8)
- Cannarsa S, Abete MC, Zanardi M, Squadrone S 2014 Polycyclic aromatic hydrocarbons (PAH) in marine sediment of the northwestern Mediterranean Sea (Italy). <http://blackmed-itjournal.org/wp-content/uploads/137-141-Vol20No2-Cannarsa.pdf> [Accessed April 17, 2019]
- Cardador L, Tella JL, Anadón JD, Abellán P, Carrete M 2019 The European trade ban on wild birds reduced invasion risks. *Conserv. Lett.*, e12631. doi: [10.1111/conl.12631](https://doi.org/10.1111/conl.12631)
- Cardinale M, Osio GC, Scarella G 2017 Mediterranean Sea: a failure of the European fisheries management system, in *Frontiers in Marine Science* (Frontiers Media SA). doi: [10.3389/fmars.2017.00072](https://doi.org/10.3389/fmars.2017.00072)
- Carillo A, Sannino G, Artale V, Ruti PM, Calmanti S et al. 2012 Steric sea level rise over the Mediterranean Sea: present climate and scenario simulations. *Clim. Dyn.* 39, 2167–2184. doi: [10.1007/s00382-012-1369-1](https://doi.org/10.1007/s00382-012-1369-1)
- Carlton JT 1989 Man's Role in Changing the Face of the Ocean: Biological Invasions and Implications for Conservation of Near-Shore Environments. *Conserv. Biol.* 3, 265–273. doi: [10.2307/2386170](https://doi.org/10.2307/2386170)
- Carlton JT, Chapman JW, Geller JB, Miller JA, Carlton DA et al. 2017 Tsunami-driven rafting: Transoceanic species dispersal and implications for marine biogeography. *Science (80-.J.* 357, 1402–1406. doi: [10.1126/science.aoa1498](https://doi.org/10.1126/science.aoa1498)
- Carranza ML, Drius M, Malavasi M, Frate L, Stanisci A et al. 2018 Assessing land take and its effects on dune carbon pools. An insight into the Mediterranean coastline. *Ecol. Indic.* 85, 951–955. doi: [10.1016/j.ecolind.2017.10.052](https://doi.org/10.1016/j.ecolind.2017.10.052)
- Carrer D, Pique G, Ferlicoq M, Ceamanos X, Ceschia E 2018 What is the potential of cropland albedo management in the fight against global warming? A case study based on the use of cover crops. *Environ. Res. Lett.* 13, 044030. doi: [10.1088/1748-9326/aab650](https://doi.org/10.1088/1748-9326/aab650)
- Carrete M, Tella J 2008 Wild-bird trade and exotic invasions: a new link of conservation concern? *Front. Ecol. Environ.* 6, 207–211. doi: [10.1890/070075](https://doi.org/10.1890/070075)
- Carrete M, Tella JL 2015 Rapid loss of antipredatory behaviour in captive-bred birds is linked to current avian invasions. *Sci. Rep.* 5, e18274. doi: [10.1038/srep18274](https://doi.org/10.1038/srep18274)
- Carta A, Taboada T, Müller JV 2018 Diachronic analysis using aerial photographs across fifty years reveals significant land use and vegetation changes on a Mediterranean island. *Appl. Geogr.* 98, 78–86. doi: [10.1016/j.apgeog.2018.07.010](https://doi.org/10.1016/j.apgeog.2018.07.010)
- Castellanos-Frías E, García de León D, Pujadas-Salva A, Dorado J, González-Andújar JL 2014 Potential distribution of *Avena sterilis* L. in Europe under climate change. *Ann. Appl. Biol.* 165, 53–61. doi: [10.1111/aab.12117](https://doi.org/10.1111/aab.12117)
- Castro-Díez P, González-Muñoz N, Alonso AM, Gallardo A, Poorter L 2009 Effects of exotic invasive trees on nitrogen cycling: A case study in Central Spain. *Biol. Invasions* 11, 1973–1986. doi: [10.1007/s10530-008-9374-3](https://doi.org/10.1007/s10530-008-9374-3)
- Castro-Díez Y, Pozo-Vázquez D, Rodrigo FS, Esteban-Parra MJ 2002 NAO and winter temperature variability in southern Europe. *Geophys. Res. Lett.* 29, 1-1–1-4. doi: [10.1029/2001gl014042](https://doi.org/10.1029/2001gl014042)
- Cavicchia L, von Storch H, Gualdi S 2014 Mediterranean Tropical-Like Cyclones in Present and Future Climate. *J. Clim.* 27, 7493–7501. doi: [10.1175/jcli-d-14-00339.1](https://doi.org/10.1175/jcli-d-14-00339.1)
- Cazenave A, WCRP Global Sea Level Budget Group 2018 Global sea-level budget 1993–present. *Earth Syst. Sci. Data* 10, 1551–1590. doi: [10.5194/essd-10-1551-2018](https://doi.org/10.5194/essd-10-1551-2018)
- Cancapapa A, Masiá A, Andreu V, Picó Y 2015 Spatio-temporal patterns of pesticide residues in the Turia and Júcar Rivers (Spain). *Sci. Total Environ.* 540, 200–210. doi: [10.1016/j.scitotenv.2015.06.063](https://doi.org/10.1016/j.scitotenv.2015.06.063)
- CEAM 2019 Mediterranean Sea Surface Temperature report (Summer 2019). doi: [10.13140/RG.2.2.23375.23209](https://doi.org/10.13140/RG.2.2.23375.23209)
- Cebrián E, Tomas F, López-Sendino P, Vilà M, Ballesteros E 2018 Biodiversity influences invasion success of a facultative epiphytic seaweed in a marine forest. *Biol. Invasions* 20, 2839–2848. doi: [10.1007/s10530-018-1736-x](https://doi.org/10.1007/s10530-018-1736-x)
- Ceccherelli G, Piazzi L, Balata D 2002 Spread of introduced *Caulerpa* species in macroalgal habitats. *J. Exp. Mar. Bio. Ecol.* 280, 1–11. doi: [10.1016/s0022-0981\(02\)00336-2](https://doi.org/10.1016/s0022-0981(02)00336-2)
- Ceccherini G, Russo S, Ameztoy I, Marchese AF, Carmo-na-Moreno C 2017 Heat waves in Africa 1981–2015, observations and reanalysis. *Nat. Hazards Earth Syst. Sci.* 17, 115–125. doi: [10.5194/nhess-2016-90](https://doi.org/10.5194/nhess-2016-90)
- Cecchi P, Garrido M, Collos Y, Pasqualini V 2016 Water flux management and phytoplankton communities in a Mediterranean coastal lagoon. Part II: Mixotrophy of dinoflagellates as an adaptive strategy? *Mar. Pollut. Bull.* 108, 120–133. doi: [10.1016/j.marpolbul.2016.04.041](https://doi.org/10.1016/j.marpolbul.2016.04.041)
- Ceglar A, Zampieri M, Toreti A, Dentener FJ 2019 Observed Northward Migration of Agro-Climate Zones in Europe Will Further Accelerate Under Climate Change. *Earth's Futur.* 7, 1088–1101. doi: [10.1029/2019ef001178](https://doi.org/10.1029/2019ef001178)
- Celesti-Grapow L, Bassi L, Brundu G, Camarda I, Carli E et al. 2016 Plant invasions on small Mediterranean islands: An overview. *Plant Biosyst. - An Int. J. Deal. with all Asp. Plant Biol.* 150, 1119–1133. doi: [10.1080/11263504.2016.1218974](https://doi.org/10.1080/11263504.2016.1218974)
- Cerdan O, Le Bissonnais Y, Govers G, Lecomte V, van Oost K et al. 2004 Scale effect on runoff from experimental plots to catchments in agricultural areas in Normandy. *J. Hydrol.* 299, 4–14. doi: [10.1016/J.JHYDROL.2004.02.017](https://doi.org/10.1016/J.JHYDROL.2004.02.017)
- Chang EKM, Guo Y, Xia X 2012 CMIP5 multimodel ensemble

- projection of storm track change under global warming. *JGR Atmos.* 117. doi: [10.1029/2012jd018578](https://doi.org/10.1029/2012jd018578)
- Chatha SAS, Asgher M, Iqbal HMN 2017 Enzyme-based solutions for textile processing and dye contaminant biodegradation—a review. *Environ. Sci. Pollut. Res.* 24, 14005–14018. doi: [10.1007/s11356-017-8998-1](https://doi.org/10.1007/s11356-017-8998-1)
- Cheggour A, Simonneaux V, Roose E 2012 Effets des plantations forestières sur banquettes sur le ruissellement et l'érosion par rapport aux parcours dans les montagnes semi-arides du Haut-Atlas de Marrakech (Maroc), in *Lutte antiérosive: réhabilitation des sols tropicaux et protection contre les pluies exceptionnelles*, eds. Roose E, Duchaoufou H (Marseille, France).
- Cheminée A, Sala E, Pastor J, Bodilis P, Thiriet P et al. 2013 Nursery value of *Cystoseira* forests for Mediterranean rocky reef fishes. *J. Exp. Mar. Bio. Ecol.* 442, 70–79. doi: [10.1016/j.jembe.2013.02.003](https://doi.org/10.1016/j.jembe.2013.02.003)
- Chen Y, Paytan A, Chase Z, Measures C, Beck AJ et al. 2008 Sources and fluxes of atmospheric trace elements to the Gulf of Aqaba, Red Sea. *JGR Atmos.* 113. doi: [10.1029/2007JD009110](https://doi.org/10.1029/2007JD009110)
- Cheung WWL, Lam VWY, Sarmiento JL, Kearney K, Watson R et al. 2009 Projecting global marine biodiversity impacts under climate change scenarios. *Fish Fish.* 10, 235–251. doi: [10.1111/j.1467-2979.2008.00315.x](https://doi.org/10.1111/j.1467-2979.2008.00315.x)
- Cheung WWL, Lam VWY, Sarmiento JL, Kearney K, Watson R et al. 2010 Large-scale redistribution of maximum fisheries catch potential in the global ocean under climate change. *Glob. Chang. Biol.* 16, 24–35. doi: [10.1111/j.1365-2486.2009.01995.x](https://doi.org/10.1111/j.1365-2486.2009.01995.x)
- Cheung WWL, Sarmiento JL, Dunne J, Frölicher TL, Lam VWY et al. 2013a Shrinking of fishes exacerbates impacts of global ocean changes on marine ecosystems. *Nat. Clim. Chang.* 3, 254–258. doi: [10.1038/nclimate1691](https://doi.org/10.1038/nclimate1691)
- Cheung WWL, Watson R, Pauly D 2013b Signature of ocean warming in global fisheries catch. *Nature* 497, 365–368. doi: [10.1038/nature12156](https://doi.org/10.1038/nature12156)
- Chiaccio M, Wild M 2010 Influence of NAO and clouds on long-term seasonal variations of surface solar radiation in Europe. *JGR Atmos.* 115, D00D22. doi: [10.1029/2009JD012182](https://doi.org/10.1029/2009JD012182)
- Cho S-H, Richmond-Bryant J, Thornburg J, Portzer J, Vanderpool R et al. 2011 A literature review of concentrations and size distributions of ambient airborne Pb-containing particulate matter. *Atmos. Environ.* 45, 5005–5015. doi: [10.1016/J.ATMOSENV.2011.05.009](https://doi.org/10.1016/J.ATMOSENV.2011.05.009)
- Christensen JH, Carter TR, Giorgi F 2002 PRUDENCE employs new methods to assess European climate change. *Eos, Trans. Am. Geophys. Union* 83, 147. doi: [10.1029/2002eo000094](https://doi.org/10.1029/2002eo000094)
- Christensen JH, Hewitson B, Busuioc A, Chen A, Gao X et al. 2007 Regional Climate Projections, in *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Solomon S, Qin D, Manning M, Chen Z, Marquis M et al., 847–940.
- Chu Y, Salles C, Tournoud M-G, Got P, Troussellier M et al. 2011 Faecal bacterial loads during flood events in Northwestern Mediterranean coastal rivers. *J. Hydrol.* 405, 501–511. doi: [10.1016/j.jhydrol.2011.05.047](https://doi.org/10.1016/j.jhydrol.2011.05.047)
- Chuine I, Morin X, Sonié L, Collin C, Fabreguettes J et al. 2012 Climate change might increase the invasion potential of the alien C₄ grass *Setaria parviflora* (Poaceae) in the Mediterranean Basin. *Divers. Distrib.* 18, 661–672. doi: [10.1111/j.1472-4642.2011.00880.x](https://doi.org/10.1111/j.1472-4642.2011.00880.x)
- Church JA, Clark PU, Cazenave A, Gregory JM, Jevrejeva S et al. 2013 Sea level change. PM Cambridge University Press
- Chytrý M, Maskell LC, Pino J, Pyšek P, Vilà M et al. 2008 Habitat invasions by alien plants: a quantitative comparison among Mediterranean, subcontinental and oceanic regions of Europe. *J. Appl. Ecol.* 45, 448–458. doi: [10.1111/j.1365-2664.2007.01398.x](https://doi.org/10.1111/j.1365-2664.2007.01398.x)
- Ciminiello P, Dell'Aversano C, Dello Iacovo E, Fattorusso E, Forino M et al. 2014 First finding of *Ostreopsis cf. ovata* toxins in marine aerosols. *Environ. Sci. Technol.* 48, 3532–3540. doi: [10.1021/es405617d](https://doi.org/10.1021/es405617d)
- Cincinelli A, Martellini T, Guerranti C, Scopetani C, Chelazzi D et al. 2019 A potpourri of microplastics in the sea surface and water column of the Mediterranean Sea. *TrAC - Trends Anal. Chem.* 110, 321–326. doi: [10.1016/j.trac.2018.10.026](https://doi.org/10.1016/j.trac.2018.10.026)
- Cini A, Anfora G, Escudero-Colomar LA, Grassi A, Santosuosso U et al. 2014 Tracking the invasion of the alien fruit pest *Drosophila suzukii* in Europe. *J. Pest Sci.* (2004). 87, 559–566. doi: [10.1007/s10340-014-0617-z](https://doi.org/10.1007/s10340-014-0617-z)
- Clarke L, Jiang K, Akimoto K, Babiker M, Blanford G et al. 2014 Assessing Transformation Pathways, in *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Edenhofer O, Pichs-Madruga R, Sokona Y, Farahani E, Kadner S et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press).
- Clavero M, Brotons L, Pons P, Sol D 2009 Prominent role of invasive species in avian biodiversity loss. *Biol. Conserv.* 142, 2043–2049. doi: [10.1016/j.biocon.2009.03.034](https://doi.org/10.1016/j.biocon.2009.03.034)
- Clotet M, Basnou C, Bagaria G, Pino J 2016 Contrasting historical and current land-use correlation with diverse components of current alien plant invasions in Mediterranean habitats. *Biol. Invasions* 18, 2897–2909. doi: [10.1007/s10530-016-1181-7](https://doi.org/10.1007/s10530-016-1181-7)
- Cofiño AS, San-Martín D, Gutiérrez JM 2007 A web portal for regional projection of weather forecast using GRID middleware, in *Lecture Notes in Computer Science*, 82–89.
- Coll M, Piroddi C, Steenbeek J, Kaschner K, Ben Rais Lasram F et al. 2010 The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS One* 5, e11842. doi: [10.1371/journal.pone.0011842](https://doi.org/10.1371/journal.pone.0011842)
- Collins M, Knutti R, Arblaster J, Dufresne J-L, Fichefet T et al. 2013 Long-term Climate Change: Projections, Commitments and Irreversibility, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK et al. (Cambridge, United Kingdom and New York, NY, USA), 1029–1136. doi: [10.1017/CBO9781107415324.024](https://doi.org/10.1017/CBO9781107415324.024)

- Colloca F, Cardinale M, Maynou F, Giannoulaki M, Scarella G et al. 2013 Rebuilding Mediterranean fisheries: a new paradigm for ecological sustainability. *Fish Fish.* 14, 89–109. doi: [10.1111/j.1467-2979.2011.00453.x](https://doi.org/10.1111/j.1467-2979.2011.00453.x)
- Colloca F, Scarella G, Libralato S 2017 Recent Trends and Impacts of Fisheries Exploitation on Mediterranean Stocks and Ecosystems. *Front. Mar. Sci.* 4. doi: [10.3389/fmars.2017.00244](https://doi.org/10.3389/fmars.2017.00244)
- Colucci RR, Guglielmin M 2019 Climate change and rapid ice melt: Suggestions from abrupt permafrost degradation and ice melting in an alpine ice cave. *Prog. Phys. Geogr. Earth Environ.* 43. doi: [10.1177%2F030913319846056](https://doi.org/10.1177%2F030913319846056)
- Colucci RR, Žebre M 2016 Late Holocene evolution of glaciers in the southeastern Alps. *J. Maps* 12, 289–299. doi: [10.1080/17445647.2016.1203216](https://doi.org/10.1080/17445647.2016.1203216)
- Compa M, Alomar C, Wilcox C, van Sebille E, Lebreton L et al. 2019 Risk assessment of plastic pollution on marine diversity in the Mediterranean Sea. *Sci. Total Environ.* 678, 188–196. doi: [10.1016/j.scitotenv.2019.04.355](https://doi.org/10.1016/j.scitotenv.2019.04.355)
- Constán-Nava S, Soliveres S, Torices R, Serra L, Bonet A 2015 Direct and indirect effects of invasion by the alien tree *Ailanthus altissima* on riparian plant communities and ecosystem multifunctionality. *Biol. Invasions* 17, 1095–1108. doi: [10.1007/s10530-014-0780-4](https://doi.org/10.1007/s10530-014-0780-4)
- Conte D, Lionello P 2013 Characteristics of large positive and negative surges in the Mediterranean Sea and their attenuation in future climate scenarios. *Glob. Planet. Change* 111, 159–173. doi: [10.1016/j.gloplacha.2013.09.006](https://doi.org/10.1016/j.gloplacha.2013.09.006)
- Conti G, Fagarazzi L 2015 Forest expansion in mountain ecosystems: “environmentalist’s dream” or societal nightmare? *Planum* 11, 1–20.
- Conversi A, Fonda Umani S, Peluso T, Molinero JC, Santojanni A et al. 2010 The Mediterranean Sea regime shift at the end of the 1980s, and intriguing parallelisms with other European basins. *PLoS One* 5, e10633. doi: [10.1371/journal.pone.0010633](https://doi.org/10.1371/journal.pone.0010633)
- Cook CM, Vardaka E, Lanaras T 2004 Toxic cyanobacteria in Greek freshwaters, 1987–2000: Occurrence, toxicity, and impacts in the Mediterranean region. *Acta Hydrochim. Hydrobiol.* 32, 107–124. doi: [10.1002/ahed.200300523](https://doi.org/10.1002/ahed.200300523)
- Cook KH, Vizy EK 2015 Detection and Analysis of an Amplified Warming of the Sahara Desert. *J. Clim.* 28, 6560–6580. doi: [10.1175/jcli-d-14-00230.1](https://doi.org/10.1175/jcli-d-14-00230.1)
- Coppola E, Sobolowski S, Pichelli E, Raffaele F, Ahrens B et al. 2020 A first-of-its-kind multi-model convection permitting ensemble for investigating convective phenomena over Europe and the Mediterranean. *Clim. Dyn.* 55, 3–34. doi: [10.1007/s00382-018-4521-8](https://doi.org/10.1007/s00382-018-4521-8)
- Corrales X, Coll M, Ofir E, Heymans JJ, Steenbeek J et al. 2018 Future scenarios of marine resources and ecosystem conditions in the Eastern Mediterranean under the impacts of fishing, alien species and sea warming. *Sci. Rep.* 8, 14284. doi: [10.1038/s41598-018-32666-x](https://doi.org/10.1038/s41598-018-32666-x)
- Corte-Real JM, Zhang X, Wang X 1995 Large-scale circulation regimes and surface climatic anomalies over the Mediterranean. *Int. J. Climatol.* 15, 1135–1150. doi: [10.1002/joc.3370151006](https://doi.org/10.1002/joc.3370151006)
- Cosentino SL, Porqueddu C, Copani V, Patané C, Testa G et al. 2014 European grasslands overview: Mediterranean region. *Grassl. Sci. Eur.* 19, 41–56.
- Coumou D, Rahmstorf S 2012 A decade of weather extremes. *Nat. Clim. Chang.* 2, 491–496. doi: [10.1038/nclimate1452](https://doi.org/10.1038/nclimate1452)
- Covas L, Senar JC, Roqué L, Quesada J 2017 Records of fatal attacks by Rose-ringed Parakeets *Psittacula krameri* on native avifauna. http://ornitologia.org/mm/file/2017_06.pdf [Accessed May 14, 2019].
- Cózar A, Sanz-Martín M, Martí E, Ignacio González-Gordillo J, Ubeda B et al. 2015 Plastic accumulation in the Mediterranean Sea. *PLoS One* 10. doi: [10.1371/journal.pone.0121762](https://doi.org/10.1371/journal.pone.0121762)
- Crain CM, Kroeker KJ, Halpern BS 2008 Interactive and cumulative effects of multiple human stressors in marine systems. *Ecol. Lett.* 11, 1304–1315. doi: [10.1111/j.1461-0248.2008.01253.x](https://doi.org/10.1111/j.1461-0248.2008.01253.x)
- Craine JM, Elmore AJ, Olson KC, Tolleson D 2010 Climate change and cattle nutritional stress. *Glob. Chang. Biol.* 16, 2901–2911. doi: [10.1111/j.1365-2486.2009.02060.x](https://doi.org/10.1111/j.1365-2486.2009.02060.x)
- Cramer W, Guiot J, Fader M, Garrabou J, Gattuso J-P et al. 2018 Climate change and interconnected risks to sustainable development in the Mediterranean. *Nat. Clim. Chang.* 8, 972–980. doi: [10.1038/s41558-018-0299-2](https://doi.org/10.1038/s41558-018-0299-2)
- Crooks JA, Chang AL, Ruiz GM 2011 Aquatic pollution increases the relative success of invasive species. *Biol. Invasions* 13, 165–176. doi: [10.1007/s10530-010-9799-3](https://doi.org/10.1007/s10530-010-9799-3)
- Crooks JA, Suarez A 2006 Hyperconnectivity, invasive species, and the breakdown of barriers to dispersal, in *Connectivity Conservation*, eds. Crooks KR, Sanjayan MA (Cambridge, New York, USA: Cambridge University Press), 451–478.
- Cuartera JC 2017 Income projections for climate change research: A framework based on human capital dynamics. *Glob. Environ. Chang.* 42, 226–236. doi: [10.1016/j.gloenvcha.2015.02.012](https://doi.org/10.1016/j.gloenvcha.2015.02.012)
- Cudennec C, Leduc C, Koutsoyiannis D 2007 Dryland hydrology in Mediterranean regions—a review. *Hydrol. Sci. J.* 52, 1077–1087. doi: [10.1623/hysj.52.6.1077](https://doi.org/10.1623/hysj.52.6.1077)
- Cullen HM, DeMenocal PB 2000 North Atlantic influence on Tigris-Euphrates streamflow. *Int. J. Climatol.* 20, 853–863. doi: [10.1002/1097-0088\(20000630\)20:8<853::AID-JOC497>3.0.CO;2-M](https://doi.org/10.1002/1097-0088(20000630)20:8<853::AID-JOC497>3.0.CO;2-M)
- Cyrus J, Eeftens M, Heinrich J, Ampe C, Armengaud A et al. 2012 Variation of NO₂ and NO_x concentrations between and within 36 European study areas: Results from the ESCAPE study. *Atmos. Environ.* 62, 374–390. doi: [10.1016/j.atmosenv.2012.07.080](https://doi.org/10.1016/j.atmosenv.2012.07.080)
- D’Agostino R, Lionello P 2020 The atmospheric moisture budget in the Mediterranean: Mechanisms for seasonal changes in the Last Glacial Maximum and future warming scenario. *Quat. Sci. Rev.* 241, 106392. doi: [10.1016/j.quascirev.2020.106392](https://doi.org/10.1016/j.quascirev.2020.106392)
- D’Agostino R, Lionello P, Adam O, Schneider T 2017 Factors controlling Hadley circulation changes from the Last Glacial Maximum to the end of the 21st century. *Geophys. Res. Lett.* 44, 8585–8591. doi: [10.1002/2017gl074533](https://doi.org/10.1002/2017gl074533)
- D’Agostino R, Scambati AL, Jungclaus J, Lionello P 2020 Poleward Shift of Northern Subtropics in Winter: Time of Emergence of Zonal Versus Regional Signals. *Geophys. Res. Lett.* 47, e2020GL089325. doi: [10.1029/2020GL089325](https://doi.org/10.1029/2020GL089325)

- D'Silva MS, Anil AC, Naik RK, D'Costa PM 2012 Algal blooms: a perspective from the coasts of India. *Nat. Hazards* 63, 1225–1253. doi: [10.1007/s11069-012-0190-9](https://doi.org/10.1007/s11069-012-0190-9)
- Dafka S, Toreti A, Zanis P, Xoplaki E, Luterbacher J 2019 Twenty-first-century changes in the Eastern Mediterranean Etearians and associated midlatitude atmospheric circulation. *JGR Atmos.* 124, 12741–12754. doi: [10.1029/2019JD031203](https://doi.org/10.1029/2019JD031203)
- Dahshan H, Megahed AM, Abd-Elall AMM, Abd-El-Kader MA-G, Nabawy E et al. 2016 Monitoring of pesticides water pollution-The Egyptian River Nile. *J. Environ. Heal. Sci. Eng.* 14, 15. doi: [10.1186/s40201-016-0259-6](https://doi.org/10.1186/s40201-016-0259-6)
- Dai A 2013 Increasing drought under global warming in observations and models. *Nat. Clim. Chang.* 3, 52–58. doi: [10.1038/nclimate1633](https://doi.org/10.1038/nclimate1633)
- DAISIE 2009 *Handbook of Alien Species in Europe*. Springer Netherlands doi: [10.1007/978-1-4020-8280-1](https://doi.org/10.1007/978-1-4020-8280-1)
- Danovaro R, Fonda Umani S, Pusceddu A, Umani SF, Pusceddu A et al. 2009 Climate Change and the Potential Spreading of Marine Mucilage and Microbial Pathogens in the Mediterranean Sea. *PLoS One* 4, e7006. doi: [10.1371/journal.pone.0007006](https://doi.org/10.1371/journal.pone.0007006)
- Danti R, Della Rocca G 2017 Epidemiological History of Cypress Canker Disease in Source and Invasion Sites. *Forests* 8, 121. doi: [10.3390/f8040121](https://doi.org/10.3390/f8040121)
- Darmaraki S, Somot S, Sevault F, Nabat P 2019a Past Variability of Mediterranean Sea Marine Heatwaves. *Geophys. Res. Lett.* doi: [10.1029/2019gl082933](https://doi.org/10.1029/2019gl082933)
- Darmaraki S, Somot S, Sevault F, Nabat P, Cabos Narvaez WD et al. 2019b Future evolution of Marine Heatwaves in the Mediterranean Sea. *Clim. Dyn.* 53, 1371–1392. doi: [10.1007/s00382-019-04661-z](https://doi.org/10.1007/s00382-019-04661-z)
- Dayan U, Ricaud P, Zbinden R, Dulac F 2017 Atmospheric pollution over the eastern Mediterranean during summer - A review. *Atmos. Chem. Phys.* 17, 13233–13263. doi: [10.5194/acp-17-13233-2017](https://doi.org/10.5194/acp-17-13233-2017)
- de Giglio O, Caggiano G, Bagordo F, Barbuti G, Brigida S et al. 2017 Enteric viruses and fecal bacteria indicators to assess groundwater quality and suitability for irrigation. *Int. J. Environ. Res. Public Health* 14, 558. doi: [10.3390/ijerph14060558](https://doi.org/10.3390/ijerph14060558)
- de Montmollin B, Strahm W 2005 *The Top 50 Mediterranean Island Plants. Wild Plants At the Brink of Extinction, and What is Needed to Save Them*. IUCN/SSC Mediterranean Islands Plant Specialist Group. IUCN, Gland, Switzerland and Cambridge doi: [10.1017/s0030605306270200](https://doi.org/10.1017/s0030605306270200)
- Debolini M, Marraccini E, Dubœuf JP, Geijzendorffer IR, Guerra CA et al. 2018 Land and farming system dynamics and their drivers in the Mediterranean Basin. *Land use policy* 75, 702–710. doi: [10.1016/j.landusepol.2017.07.010](https://doi.org/10.1016/j.landusepol.2017.07.010)
- Deidun A 2018 Back with a bang – an unexpected massive bloom of *Cassiopea andromeda* (Forskaal, 1775) in the Maltese Islands, nine years after its first appearance. *Bio-Invasions Rec.* 7, 399–404. doi: [10.3391/bir.2018.7.4.07](https://doi.org/10.3391/bir.2018.7.4.07)
- Dell'Aquila A, Mariotti A, Bastin S, Calmant S, Cavicchia L et al. 2018 Evaluation of simulated decadal variations over the Euro-Mediterranean region from ENSEMBLES to Med-CORDEX. *Clim. Dyn.* 51, 857–876. doi: [10.1007/s00382-016-3143-2](https://doi.org/10.1007/s00382-016-3143-2)
- Dellink R, Chateau J, Lanzi E, Magné B 2017 Long-term economic growth projections in the Shared Socioeconomic Pathways. *Glob. Environ. Chang.* 42, 200–214. doi: [10.1016/j.gloenvcha.2015.06.004](https://doi.org/10.1016/j.gloenvcha.2015.06.004)
- Déqué M 2007 Frequency of precipitation and temperature extremes over France in an anthropogenic scenario: Model results and statistical correction according to observed values. *Glob. Planet. Change* 57, 16–26. doi: [10.1016/j.glopacha.2006.11.030](https://doi.org/10.1016/j.glopacha.2006.11.030)
- Déqué M, Somot S, Sánchez-Gómez E, Goodess CM, Jacob D et al. 2012 The spread amongst ENSEMBLES regional scenarios: Regional climate models, driving general circulation models and interannual variability. *Clim. Dyn.* 38, 951–964. doi: [10.1007/s00382-011-1053-x](https://doi.org/10.1007/s00382-011-1053-x)
- Deser C, Hurrell JW, Phillips AS 2017 The role of the North Atlantic Oscillation in European climate projections. *Clim. Dyn.* 49, 3141–3157. doi: [10.1007/s00382-016-3502-z](https://doi.org/10.1007/s00382-016-3502-z)
- Desneux N, Wajnberg E, Wyckhuys KAG, Burgio G, Arpaia S et al. 2010 Biological invasion of European tomato crops by *Tuta absoluta*: Ecology, geographic expansion and prospects for biological control. *J. Pest Sci.* (2004). 83, 197–215. doi: [10.1007/s10340-010-0321-6](https://doi.org/10.1007/s10340-010-0321-6)
- DG/ACTA 1993 Stratégie nationale pour la conservation des eaux et des sols. Publication de la direction de la conservation des eaux et des sols.
- di Castri F 1991 An ecological overview of the five regions of the world with a mediterranean climate, in *Biogeography of Mediterranean Invasions*, eds. Groves RH, di Castri F (Cambridge University Press), 3–16. doi: [10.1017/cbo9780511525544.002](https://doi.org/10.1017/cbo9780511525544.002)
- Diapouli E, Manousakas M, Vratolis S, Vasilatou V, Maggos T et al. 2017 Evolution of air pollution source contributions over one decade, derived by PM10 and PM2.5 source apportionment in two metropolitan urban areas in Greece. *Atmos. Environ.* 164, 416–430. doi: [10.1016/j.atmosenv.2017.06.016](https://doi.org/10.1016/j.atmosenv.2017.06.016)
- Diappi L 2015 City Size and Urbanization in Mediterranean Cities. *Sci. Reg.*, 129–137. doi: [10.3280/SCRE2015-001011](https://doi.org/10.3280/SCRE2015-001011)
- Diffenbaugh NS, Giorgi F 2012 Climate change hotspots in the CMIP5 global climate model ensemble. *Clim. Change* 114, 813–822. doi: [10.1007/s10584-012-0570-x](https://doi.org/10.1007/s10584-012-0570-x)
- Diffenbaugh NS, Pal JS, Giorgi F, Gao X 2007 Heat stress intensification in the Mediterranean climate change hotspot. *Geophys. Res. Lett.* 34, L11706. doi: [10.1029/2007gl030000](https://doi.org/10.1029/2007gl030000)
- Dimarchopoulou D, Stergiou KI, Tsikliras AC 2017 Gap analysis on the biology of Mediterranean marine fishes. *PLoS One* 12, e0175949. doi: [10.1371/journal.pone.0175949](https://doi.org/10.1371/journal.pone.0175949)
- Doblas-Miranda E, Alonso R, Arnan X, Bermejo V, Brotons L et al. 2017 A review of the combination among global change factors in forests, shrublands and pastures of the Mediterranean Region: Beyond drought effects. *Glob. Planet. Change* 148, 42–54. doi: [10.1016/j.glopacha.2016.11.012](https://doi.org/10.1016/j.glopacha.2016.11.012)
- Dobrynin M, Murawsky J, Yang S 2012 Evolution of the global wind wave climate in CMIP5 experiments. *Geophys. Res. Lett.* 39. doi: [10.1029/2012gl052843](https://doi.org/10.1029/2012gl052843)
- Domon G 2011 Landscape as resource: Consequences, challenges and opportunities for rural development. *Landsc. Urban Plan.* 100, 338–340. doi: [10.1016/j.landurbplan.2011.02.014](https://doi.org/10.1016/j.landurbplan.2011.02.014)

- Donat MG, Peterson TC, Brunet M, King AD, Almazroui M et al. 2014 Changes in extreme temperature and precipitation in the Arab region: long-term trends and variability related to ENSO and NAO. *Int. J. Climatol.* 34, 581–592. doi: [10.1002/joc.3707](https://doi.org/10.1002/joc.3707)
- Donat MG, Renggli D, Wild S, Alexander L V., Leckebusch GC et al. 2011 Reanalysis suggests long-term upward trends in European storminess since 1871. *Geophys. Res. Lett.* 38, n/a-n/a. doi: [10.1029/2011GL047995](https://doi.org/10.1029/2011GL047995)
- Dorn W, Dethloff K, Rinke A, Roeckner E 2003 Competition of NAO regime changes and increasing greenhouse gases and aerosols with respect to Arctic climate projections. *Clim. Dyn.* 21, 447–458. doi: [10.1007/s00382-003-0344-2](https://doi.org/10.1007/s00382-003-0344-2)
- Dosio A, Fischer EM 2018 Will Half a Degree Make a Difference? Robust Projections of Indices of Mean and Extreme Climate in Europe Under 1.5°C, 2°C, and 3°C Global Warming. *Geophys. Res. Lett.* 45, 935–944. doi: [10.1002/2017gl076222](https://doi.org/10.1002/2017gl076222)
- Douville H, Ribes A, Decharme B, Alkama R, Sheffield J 2013 Anthropogenic influence on multidecadal changes in reconstructed global evapotranspiration. *Nat. Clim. Chang.* 3, 59–62. doi: [10.1038/nclimate1632](https://doi.org/10.1038/nclimate1632)
- Drira Z, Kmihi-Megdiche S, Sahnoun H, Hammami A, Allouche N et al. 2016 Assessment of anthropogenic inputs in the surface waters of the southern coastal area of Sfax during spring [Tunisia, Southern Mediterranean Sea]. *Mar. Pollut. Bull.* 104, 355–363. doi: [10.1016/j.marpolbul.2016.01.035](https://doi.org/10.1016/j.marpolbul.2016.01.035)
- Dris R, Gasperi J, Mirande C, Mandin C, Guerrouache M et al. 2017 A first overview of textile fibers, including microplastics, in indoor and outdoor environments. *Environ. Pollut.* 221, 453–458. doi: [10.1016/J.ENVPOL.2016.12.013](https://doi.org/10.1016/J.ENVPOL.2016.12.013)
- Dris R, Gasperi J, Saad M, Mirande C, Tassin B 2016 Synthetic fibers in atmospheric fallout: A source of microplastics in the environment? *Mar. Pollut. Bull.* 104, 290–293. doi: [10.1016/J.MARPOLBUL.2016.01.006](https://doi.org/10.1016/J.MARPOLBUL.2016.01.006)
- Dubois A, Lacouture L, Feuillet C 2010 Les pesticides dans les milieux aquatiques : données 2007. *Etudes Doc. du Commissariat Général au Développement Durable* 26, 50. <https://www.statistiques.developpement-durable.gouv.fr/sites/default/files/2019-01/les-pesticides-dans-les-milieux-aquatiques-etudes-et-documents-26-juillet2010.pdf>
- Dubois C, Somot S, Calmant S, Carillo A, Déqué M et al. 2012 Future projections of the surface heat and water budgets of the Mediterranean Sea in an ensemble of coupled atmosphere-ocean regional climate models. *Clim. Dyn.* 39, 1859–1884. doi: [10.1007/s00382-011-1261-4](https://doi.org/10.1007/s00382-011-1261-4)
- Dubrovský M, Hayes M, Duce P, Trnka M, Svoboda M et al. 2014 Multi-GCM projections of future drought and climate variability indicators for the Mediterranean region. *Reg. Environ. Chang.* 14, 1907–1919. doi: [10.1007/s10113-013-0562-z](https://doi.org/10.1007/s10113-013-0562-z)
- Dullinger I, Wessely J, Bossdorf O, Dawson W, Essl F et al. 2017 Climate change will increase the naturalization risk from garden plants in Europe. *Glob. Ecol. Biogeogr.* 26, 43–53. doi: [10.1111/geb.12512](https://doi.org/10.1111/geb.12512)
- Dunić N, Vilibić I, Šepić J, Mihanović H, Sevault F et al. 2019 Performance of multi-decadal ocean simulations in the Adriatic Sea. *Ocean Model.* 134, 81–109. doi: [10.1016/j.ocemod.2019.01.006](https://doi.org/10.1016/j.ocemod.2019.01.006)
- Dunkeloh A, Jacobbeit J 2003 Circulation dynamics of Mediterranean precipitation variability 1948–98. *Int. J. Climatol.* 23, 1843–1866. doi: [10.1002/joc.973](https://doi.org/10.1002/joc.973)
- Dupire S, Curt T, Bigot S 2017 Spatio-temporal trends in fire weather in the French Alps. *Sci. Total Environ.* 595, 801–817. doi: [10.1016/j.scitotenv.2017.04.027](https://doi.org/10.1016/j.scitotenv.2017.04.027)
- Early R, González-Moreno P, Murphy ST, Day R 2018 Forecasting the global extent of invasion of the cereal pest *Spodoptera frugiperda*, the fall armyworm. *bioRxiv*, 391847. doi: [10.3897/neobiota.40.28165.suppl](https://doi.org/10.3897/neobiota.40.28165.suppl)
- EC/DG for Research and Innovation 2011 *EuroMed-2030: Long term challenges for the Mediterranean area; report of an expert group*. Luxembourg: Publ. Office of the European Union.
- Edelist D, Rilov G, Golani D, Carlton JT, Spanier E 2013 Restructuring the Sea: profound shifts in the world's most invaded marine ecosystem. *Divers. Distrib.* 19, 69–77. doi: [10.1111/ddi.12002](https://doi.org/10.1111/ddi.12002)
- EEA 1999 Environmental Indicators: Typology and Overview, Technical Report No. 25. Copenhagen, Denmark <https://www.eea.europa.eu/publications/TEC25>
- Efthymiadis D, Goodess CM, Jones PD 2011 Trends in Mediterranean gridded temperature extremes and large-scale circulation influences. *Nat. Hazards Earth Syst. Sci.* 11, 2199–2214. doi: [10.5194/nhess-11-2199-2011](https://doi.org/10.5194/nhess-11-2199-2011)
- El Ayni F, Manoli E, Cherif S, Jrad A, Assimacopoulos D et al. 2013 Deterioration of a Tunisian coastal aquifer due to agricultural activities and possible approaches for better water management. *Water Environ. J.* 27, 348–361. doi: [10.1111/j.1747-6593.2012.00354.x](https://doi.org/10.1111/j.1747-6593.2012.00354.x)
- El Bakouri H, Ouassini A, Morillo J, Usero J 2008 Pesticides in ground water beneath Loukkos perimeter, Northwest Morocco. *J. Hydrol.* 348, 270–278. doi: [10.1016/j.jhydrol.2007.10.002](https://doi.org/10.1016/j.jhydrol.2007.10.002)
- El Kenawy AM, López Moreno JL, Vicente-Serrano SM 2011 Recent trends in daily temperature extremes over north-eastern Spain (1960–2006). *Nat. Hazards Earth Syst. Sci.* 11, 2583–2603. doi: [10.5194/nhess-11-2583-2011](https://doi.org/10.5194/nhess-11-2583-2011)
- El Kenawy AM, López Moreno JL, Vicente-Serrano SM, Mekld MS 2009 Temperature trends in Libya over the second half of the 20th century. *Theor. Appl. Climatol.* 98, 1–8. doi: [10.1007/s00704-008-0089-2](https://doi.org/10.1007/s00704-008-0089-2)
- El Nemr A, Abd-Allah AMA 2003 Contamination of polycyclic aromatic hydrocarbons (PAHs) in microlayer and subsurface waters along Alexandria coast, Egypt. *Chemosphere* 52, 1711–1716. doi: [10.1016/S0045-6535\(03\)00300-X](https://doi.org/10.1016/S0045-6535(03)00300-X)
- Elbakidze M, Hahn T, Zimmermann NE, Cudlin P, Friberg N et al. 2018 Direct and indirect drivers of change in biodiversity and nature's contributions to people, in *IPBES (2018): The IPBES regional assessment report on biodiversity and ecosystem services for Europe and Central Asia*, eds. Rounsevell M, Fischer M, Torre-Marín Rando A, Mader A (Bonn, Germany: Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem services), 385–568.
- Elguindi N, Somot S, Déqué M, Ludwig W 2011 Climate change evolution of the hydrological balance of the Mediterranean, Black and Caspian Seas: impact of climate model resolution. *Clim. Dyn.* 36, 205–228. doi: [10.1007/s00382-009-0715-4](https://doi.org/10.1007/s00382-009-0715-4)

- Elton CS 1958 *The Ecology of Invasions by Animals and Plants*. Chapman and Hall, New York, US.
doi: [10.1007/978-1-4899-7214-9](https://doi.org/10.1007/978-1-4899-7214-9)
- Emanuel K 2005 Genesis and maintenance of "Mediterranean hurricanes." *Adv. Geosci.* 2, 217–220.
doi: [10.5194/adgeo-2-217-2005](https://doi.org/10.5194/adgeo-2-217-2005)
- Emara HI, Said TO, El Naggar NA, Shreadah MA 2008 Aliphatic and polycyclic hydrocarbon compounds as chemical markers for pollution sources in relation to physico-chemical characteristics of the eastern harbour (Egyptian Mediterranean Sea). *Egypt. J. Aquat. Res.* 34, 1–19.
- Engelbrecht JP, Jayanty RKM 2013 Assessing sources of airborne mineral dust and other aerosols, in Iraq. *Aeolian Res.* 9, 153–160. doi: [10.1016/J.AEOLIA.2013.02.003](https://doi.org/10.1016/J.AEOLIA.2013.02.003)
- Enriquez-Alonso A, Sánchez-Lorenzo A, Calbó J, González J-A, Norris JR 2016 Cloud cover climatologies in the Mediterranean obtained from satellites, surface observations, reanalyses, and CMIP5 simulations: validation and future scenarios. *Clim. Dyn.* 47, 249–269. doi: [10.1007/s00382-015-2834-4](https://doi.org/10.1007/s00382-015-2834-4)
- EPPO 2012a Decision-support scheme for an Express Pest Risk Analysis. *EPPO Bull.* 42, 457–462. doi: [10.1111/epp.2591](https://doi.org/10.1111/epp.2591)
- EPPO 2012b EPPO prioritization process for invasive alien plants. *EPPO Bull.* 42, 463–474. doi: [10.1111/epp.2592](https://doi.org/10.1111/epp.2592)
- Eschen R, Grégoire J-C, Hengeveld GM, de Hoop BM, Rigaux L et al. 2015 Trade patterns of the tree nursery industry in Europe and changes following findings of citrus longhorn beetle, *Anoplophora chinensis* Forster. *NeoBiota* 26, 1–20. doi: [10.3897/neobiota.26.8947](https://doi.org/10.3897/neobiota.26.8947)
- Escudero M, Querol X, Ávila A, Cuevas E 2007 Origin of the exceedances of the European daily PM limit value in regional background areas of Spain. *Atmos. Environ.* 41, 730–744. doi: [10.1016/J.ATMOSENV.2006.09.014](https://doi.org/10.1016/J.ATMOSENV.2006.09.014)
- Essid N, Boufahja F, Beyrem H, Aissa P, Mahmoudi E 2013 Effects of 17- α -estradiol on a free-living marine nematode community: A microcosm experiment. *African J. Aquat. Sci.* 38, 305–311. doi: [10.2989/16085914.2013.832652](https://doi.org/10.2989/16085914.2013.832652)
- Essl F, Kobler J 2009 Spiny invaders – Patterns and determinants of cacti invasion in Europe. *Flora - Morphol. Distrib. Funct. Ecol. Plants* 204, 485–494. doi: [10.1016/j.flora.2008.06.002](https://doi.org/10.1016/j.flora.2008.06.002)
- Etteieb S, Cherif S, Kawachi A, Han J, Elayni F et al. 2016 Combining Biological and Chemical Screenings to Assess Cytotoxicity of Emerging Contaminants in Discharges into Surface Water. *Water, Air, Soil Pollut.* 227, 341. doi: [10.1007/s11270-016-3049-y](https://doi.org/10.1007/s11270-016-3049-y)
- European Food Safety Authority 2011 The 2009 European Union Report on Pesticide Residues in Food. *EFSA J.* 9, 2430. doi: [10.2903/j.efsa.2011.2430](https://doi.org/10.2903/j.efsa.2011.2430)
- European Union 2014 Regulation (EU) No 1143/2014 of the European Parliament and of the Council of 22 October 2014 on the prevention and management of the introduction and spread of invasive alien species. *Off. J. Eur. Union* 57. doi: [10.5040/978150909568.0031](https://doi.org/10.5040/978150909568.0031)
- Evan AT, Flamant C, Lavaysse C, Kocha C, Saci A 2015 Water Vapor-Forced Greenhouse Warming over the Sahara Desert and the Recent Recovery from the Sahelian Drought. *J. Clim.* 28, 108–123. doi: [10.1175/jcli-d-14-00039.1](https://doi.org/10.1175/jcli-d-14-00039.1)
- Evangelou N, Balkanski Y, Cozic A, Hao WM, Møller AP 2014 Wildfires in Chernobyl-contaminated forests and risks to the population and the environment: A new nuclear disaster about to happen? *Environ. Int.* 73, 346–358. doi: [10.1016/J.ENVINT.2014.08.012](https://doi.org/10.1016/J.ENVINT.2014.08.012)
- Ezber Y 2018 Assessment of the changes in the Etesians in the EURO-CORDEX regional model projections. *Int. J. Climatol.* 39, 1213–1229. doi: [10.1002/joc.5872](https://doi.org/10.1002/joc.5872)
- Fabbio G, Merlo M, Tosi V 2003 Silvicultural management in maintaining biodiversity and resistance of forests in Europe—the Mediterranean region. *J. Environ. Manage.* 67, 67–76. doi: [10.1016/s0301-4797\(02\)00189-5](https://doi.org/10.1016/s0301-4797(02)00189-5)
- Faccoli M, Campo G, Perrotta G, Rassati D 2016 Two newly introduced tropical bark and ambrosia beetles (Coleoptera: Curculionidae, Scolytinae) damaging figs (*Ficus carica*) in southern Italy. *Zootaxa* 4138, 189. doi: [10.11646/zootaxa.4138.1.10](https://doi.org/10.11646/zootaxa.4138.1.10)
- Fan J, Denux O, Courtin C, Bernard A, Javal M et al. 2019 Multi-component blends for trapping native and exotic long-horn beetles at potential points-of-entry and in forests. *J. Pest Sci.* (2004), 92, 281–297. doi: [10.1007/s10340-018-0997-6](https://doi.org/10.1007/s10340-018-0997-6)
- Fantini A, Raffaele F, Torma CZ, Bacer S, Coppola E et al. 2018 Assessment of multiple daily precipitation statistics in ERA-Interim driven Med-CORDEX and EURO-CORDEX experiments against high resolution observations. *Clim. Dyn.* 51, 877–900. doi: [10.1007/s00382-016-3453-4](https://doi.org/10.1007/s00382-016-3453-4)
- FAO 2015 Towards a Regional Collaborative Strategy on Sustainable Water Management and Food Security in the Near East and North Africa Region. Cairo, Egypt.
- FAO 2017a Integrated Monitoring Guide for SDG 6. Step-by-step monitoring methodology for indicator 6.4.1 on water-use efficiency.
- FAO 2017b International standards for phytosanitary measures 5. Glossary of phytosanitary terms.
- Faurès J-M, Hoogeveen J, Bruinsma J 2002 The FAO Irrigated Area Forecast for 2030. <http://www.fao.org/tempref/agl/AGLW/docs/fauresetalagadir.pdf>
- Feidas H, Makrigiannis T, Bora-Senta E 2004 Trend analysis of air temperature time series in Greece and their relationship with circulation using surface and satellite data: 1955–2001. *Theor. Appl. Climatol.* 79, 185–208. doi: [10.1007/s00704-004-0064-5](https://doi.org/10.1007/s00704-004-0064-5)
- Feki W, Hamza A, Frossard V, Abdennadher M, Hannachi I et al. 2013 What are the potential drivers of blooms of the toxic dinoflagellate *Karenia selliformis*? A 10-year study in the Gulf of Gabes, Tunisia, southwestern Mediterranean Sea. *Harmful Algae* 23, 8–18. doi: [10.1016/j.hal.2012.12.001](https://doi.org/10.1016/j.hal.2012.12.001)
- Feliks Y, Ghil M, Robertson AW 2010 Oscillatory Climate Modes in the Eastern Mediterranean and Their Synchronization with the North Atlantic Oscillation. *J. Clim.* 23, 4060–4079. doi: [10.1175/2010jcli3181.1](https://doi.org/10.1175/2010jcli3181.1)
- Felis T, Rimbu N 2010 Mediterranean climate variability documented in oxygen isotope records from northern Red Sea corals—A review. *Glob. Planet. Change* 71, 232–241. doi: [10.1016/j.glopacha.2009.10.006](https://doi.org/10.1016/j.glopacha.2009.10.006)
- Fenoglio-Marc L, Mariotti A, Sannino G, Meyssignac B, Carillo A et al. 2013 Decadal variability of net water flux at the Med-

- iterranean Sea Gibraltar Strait. *Glob. Planet. Change* 100, 1–10. doi: [10.1016/J.GLOPLACHA.2012.08.007](https://doi.org/10.1016/J.GLOPLACHA.2012.08.007)
- Fernandes PG, Ralph GM, Nieto A, Criado MG, Vasilakopoulos P et al. 2017 Coherent assessments of Europe's marine fishes show regional divergence and megafauna loss. *Nat. Ecol. Evol.* 1, 0170. doi: [10.1038/s41559-017-0170](https://doi.org/10.1038/s41559-017-0170)
- Fernández J, Frías MD, Cabos WD, Cofiño AS, Domínguez M et al. 2019 Consistency of climate change projections from multiple global and regional model intercomparison projects. *Clim. Dyn.* 52, 1139–1156. doi: [10.1007/s00382-018-4181-8](https://doi.org/10.1007/s00382-018-4181-8)
- Filahi S, Tanarhte M, Mouhir L, El Morhit M, Tramblay Y 2015 Trends in indices of daily temperature and precipitations extremes in Morocco. *Theor. Appl. Climatol.* 124, 959–972. doi: [10.1007/s00704-015-1472-4](https://doi.org/10.1007/s00704-015-1472-4)
- Fischer-Brunns I, Banse DF, Feichter J 2009 Future impact of anthropogenic sulfate aerosol on North Atlantic climate. *Clim. Dyn.* 32, 511–524. doi: [10.1007/s00382-008-0458-7](https://doi.org/10.1007/s00382-008-0458-7)
- Fischer EM, Schär C 2010 Consistent geographical patterns of changes in high-impact European heatwaves. *Nat. Geosci.* 3, 398–403. doi: [10.1038/ngeo866](https://doi.org/10.1038/ngeo866)
- Flaounas E, Drobinski P, Borga M, Calvet J-C, Delrieu G et al. 2012 Assessment of gridded observations used for climate model validation in the Mediterranean region: the HyMeX and MED-CORDEX framework. *Environ. Res. Lett.* 7, 024017. doi: [10.1088/1748-9326/7/2/024017](https://doi.org/10.1088/1748-9326/7/2/024017)
- Flocas HA, Simmonds I, Kouroutzoglou J, Keay K, Hatzaki M et al. 2010 On Cyclonic Tracks over the Eastern Mediterranean. *J. Clim.* 23, 5243–5257. doi: [10.1175/2010jcli3426.1](https://doi.org/10.1175/2010jcli3426.1)
- Floris I, Cocco A, Buffa F, Mannu R, Satta A 2018 Insect pests of *Eucalyptus* plantations in Sardinia (Italy). *Redia*, 61–71. doi: [10.19263/redia-101.18.09](https://doi.org/10.19263/redia-101.18.09)
- Floutsi AA, Korras-Carraca MB, Matsoukas C, Hatzianastassiou N, Biskos G 2016 Climatology and trends of aerosol optical depth over the Mediterranean basin during the last 12 years (2002–2014) based on Collection 006 MODIS-Aqua data. *Sci. Total Environ.* 551–552, 292–303. doi: [10.1016/J.SCITOTENV.2016.01.192](https://doi.org/10.1016/J.SCITOTENV.2016.01.192)
- Foley JA, DeFries RS, Asner GP, Barford C, Bonan G et al. 2005 Global consequences of land use. *Science (80-.).* 309, 570–574. doi: [10.1126/science.1111772](https://doi.org/10.1126/science.1111772)
- Folini D, Wild M 2011 Aerosol emissions and dimming/brightening in Europe: Sensitivity studies with ECHAM5-HAM. *JGR Atmos.* 116. doi: [10.1029/2011JD016227](https://doi.org/10.1029/2011JD016227)
- Folt CL, Chen CY, Moore M V., Burnaford J 1999 Synergism and antagonism among multiple stressors. *Limnol. Oceanogr.* 44, 864–877. doi: [10.4319/lo.1999.44.3_part_2.0864](https://doi.org/10.4319/lo.1999.44.3_part_2.0864)
- Fosser G, Khodayar S, Berg P 2015 Benefit of convection permitting climate model simulations in the representation of convective precipitation. *Clim. Dyn.* 44, 45–60. doi: [10.1007/s00382-014-2242-1](https://doi.org/10.1007/s00382-014-2242-1)
- Fourati R, Tedetti M, Guigue C, Goutx M, García N et al. 2018a Sources and spatial distribution of dissolved aliphatic and polycyclic aromatic hydrocarbons in surface coastal waters of the Gulf of Gabès (Tunisia, Southern Mediterranean Sea). *Prog. Oceanogr.* 163, 232–247. doi: [10.1016/j.pocean.2017.02.001](https://doi.org/10.1016/j.pocean.2017.02.001)
- Fourati R, Tedetti M, Guigue C, Goutx M, Zaghdén H et al. 2018b Natural and anthropogenic particulate-bound aliphatic and polycyclic aromatic hydrocarbons in surface waters of the Gulf of Gabès (Tunisia, southern Mediterranean Sea). *Environ. Sci. Pollut. Res.* 25, 2476–2494. doi: [10.1007/s11356-017-0641-7](https://doi.org/10.1007/s11356-017-0641-7)
- Francardi V, Noal A, Francescato S, Pinto R, Bruni A et al. 2017 Coexistence of *Xylosandrus crassiusculus* (Motschulsky) and *X. compactus* (Eichhoff) (Coleoptera Curculionidae Scolytinae) in the National Park of Circeo (Lazio, Italy). *Redia* 100, 149–155.
- Francour P, Boudouresque C-F, Harmelin JG, Harmelin-Vivien ML, Quignard JP 1994 Are the Mediterranean waters becoming warmer? Information from biological indicators. *Mar. Pollut. Bull.* 28, 523–526. doi: [10.1016/0025-326x\(94\)90071-x](https://doi.org/10.1016/0025-326x(94)90071-x)
- Fréjaville T, Curt T 2017 Seasonal changes in the human alteration of fire regimes beyond the climate forcing. *Environ. Res. Lett.* 12, 035006. doi: [10.1088/1748-9326/aa5d23](https://doi.org/10.1088/1748-9326/aa5d23)
- Froese R, Kesner-Reyes K 2002 Impact of Fishing on the Abundance of Marine Species. 12.
- Froese R, Winker H, Coro G, Demirel N, Tsikliras AC et al. 2018 Status and rebuilding of European fisheries. *Mar. Policy* 93, 159–170. doi: [10.1016/j.marpol.2018.04.018](https://doi.org/10.1016/j.marpol.2018.04.018)
- Frölicher TL, Fischer EM, Gruber N 2018 Marine heatwaves under global warming. *Nature* 560, 360–364. doi: [10.1038/s41586-018-0383-9](https://doi.org/10.1038/s41586-018-0383-9)
- Fumière Q, Déqué M, Nuissier O, Somot S, Alias A et al. 2019 Extreme rainfall in Mediterranean France during the fall: added value of the CNRM-AROME Convection-Permitting Regional Climate Model. *Clim. Dyn.*, 1–15. doi: [10.1007/s00382-019-04898-8](https://doi.org/10.1007/s00382-019-04898-8)
- Fyfe JC, Boer GJ, Flato GM 1999 The Arctic and Antarctic Oscillations and their projected changes under global warming. *Geophys. Res. Lett.* 26, 1601–1604. doi: [10.1029/1999GL900317](https://doi.org/10.1029/1999GL900317)
- Gačić M, Borzelli GLE, Civitarese G, Cardin V, Yari S 2010 Can internal processes sustain reversals of the ocean upper circulation? The Ionian Sea example. *Geophys. Res. Lett.* 37, n/a-n/a. doi: [10.1029/2010GL043216](https://doi.org/10.1029/2010GL043216)
- Gaertner MÁ, González-Alemán JJ, Romera R, Domínguez M, Gil V et al. 2018 Simulation of medicanes over the Mediterranean Sea in a regional climate model ensemble: impact of ocean-atmosphere coupling and increased resolution. *Clim. Dyn.* 51, 1041–1057. doi: [10.1007/s00382-016-3456-1](https://doi.org/10.1007/s00382-016-3456-1)
- Gaertner MÁ, Jacob D, Gil V, Domínguez M, Padorno E et al. 2007 Tropical cyclones over the Mediterranean Sea in climate change simulations. *Geophys. Res. Lett.* 34, L14711. doi: [10.1029/2007gl029977](https://doi.org/10.1029/2007gl029977)
- Galdies C 2012 Temperature trends in Malta (central Mediterranean) from 1951 to 2010. *Meteorol. Atmos. Phys.* 117, 135–143. doi: [10.1007/s00703-012-0187-7](https://doi.org/10.1007/s00703-012-0187-7)
- Galil BS 2007 Loss or gain? Invasive aliens and biodiversity in the Mediterranean Sea. *Mar. Pollut. Bull.* 55, 314–322. doi: [10.1016/j.marpolbul.2006.11.008](https://doi.org/10.1016/j.marpolbul.2006.11.008)
- Galindo N, Yubero E, Nicolás JF, Varea M, Crespo J 2018 Characterization of metals in PM1 and PM10 and health risk evaluation at an urban site in the western Mediterranean. *Chemosphere* 201, 243–250. doi: [10.1016/j.chemosphere.2018.02.162](https://doi.org/10.1016/j.chemosphere.2018.02.162)

- Gallardo B, Aldridge DC, González-Moreno P, Pergl J, Pizarro M et al. 2017 Protected areas offer refuge from invasive species spreading under climate change. *Glob. Chang. Biol.* 23, 5331–5343. doi: [10.1111/gcb.13798](https://doi.org/10.1111/gcb.13798)
- Gallart F, Delgado JM, Beatson SJ V, Posner H, Llorens P et al. 2011 Analysing the effect of global change on the historical trends of water resources in the headwaters of the Llobregat and Ter river basins [Catalonia, Spain]. *Phys. Chem. Earth, Parts A/B/C* 36, 655–661. doi: [10.1016/j.pce.2011.04.009](https://doi.org/10.1016/j.pce.2011.04.009)
- Galloway TS 2015 Micro- and nano-plastics and human health, in *Marine Anthropogenic Litter* (Cham: Springer International Publishing), 343–366. doi: [10.1007/978-3-319-16510-3_13](https://doi.org/10.1007/978-3-319-16510-3_13)
- Ganor E, Osetinsky I, Stupp A, Alpert P 2010 Increasing trend of African dust, over 49 years, in the eastern Mediterranean. *JGR Atmos.* 115, D07201. doi: [10.1029/2009JD012500](https://doi.org/10.1029/2009JD012500)
- Gao J 2017 Downscaling Global Spatial Population Projections from 1/8-degree to 1-km Grid Cells. doi: [10.5065/d60z721h](https://doi.org/10.5065/d60z721h)
- Gao X, Pal JS, Giorgi F 2006 Projected changes in mean and extreme precipitation over the Mediterranean region from a high resolution double nested RCM simulation. *Geophys. Res. Lett.* 33, 2–5. doi: [10.1029/2005GL024954](https://doi.org/10.1029/2005GL024954)
- Garcés E, Camp J 2012 Habitat changes in the Mediterranean Sea and the consequences for Harmful Algal Blooms formation, in *Life in the Mediterranean Sea: A Look at Habitat Changes*, ed. Stambler N (Nova Science Publishers, Inc.), 519–541.
- García-Ayllón S 2018 Predictive Diagnosis of Agricultural Peri-urban Areas Based on Territorial Indicators: Comparative Landscape Trends of the So-Called “Orchard of Europe.” *Sustainability* 10, 1820. doi: [10.3390/su10061820](https://doi.org/10.3390/su10061820)
- García-Gómez H, Izquierdo-Rojano S, Aguilera L, González-Fernández I, Valiño F et al. 2018 Joining empirical and modelling approaches to estimate dry deposition of nitrogen in Mediterranean forests. *Environ. Pollut.* 243, 427–436. doi: [10.1016/j.envpol.2018.09.015](https://doi.org/10.1016/j.envpol.2018.09.015)
- García-Serrano H, Sans FX, Escarré J 2007 Interspecific competition between alien and native congeneric species. *Acta Oecologica* 31, 69–78. doi: [10.1016/J.ACTAO.2006.09.005](https://doi.org/10.1016/J.ACTAO.2006.09.005)
- Garnas JR, Auger-Rozenberg M-A, Roques A, Bertelsmeier C, Wingfield MJ et al. 2016 Complex patterns of global spread in invasive insects: eco-evolutionary and management consequences. *Biol. Invasions* 18, 935–952. doi: [10.1007/s10530-016-1082-9](https://doi.org/10.1007/s10530-016-1082-9)
- Garrabou J, Coma R, Bensoussan N, Bally M, Chevaldonné P et al. 2009 Mass mortality in Northwestern Mediterranean rocky benthic communities: Effects of the 2003 heat wave. *Glob. Chang. Biol.* 15, 1090–1103. doi: [10.1111/j.1365-2486.2008.01823.x](https://doi.org/10.1111/j.1365-2486.2008.01823.x)
- Garrido A, Martínez-Santos P, Llamas MR 2006 Groundwater irrigation and its implications for water policy in semiarid countries: the Spanish experience. *Hydrogeol. J.* 14, 340–349. doi: [10.1007/s10040-005-0006-z](https://doi.org/10.1007/s10040-005-0006-z)
- Garrido M, Cecchi P, Collos Y, Agostini S, Pasqualini V 2016 Water flux management and phytoplankton communities in a Mediterranean coastal lagoon. Part I: How to promote dinoflagellate dominance? *Mar. Pollut. Bull.* 104, 139–152. doi: [10.1016/j.marpolbul.2016.01.049](https://doi.org/10.1016/j.marpolbul.2016.01.049)
- Gasperi J, Wright SL, Dris R, Collard F, Mandin C et al. 2018 Microplastics in air: Are we breathing it in? *Curr. Opin. Environ. Sci. Heal.* 1, 1–5. doi: [10.1016/J.COESH.2017.10.002](https://doi.org/10.1016/J.COESH.2017.10.002)
- Gassó N, Pino J, Font X, Vilà M 2012 Regional context affects native and alien plant species richness across habitat types. *Appl. Veg. Sci.* 15, 4–13. doi: [10.1111/j.1654-109x.2011.01159.x](https://doi.org/10.1111/j.1654-109x.2011.01159.x)
- Gattuso J-P, Magnan A, Billé R, Cheung WWL, Howes EL et al. 2015 Contrasting futures for ocean and society from different anthropogenic CO₂ emissions scenarios. *Science (80-).* 349. doi: [10.1126/science.aac4722](https://doi.org/10.1126/science.aac4722)
- Gea-Izquierdo G, Nicault A, Battipaglia G, Dorado Liñán I, Gutiérrez E et al. 2017 Risky future for Mediterranean forests unless they undergo extreme carbon fertilization. *Glob. Chang. Biol.*, 1–13. doi: [10.1111/gcb.13597](https://doi.org/10.1111/gcb.13597)
- Geng Q, Sugi M 2003 Possible Change of Extratropical Cyclone Activity due to Enhanced Greenhouse Gases and Sulfate Aerosols—Study with a High-Resolution AGCM. *J. Clim.* 16, 2262–2274. doi: [10.1175/1520-0442\[2003\]16<2262:pcoeca>2.0.co;2](https://doi.org/10.1175/1520-0442[2003]16<2262:pcoeca>2.0.co;2)
- Genovesi P, Bacher S, Kobelt M, Pascal M, Scalera R 2009 Alien mammals of Europe, in *Handbook of alien species in Europe*, ed. DAISIE (Springer Netherlands), 119–128. doi: [10.1007/978-1-4020-8280-1_9](https://doi.org/10.1007/978-1-4020-8280-1_9)
- Gerard F, Petit S, Smith G, Thomson A, Brown N et al. 2010 Land cover change in Europe between 1950 and 2000 determined employing aerial photography. *Prog. Phys. Geogr. Earth Environ.* 34, 183–205. doi: [10.1177/0309133309360141](https://doi.org/10.1177/0309133309360141)
- Gerasopoulos E, Kouvarakis G, Babasaki P, Vrekoussis M, Putaud JP et al. 2006 Origin and variability of particulate matter (PM10) mass concentrations over the Eastern Mediterranean. *Atmos. Environ.* 40, 4679–4690. doi: [10.1016/j.atmosenv.2006.04.020](https://doi.org/10.1016/j.atmosenv.2006.04.020)
- Geyer R, Jambeck JR, Law KL 2017 Production, use, and fate of all plastics ever made. *Sci. Adv.* 3, e1700782. doi: [10.1126/sciadv.1700782](https://doi.org/10.1126/sciadv.1700782)
- Ghelardini L, Luchi N, Pecori F, Pepori AL, Danti R et al. 2017 Ecology of invasive forest pathogens. *Biol. Invasions* 19, 3183–3200. doi: [10.1007/s10530-017-1487-0](https://doi.org/10.1007/s10530-017-1487-0)
- Giakoumi S, Pey A, Huseyinoglu MF 2019 Assessing the state of invasive fishes in two Mediterranean marine protected areas and adjacent unprotected areas. in *1st Mediterranean Symposium on Non-Indigenous Species, Antalia, Turkey* [Frontiers Media SA], 53–58. doi: [10.3389/fmars.2017.00049](https://doi.org/10.3389/fmars.2017.00049)
- Giannakopoulou E-M, Toumi R 2012 Impacts of the Nile Delta land-use on the local climate. *Atmos. Sci. Lett.* 13, 208–215. doi: [10.1002/asl.381](https://doi.org/10.1002/asl.381)
- Gil-Tena A, Brotons L, Saura S 2010 Effects of forest landscape change and management on the range expansion of forest bird species in the Mediterranean region. *For. Ecol. Manage.* 259, 1338–1346. doi: [10.1016/j.foreco.2009.10.026](https://doi.org/10.1016/j.foreco.2009.10.026)
- Gillet NP, Fyfe JC, Parker DE 2013 Attribution of observed sea level pressure trends to greenhouse gas, aerosol, and ozone changes. *Geophys. Res. Lett.* 40, 2302–2306. doi: [10.1002/grl.50500](https://doi.org/10.1002/grl.50500)
- Gillet NP, Graf HF, Osborn TJ 2003 Climate change and the North Atlantic oscillation. *Geophys. Monogr. Ser.* 134, 193–209. doi: [10.1029/134GM09](https://doi.org/10.1029/134GM09)

- Giménez Papiol G, Casanova A, Fernández-Tejedor M, de la Iglesia P, Diogène J 2013 Management of domoic acid monitoring in shellfish from the Catalan coast. *Environ. Monit. Assess.* 185, 6653–6666. doi: [10.1007/s10661-012-3054-6](https://doi.org/10.1007/s10661-012-3054-6)
- Gimeno I, Vilà M, Hulme PE 2006 Are islands more susceptible to plant invasion than continents? A test using *Oxalis pes-caprae* L. in the western Mediterranean. *J. Biogeogr.* 33, 1559–1565. doi: [10.1111/j.1365-2699.2006.01525.x](https://doi.org/10.1111/j.1365-2699.2006.01525.x)
- Giorgi F 2006 Climate change hotspots. *Geophys. Res. Lett.* 33, L08707. doi: [10.1029/2006GL025734](https://doi.org/10.1029/2006GL025734)
- Giorgi F, Bi X 2005 Updated regional precipitation and temperature changes for the 21st century from ensembles of recent AOGCM simulations. *Geophys. Res. Lett.* 32, L21715. doi: [10.1029/2005GL024288](https://doi.org/10.1029/2005GL024288)
- Giorgi F, Coppola E 2007 European climate-change oscillation (ECO). *Geophys. Res. Lett.* 34, L21703. doi: [10.1029/2007gl031223](https://doi.org/10.1029/2007gl031223)
- Giorgi F, Coppola E 2009 Projections of twenty-first century climate over Europe. *EPJ Web Conf.* 1, 29–46. doi: [10.1140/epjconf/e2009-00908-9](https://doi.org/10.1140/epjconf/e2009-00908-9)
- Giorgi F, Coppola E, Raffaele F 2014 A consistent picture of the hydroclimatic response to global warming from multiple indices: Models and observations. *J. Geophys. Res. Atmos.* 119, 11,611–695,708. doi: [10.1002/2014JD022238](https://doi.org/10.1002/2014JD022238)
- Giorgi F, Im E-S, Coppola E, Daffenbaugh NS, Gao XJ et al. 2011 Higher Hydroclimatic Intensity with Global Warming. *J. Clim.* 24, 5309–5324. doi: [10.1175/2011JCLI3979.1](https://doi.org/10.1175/2011JCLI3979.1)
- Giorgi F, Jones C, Asrar GR 2009 Addressing climate information needs at the regional level: The CORDEX framework. *WMO Bull.* 58, 175.
- Giorgi F, Lionello P 2008 Climate change projections for the Mediterranean region. *Glob. Planet. Change* 63, 90–104. doi: [10.1016/j.gloplacha.2007.09.005](https://doi.org/10.1016/j.gloplacha.2007.09.005)
- Giorgi F, Raffaele F, Coppola E 2019 The response of precipitation characteristics to global warming from climate projections. *Earth Syst. Dynam.* 10, 73–89. doi: [10.5194/esd-10-73-2019](https://doi.org/10.5194/esd-10-73-2019)
- Giorgi F, Torma CZ, Coppola E, Ban N, Schär C et al. 2016 Enhanced summer convective rainfall at Alpine high elevations in response to climate warming. *Nat. Geosci.* 9, 584–589. doi: [10.1038/ngeo2761](https://doi.org/10.1038/ngeo2761)
- Gitay H, Brown S, Easterling W, Jallow B, Antle J et al. 2001 Ecosystems and their goods and services, in *Climate Change 2001: Impacts, Adaptation, and Vulnerability*, eds. McCarthy JJ, Canziani OF, Leary NA, Dokken DJ, White KS (Cambridge, UK and New York, NY, USA), 235–342.
- Givan O, Edelist D, Sonin O, Belmaker J 2017a Thermal affinity as the dominant factor changing Mediterranean fish abundances. *Glob. Chang. Biol.* 24, e80–e89. doi: [10.1111/gcb.13835](https://doi.org/10.1111/gcb.13835)
- Givan O, Parravicini V, Kulbicki M, Belmaker J 2017b Trait structure reveals the processes underlying fish establishment in the Mediterranean. *Glob. Ecol. Biogeogr.* 26, 142–153. doi: [10.1111/geb.12523](https://doi.org/10.1111/geb.12523)
- Godinho S, Gil A, Guiomar N, Costa MJ, Neves N 2016 Assessing the role of Mediterranean evergreen oaks canopy cover in land surface albedo and temperature using a remote sensing-based approach. *Appl. Geogr.* 74, 84–94. doi: [10.1016/j.apgeog.2016.07.004](https://doi.org/10.1016/j.apgeog.2016.07.004)
- Gogou A, Bouloubassi I, Stephanou EG 2000 Marine organic geochemistry of the Eastern Mediterranean: 1. Aliphatic and polyaromatic hydrocarbons in Cretan Sea surficial sediments. *Mar. Chem.* 68, 265–282. doi: [10.1016/S0304-4203\(99\)00082-1](https://doi.org/10.1016/S0304-4203(99)00082-1)
- Gómez-Gutiérrez AI, Jover E, Bodineau L, Albaiges J, Bayona JM 2006 Organic contaminant loads into the Western Mediterranean Sea: Estimate of Ebro River inputs. *Chemosphere* 65, 224–236. doi: [10.1016/j.chemosphere.2006.02.058](https://doi.org/10.1016/j.chemosphere.2006.02.058)
- Gómez F, González N, Echevarría F, García CM 2000 Distribution and fluxes of dissolved nutrients in the Strait of Gibraltar and its relationships to microphytoplankton biomass. *Estuar. Coast. Shelf Sci.* 51, 439–449. doi: [10.1006/ecss.2000.0689](https://doi.org/10.1006/ecss.2000.0689)
- González-Alemán JJ, Pascale S, Gutierrez-Fernandez J, Murakami H, Gaertner MÁ et al. 2019 Potential Increase in Hazard From Mediterranean Hurricane Activity With Global Warming. *Geophys. Res. Lett.* 46, 1754–1764. doi: [10.1029/2018GL081253](https://doi.org/10.1029/2018GL081253)
- González-Moreno P, Lazzaro L, Vilà M, Preda C, Adriaens T et al. 2019 Consistency of impact assessment protocols for non-native species. *NeoBiota* 44, 1–25. doi: [10.3897/neobiota.44.31650](https://doi.org/10.3897/neobiota.44.31650)
- González-Moreno P, Pino J, Gassó N, Vilà M 2013 Landscape context modulates alien plant invasion in Mediterranean forest edges. *Biol. Invasions* 15, 547–557. doi: [10.1007/s10530-012-0306-x](https://doi.org/10.1007/s10530-012-0306-x)
- Gormley-Gallagher AM, Sterl S, Hirsch AL, Seneviratne SI, Davin EL et al. 2020 Agricultural management effects on mean and extreme temperature trends. *Earth Syst. Dynam. Discuss.*, 1–27. doi: [10.5194/esd-2020-35](https://doi.org/10.5194/esd-2020-35)
- Gottipolu RR, Landa ER, Schladweiler MC, McGee JK, Ledbetter AD et al. 2008 Cardiopulmonary Responses of Intratracheally Instilled Tire Particles and Constituent Metal Components. *Inhal. Toxicol.* 20, 473–484. doi: [10.1080/08958370701858427](https://doi.org/10.1080/08958370701858427)
- Goubanova K, Li L 2007 Extremes in temperature and precipitation around the Mediterranean basin in an ensemble of future climate scenario simulations. *Glob. Planet. Change* 57, 27–42. doi: [10.1016/j.gloplacha.2006.11.012](https://doi.org/10.1016/j.gloplacha.2006.11.012)
- Goubert C, Minard G, Vieira C, Boulesteix M 2016 Population genetics of the Asian tiger mosquito *Aedes albopictus*, an invasive vector of human diseases. *Heredity (Edinb.)* 117, 125–134. doi: [10.1038/hdy.2016.35](https://doi.org/10.1038/hdy.2016.35)
- Goyet C, Hassoun AER, Gemayel E, Touratier F, Abboud-Abi Saab M et al. 2016 Thermodynamic forecasts of the Mediterranean Sea acidification. *Mediterr. Mar. Sci.* 17, 508–518. doi: [10.12681/mms.1487](https://doi.org/10.12681/mms.1487)
- Gramberger M, Harrison PA, Jäger J, Kok K, Libbrecht S et al. 2013 Report on the third CLIMSAVE European stakeholder workshop. CLIMSAVE Project.
- Grimm NB, Faeth SH, Golubiewski NE, Redman CL, Wu J et al. 2008 Global change and the ecology of cities. *Science (80-.).* 319, 756–760. doi: [10.1126/science.1150195](https://doi.org/10.1126/science.1150195)
- Grise KM, Davis SM, Simpson IR, Waugh DW, Fu Q et al. 2019

- Recent tropical expansion: natural variability or forced response? *J. Clim.* 32, 1551–1571.
doi: [10.1175/jcli-d-18-0444.1](https://doi.org/10.1175/jcli-d-18-0444.1)
- Grisogono B, Belušić D 2009 A review of recent advances in understanding the meso- and microscale properties of the severe Bora wind. *Tellus A Dyn. Meteorol. Oceanogr.* 61 A, 1–16. doi: [10.1111/j.1600-0870.2008.00369.x](https://doi.org/10.1111/j.1600-0870.2008.00369.x)
- Gritti ES, Smith B, Sykes MT 2006 Vulnerability of Mediterranean Basin ecosystems to climate change and invasion by exotic plant species. *J. Biogeogr.* 33, 145–157.
doi: [10.1111/j.1365-2699.2005.01377.x](https://doi.org/10.1111/j.1365-2699.2005.01377.x)
- Gros M, Petrović M, Ginebreda A, Barceló D 2010 Removal of pharmaceuticals during wastewater treatment and environmental risk assessment using hazard indexes. *Environ. Int.* 36, 15–26. doi: [10.1016/j.envint.2009.09.002](https://doi.org/10.1016/j.envint.2009.09.002)
- Gualdi S, Somot S, Li L, Artale V, Adani M et al. 2013 The CIRCE Simulations: Regional Climate Change Projections with Realistic Representation of the Mediterranean Sea. *Bull. Am. Meteorol. Soc.* 94. doi: [10.1175/BAMS-D-11-00136.1](https://doi.org/10.1175/BAMS-D-11-00136.1)
- Guichard L, Dedieu F, Jeuffroy M-H, Meynard J-M, Reau R et al. 2017 Le plan Ecophyto de réduction d'usage des pesticides en France : décryptage d'un échec et raisons d'espérer. *Cah. Agric.* 26, 14002. doi: [10.1051/cagri/2017004](https://doi.org/10.1051/cagri/2017004)
- Guigue C, Tedetti M, Dang DH, Mullot J-U, Garnier C et al. 2017 Remobilization of polycyclic aromatic hydrocarbons and organic matter in seawater during sediment resuspension experiments from a polluted coastal environment: Insights from Toulon Bay (France). *Environ. Pollut.* 229, 627–638.
doi: [10.1016/j.envpol.2017.06.090](https://doi.org/10.1016/j.envpol.2017.06.090)
- Guigue C, Tedetti M, Ferretto N, García N, Méjanelle L et al. 2014 Spatial and seasonal variabilities of dissolved hydrocarbons in surface waters from the Northwestern Mediterranean Sea: Results from one year intensive sampling. *Sci. Total Environ.* 466–467, 650–662.
doi: [10.1016/j.scitotenv.2013.07.082](https://doi.org/10.1016/j.scitotenv.2013.07.082)
- Guigue C, Tedetti M, Giorgi S, Goutx M 2011 Occurrence and distribution of hydrocarbons in the surface microlayer and subsurface water from the urban coastal marine area off Marseilles, Northwestern Mediterranean Sea. *Mar. Pollut. Bull.* 62, 2741–2752. doi: [10.1016/j.marpolbul.2011.09.013](https://doi.org/10.1016/j.marpolbul.2011.09.013)
- Gutiérrez C, Somot S, Nabat P, Mallet M, Corre L et al. 2020 Future evolution of surface solar radiation and photovoltaic potential in Europe: investigating the role of aerosols. *Environ. Res. Lett.* 15, 034035. doi: [10.1088/1748-9326/ab6666](https://doi.org/10.1088/1748-9326/ab6666)
- Guy-Haim T 2017 The impact of ocean warming and acidification on coastal benthic species and communities.
- Guy-Haim T, Silverman J, Raddatz S, Wahl M, Rilov G 2016 Shifted coastal communities and ecosystem functions under predicted warming and acidification. In *41st CIESM Congress, Kiel, Germany*.
- Guzmán GI, González de Molina M, Soto Fernández D, Infante-Amate J, Aguilera E 2018 Spanish agriculture from 1900 to 2008: a long-term perspective on agroecosystem energy from an agroecological approach. *Reg. Environ. Chang.* 18, 995–1008. doi: [10.1007/s10113-017-1136-2](https://doi.org/10.1007/s10113-017-1136-2)
- Haarsma RJ, Selten F, Van den Hurk B, Hazeleger W, Wang X 2009 Drier Mediterranean soils due to greenhouse warm-
ing bring easterly winds over summertime central Europe. *Geophys. Res. Lett.* 36. doi: [10.1029/2008gl036617](https://doi.org/10.1029/2008gl036617)
- Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M et al. 2008 Seawater carbonate chemistry in Ischia, Italy, 2008. doi: [10.1594/pangaea.819633](https://doi.org/10.1594/pangaea.819633)
- Hallegraeff GM 2010 Ocean climate change, phytoplankton community responses, and harmful algal blooms: A formidable predictive challenge. *J. Phycol.* 46, 220–235.
doi: [10.1111/j.1529-8817.2010.00815.x](https://doi.org/10.1111/j.1529-8817.2010.00815.x)
- Halpern BS, McLeod KL, Rosenberg AA, Crowder LB 2008a Managing for cumulative impacts in ecosystem-based management through ocean zoning. *Ocean Coast. Manag.* 51, 203–211. doi: [10.1016/j.ocecoaman.2007.08.002](https://doi.org/10.1016/j.ocecoaman.2007.08.002)
- Halpern BS, Walbridge S, Selkoe KA, Kappel C V., Micheli F et al. 2008b A global map of human impact on marine ecosystems. *Science (80-)* 319, 948–952.
doi: [10.1126/science.1149345](https://doi.org/10.1126/science.1149345)
- Handmer J, Honda Y, Kundzewicz ZW, Arnell NW, Benito G et al. 2012 Changes in impacts of climate extremes: human systems and ecosystems, in *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation* A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC), eds. Field CB, Barros V, Stocker TF, Qin D, Dokken DJ et al. (Cambridge, United Kingdom and New York, NY, USA), 231–290.
- Hannah L, Roehrdanz PR, Ikegami M, Shepard A V., Shaw MR et al. 2013 Climate change, wine, and conservation. *Proc. Natl. Acad. Sci. U. S. A.* 110, 6907–6912.
doi: [10.1073/pnas.1210127110](https://doi.org/10.1073/pnas.1210127110)
- Hansen MC, DeFries RS 2004 Detecting Long-term Global Forest Change Using Continuous Fields of Tree-Cover Maps from 8-km Advanced Very High Resolution Radiometer [AVHRR] Data for the Years 1982?99. *Ecosystems* 7, 695–716. doi: [10.1007/s10021-004-0243-3](https://doi.org/10.1007/s10021-004-0243-3)
- Hanson S, Nicholls RJ 2012 Extreme flood events and port cities through the twenty-first century: implications of climate change and other drivers, in *Maritime Transport and the Climate Change Challenge*, eds. Asariotis R, Benamara H (Abingdon, GB: Earthscan from Routledge), 243–265.
- Hanzer F, Förster K, Nemec J, Strasser U 2018 Projected cryospheric and hydrological impacts of 21st century climate change in the Ötztal Alps (Austria) simulated using a physically based approach. *Hydrol. Earth Syst. Sci.* 22, 1593–1614.
doi: [10.5194/hess-22-1593-2018](https://doi.org/10.5194/hess-22-1593-2018)
- Hao R, Yu D, Liu Y, Liu Y, Qiao J et al. 2017 Impacts of changes in climate and landscape pattern on ecosystem services. *Sci. Total Environ.* 579, 718–728.
doi: [10.1016/j.scitotenv.2016.11.036](https://doi.org/10.1016/j.scitotenv.2016.11.036)
- Harris I, Osborn TJ, Jones PD, Lister D 2020 Version 4 of the CRU TS monthly high-resolution gridded multivariate climate dataset. *Sci. Data* 7, 109. doi: [10.1038/s41597-020-0453-3](https://doi.org/10.1038/s41597-020-0453-3)
- Harvey BJ, Shaffrey LC, Woollings TJ, Zappa G, Hodges KI 2012 How large are projected 21st century storm track changes. *Geophys. Res. Lett.* 39. doi: [10.1029/2012GL052873](https://doi.org/10.1029/2012GL052873)
- Harzallah A, Jordà G, Dubois C, Sannino G, Carillo A et al. 2018 Long term evolution of heat budget in the Mediterranean Sea from Med-CORDEX forced and coupled simulations.

- Clim. Dyn.* 51, 1145–1165. doi: [10.1007/s00382-016-3363-5](https://doi.org/10.1007/s00382-016-3363-5)
- Hasanean HM 2004 Precipitation variability over the Mediterranean and its linkage with El Niño Southern Oscillation (ENSO). *J. Meteorol.* 29, 151–160.
- Hassanien MA, Abdel-Latif NM 2008 Polycyclic aromatic hydrocarbons in road dust over Greater Cairo, Egypt. *J. Hazard. Mater.* 151, 247–254. doi: [10.1016/j.jhazmat.2007.05.079](https://doi.org/10.1016/j.jhazmat.2007.05.079)
- Hatfield JL, Boote KJ, Kimball BA, Ziska LH, Izaurralde RC et al. 2011 Climate Impacts on Agriculture: Implications for Crop Production. *Agron. J.* 103, 351. doi: [10.2134/agronj2010.0303](https://doi.org/10.2134/agronj2010.0303)
- Hatzaki M, Flocas HA, Giannakopoulos C, Maher P 2009 The impact of the eastern Mediterranean teleconnection pattern on the Mediterranean climate. *J. Clim.* 22, 977–992. doi: [10.1175/2008JCLI2519.1](https://doi.org/10.1175/2008JCLI2519.1)
- Hatzaki M, Flocas HA, Simmonds I, Kouroutzoglou J, Keay K et al. 2014 Seasonal Aspects of an Objective Climatology of Anticyclones Affecting the Mediterranean. *J. Clim.* 27, 9272–9289. doi: [10.1175/JCLI-D-14-00186.1](https://doi.org/10.1175/JCLI-D-14-00186.1)
- Haywood JM, Bellouin N, Jones A, Boucher O, Wild M et al. 2011 The roles of aerosol, water vapor and cloud in future global dimming/brightening. *JGR Atmos.* 116, D20203. doi: [10.1029/2011JD016000](https://doi.org/10.1029/2011JD016000)
- Hellmann JJ, Byers JE, Bierwagen BG, Dukes JS 2008 Five Potential Consequences of Climate Change for Invasive Species. *Conserv. Biol.* 22, 534–543. doi: [10.1111/j.1523-1739.2008.00951.x](https://doi.org/10.1111/j.1523-1739.2008.00951.x)
- Hentgen L, Ban N, Kröner N, Leutwyler D, Schär C 2019 Clouds in Convection-Resolving Climate Simulations Over Europe. *JGR Atmos.* 124, 3849–3870. doi: [10.1029/2018JD030150](https://doi.org/10.1029/2018JD030150)
- Hernández-Brito D, Carrete M, Ibáñez C, Juste J, Tella J 2018 Nest-site competition and killing by invasive parakeets cause the decline of a threatened bat population. *R. Soc. Open Sci.* 5. doi: [10.1098/rsos.172477](https://doi.org/10.1098/rsos.172477)
- Hernández-Brito D, Carrete M, Popa-Lisseanu AG, Ibáñez C, Tella JL 2014 Crowding in the City: Losing and Winning Competitors of an Invasive Bird. *PLoS One* 9, e100593. doi: [10.1371/journal.pone.0100593](https://doi.org/10.1371/journal.pone.0100593)
- Hernandez-Vargas G, Sosa-Hernández JE, Saldarriaga-Hernández S, Villalba-Rodríguez AM, Parra-Saldívar R et al. 2018 Electrochemical biosensors: A solution to pollution detection with reference to environmental contaminants. *Biosensors* 8, 1–21. doi: [10.3390/bios8020029](https://doi.org/10.3390/bios8020029)
- Hernández A, Miranda M, Arellano EC, Saura S, Ovalle C 2015 Landscape dynamics and their effect on the functional connectivity of a Mediterranean landscape in Chile. *Ecol. Indic.* 48, 198–206. doi: [10.1016/j.ecolind.2014.08.010](https://doi.org/10.1016/j.ecolind.2014.08.010)
- Hernández F, Alonso-Pérez S, Hernández-Armas J, Cuevas E, Karlsson L et al. 2005 Influence of major African dust intrusions on the 137Cs and 40K activities in the lower atmosphere at the Island of Tenerife. *Atmos. Environ.* 39, 4111–4118. doi: [10.1016/J.ATMOSENV.2005.03.032](https://doi.org/10.1016/J.ATMOSENV.2005.03.032)
- Herrero-Hernández E, Andrade MS, Álvarez-Martín A, Pose-Juan E, Rodríguez-Cruz MS et al. 2013 Occurrence of pesticides and some of their degradation products in waters in a Spanish wine region. *J. Hydrol.* 486, 234–245. doi: [10.1016/j.jhydrol.2013.01.025](https://doi.org/10.1016/j.jhydrol.2013.01.025)
- Herrmann M, Somot S, Calmanti S, Dubois C, Sevault F 2011 Representation of spatial and temporal variability of daily wind speed and of intense wind events over the Mediterranean Sea using dynamical downscaling: impact of the regional climate model configuration. *Nat. Hazards Earth Syst. Sci.*, 1983–2011. <https://hal.archives-ouvertes.fr/hal-00766447/> [Accessed September 17, 2019]
- Hertig E, Seubert S, Jacobbeit J 2010 Temperature extremes in the Mediterranean area: Trends in the past and assessments for the future. *Nat. Hazards Earth Syst. Sci.* 10, 2039–2050. doi: [10.5194/nhess-10-2039-2010](https://doi.org/10.5194/nhess-10-2039-2010)
- Hewitson B, Janetos AC, Carter TR, Giorgi F, Jones RG et al. 2014 Regional context, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ et al. (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press), 1133–1197.
- Hildebrandt A, Guillamón M, Lacorte S, Tauler R, Barceló D 2008 Impact of pesticides used in agriculture and vineyards to surface and groundwater quality (North Spain). *Water Res.* 42, 3315–26. doi: [10.1016/j.watres.2008.04.009](https://doi.org/10.1016/j.watres.2008.04.009)
- Hill J, Stellmes M, Udelhoven T, Röder A, Sommer S 2008 Mediterranean desertification and land degradation. *Glob. Planet. Change* 64, 146–157. doi: [10.1016/j.gloplacha.2008.10.005](https://doi.org/10.1016/j.gloplacha.2008.10.005)
- Hirai H, Takada H, Ogata Y, Yamashita R, Mizukawa K et al. 2011 Organic micropollutants in marine plastics debris from the open ocean and remote and urban beaches. *Mar. Pollut. Bull.* 62, 1683–1692. doi: [10.1016/j.marpolbul.2011.06.004](https://doi.org/10.1016/j.marpolbul.2011.06.004)
- Hirche A, Salamani M, Tarhouni M, Nedjaoui D, El Hag M et al. 2018 The Maghreb (North Africa) rangelands evolution over forty years: re-greening or degradation, in *Desertification: Past, Current and Future Trends*, eds. Squires VR, Ariapour A (New York, United States: Nova Science Publishers), 73–106.
- Hirose K, Povinec PP 2015 Sources of plutonium in the atmosphere and stratosphere-troposphere mixing. *Sci. Rep.* 5, 15707. doi: [10.1038/srep15707](https://doi.org/10.1038/srep15707)
- Hoagland P, Anderson DM, Kaoru Y, White AW 2002 The economic effects of harmful algal blooms in the United States: Estimates, assessment issues, and information needs. *Estuaries* 25, 819–837. doi: [10.1007/BF02804908](https://doi.org/10.1007/BF02804908)
- Hobbs RJ, Huenneke LF 1992 Disturbance, Diversity, and Invasion: Implications for Conservation. *Conserv. Biol.* 6, 324–337. doi: [10.1046/j.1523-1739.1992.06030324.x](https://doi.org/10.1046/j.1523-1739.1992.06030324.x)
- Hock R, Bliss A, Marzeion B, Giesen RH, Hirabayashi Y et al. 2019 GlacierMIP – A model intercomparison of global-scale glacier mass-balance models and projections. *J. Glaciol.* 65, 453–467. doi: [10.1017/jog.2019.22](https://doi.org/10.1017/jog.2019.22)
- Hódar JA, Zamora R 2004 Herbivory and climatic warming: a Mediterranean outbreeding caterpillar attacks a relict, boreal pine species. *Biodivers. Conserv.* 13, 493–500. doi: [10.1023/b:bioc.0000009495.95589.a7](https://doi.org/10.1023/b:bioc.0000009495.95589.a7)
- Hoerling M, Eischeid J, Perlitz J, Quan X, Zhang T et al. 2012 On the Increased Frequency of Mediterranean Drought. *J. Clim.* 25, 2146–2161. doi: [10.1175/JCLI-D-11-00296.1](https://doi.org/10.1175/JCLI-D-11-00296.1)

- Horton R, Rosenzweig C, Gornitz V, Bader D, O'Grady M 2010 Climate risk information: climate change scenarios and implications for NYC infrastructure. New York City Panel on Climate Change. *Ann. N. Y. Acad. Sci.* 1196, 147–228. doi: [10.1111/j.1749-6632.2010.05323.x](https://doi.org/10.1111/j.1749-6632.2010.05323.x)
- Hoskins BJ, Hodges KI 2002 New perspectives on the northern hemisphere winter storm tracks. *J. Atmos. Sci.* 59, 1041–1061. doi: [10.1175/1520-0469\(2002\)059<1041:nptn>2.0.co;2](https://doi.org/10.1175/1520-0469(2002)059<1041:nptn>2.0.co;2)
- Hostert P, Röder A, Hill J, Udelhoven T, Tsioroulis G 2003 Retrospective studies of grazing-induced land degradation: A case study in central Crete, Greece. *Int. J. Remote Sens.* 24, 4019–4034. doi: [10.1080/014316031000103844](https://doi.org/10.1080/014316031000103844)
- Houpert L, de Madron XD, Testor P, Bosse A, D'Ortenzio F et al. 2016 Observations of open-ocean deep convection in the northwestern Mediterranean Sea: Seasonal and interannual variability of mixing and deep water masses for the 2007–2013 Period. *JGR Ocean.* 121, 8139–8171. doi: [10.1002/2016JC011857](https://doi.org/10.1002/2016JC011857)
- Hu ZZ, Wu Z 2004 The intensification and shift of the annual North Atlantic Oscillation in a global warming scenario simulation. *Tellus A Dyn. Meteorol. Oceanogr.* 56, 112–124. doi: [10.1111/j.1600-0870.2004.00050.x](https://doi.org/10.1111/j.1600-0870.2004.00050.x)
- Hueging H, Haas R, Born K, Jacob D, Pinto JG 2013 Regional changes in wind energy potential over Europe using regional climate model ensemble projections. *J. Appl. Meteorol. Climatol.* 52, 903–917. doi: [10.1175/JAMC-D-12-086.1](https://doi.org/10.1175/JAMC-D-12-086.1)
- Hughes PD 2018 Little Ice Age glaciers and climate in the Mediterranean mountains: a new analysis. *Geogr. Res. Lett.* 44, 15–45. doi: [10.18172/cig.3362](https://doi.org/10.18172/cig.3362)
- Hulme PE 2009 Trade, transport and trouble: managing invasive species pathways in an era of globalization. *J. Appl. Ecol.* 46, 10–18. doi: [10.1111/j.1365-2664.2008.01600.x](https://doi.org/10.1111/j.1365-2664.2008.01600.x)
- Hulme PE, Brundu G, Camarda I, Dalias P, Lambdon PW et al. 2008 Assessing the risks to Mediterranean islands ecosystems from alien plant introductions, in *Plant invasions: human perception, ecological impacts and management*, eds. Tokarska-Guzik B, Brock JH, Brundu G, Child L, Daehler CC et al. (Buckhuys Publishers, Leiden), 39–56.
- Hunt A, Watkiss P 2011 Climate change impacts and adaptation in cities: a review of the literature. *Clim. Change* 104, 13–49. doi: [10.1007/s10584-010-9975-6](https://doi.org/10.1007/s10584-010-9975-6)
- Hurrell JW 1995 Decadal Trends in the North Atlantic Oscillation: Regional Temperatures and Precipitation. *Science (80-).* 269, 676–679. doi: [10.1126/science.269.5224.676](https://doi.org/10.1126/science.269.5224.676)
- Hurrell JW, Kushnir Y, Ottersen G, Visbeck M 2003 *The North Atlantic Oscillation: Climatic Significance and Environmental Impact*. Washington DC: American Geophysical Union doi: [10.1029/gm134p0vii](https://doi.org/10.1029/gm134p0vii)
- Hurrell JW, Van Loon H 1997 Decadal variations in climate associated with the North Atlantic oscillation. *Clim. Change* 36, 301–326. doi: [10.1023/A:1005314315270](https://doi.org/10.1023/A:1005314315270)
- Hussein RA, Al-Ghanim KA, Abd-El-Atty MM, Mohamed LA 2016 Contamination of red sea shrimp (*Palaemon serratus*) with polycyclic aromatic hydrocarbons: A health risk assessment study. *Polish J. Environ. Stud.* 25, 615–620. doi: [10.1524/pjoes/60767](https://doi.org/10.1524/pjoes/60767)
- Ignatiades L, Gotsis-Skretas O, Pagou K, Krasakopoulou E 2009 Diversification of phytoplankton community structure and related parameters along a large-scale longitudinal east-west transect of the Mediterranean Sea. *J. Plankton Res.* 31, 411–428. doi: [10.1093/plankt/fbn124](https://doi.org/10.1093/plankt/fbn124)
- Iles C, Hegerl GC 2017 Role of the North Atlantic Oscillation in decadal temperature trends. *Environ. Res. Lett.* 12, 114010. doi: [10.1088/1748-9326/aa9152](https://doi.org/10.1088/1748-9326/aa9152)
- Im U 2013 Impact of sea-salt emissions on the model performance and aerosol chemical composition and deposition in the East Mediterranean coastal regions. *Atmos. Environ.* 75, 329–340. doi: [10.1016/J.ATMOSENV.2013.04.034](https://doi.org/10.1016/J.ATMOSENV.2013.04.034)
- Inghilesi AF, Mazza G, Cervo R, Gherardi F, Sposimo P et al. 2013 Alien Insects in Italy: Comparing Patterns from the Regional to European Level. *J. Insect Sci.* 13, 1–13. doi: [10.1673/031.013.7301](https://doi.org/10.1673/031.013.7301)
- IPCC 2007 *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, , eds. Solomon S, Qin D, Manning M, Chen Z, Marquis M et al. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.
- IPCC 2013a *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, , eds. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK et al. Cambridge, United Kingdom and New York, NY, USA.
- IPCC 2013b Summary for Policymakers, in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, eds. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK et al. [Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press].
- IPCC 2018 *Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change*, , eds. Masson-Delmotte V, Zhai P, Pörtner HO, Roberts D, Skea J et al. In press.
- IPMED 2011 Tomorrow, the Mediterranean Scenarios and projections for 2030. Paris.
- Ivar do Sul JA, Costa MF 2014 The present and future of microplastic pollution in the marine environment. *Environ. Pollut.* 185, 352–364. doi: [10.1016/j.envpol.2013.10.036](https://doi.org/10.1016/j.envpol.2013.10.036)
- Ivy-Ochs S, Kerschner H, Maisch M, Christl M, Kubik PW et al. 2009 Latest Pleistocene and Holocene glacier variations in the European Alps. *Quat. Sci. Rev.* 28, 2137–2149. doi: [10.1016/j.quascirev.2009.03.009](https://doi.org/10.1016/j.quascirev.2009.03.009)
- Izaurrealde RC, Thomson AM, Morgan JA, Fay PA, Polley HW et al. 2011 Climate Impacts on Agriculture: Implications for Forage and Rangeland Production. *Agron. J.* 103, 371. doi: [10.2134/agronj2010.0304](https://doi.org/10.2134/agronj2010.0304)
- Jacob D, Petersen J, Eggert B, Alias A, Christensen OB et al. 2014 EURO-CORDEX: new high-resolution climate change projections for European impact research. *Reg. Environ. Chang.* 14, 563–578. doi: [10.1007/s10113-013-0499-2](https://doi.org/10.1007/s10113-013-0499-2)
- Jacobbeit J, Hertig E, Seubert S, Lutz K 2014 Statistical down-

- scaling for climate change projections in the Mediterranean region: methods and results. *Reg. Environ. Chang.* 14, 1891–1906. doi: [10.1007/s10113-014-0605-0](https://doi.org/10.1007/s10113-014-0605-0)
- Jaeger EB, Seneviratne SI 2011 Impact of soil moisture–atmosphere coupling on European climate extremes and trends in a regional climate model. *Clim. Dyn.* 36, 1919–1939. doi: [10.1007/s00382-010-0780-8](https://doi.org/10.1007/s00382-010-0780-8)
- Javal M, Roques A, Haran J, Hérad F, Keena M et al. 2019 Complex invasion history of the Asian long-horned beetle: fifteen years after first detection in Europe. *J. Pest Sci.* (2004). 92, 173–187. doi: [10.1007/s10340-017-0917-1](https://doi.org/10.1007/s10340-017-0917-1)
- Jeanmonod D, Schlüssel A, Gamisans J 2011 Status and trends in the alien flora of Corsica. *EPPO Bull.* 41, 85–99. doi: [10.1111/j.1365-2338.2011.02440.x](https://doi.org/10.1111/j.1365-2338.2011.02440.x)
- Jebari S 2009 Water erosion modeling using fractal rainfall disaggregation - A study in semi-arid Tunisia.
- Jebari S, Berndtsson R, Bahri A, Boufaroua M 2010 Spatial soil loss risk and reservoir siltation in semi-arid Tunisia. *Hydrol. Sci. J.* 55, 121–137. doi: [10.1080/0262660903529049](https://doi.org/10.1080/0262660903529049)
- Jenkins P 1999 Trade and exotic species introductions, in *Invasive species and biodiversity management*, eds. Sandlund O, Schei P, Viken A (Kluwer, Dordrecht).
- Jeschke JM, Strayer DL 2005 Invasion success of vertebrates in Europe and North America. *Proc. Natl. Acad. Sci. U. S. A.* 102, 7198–7202. doi: [10.1073/pnas.0501271102](https://doi.org/10.1073/pnas.0501271102)
- Jeschke JM, Strayer DL 2006 Determinants of vertebrate invasion success in Europe and North America. *Glob. Chang. Biol.* 12, 1608–1619. doi: [10.1111/j.1365-2486.2006.01213.x](https://doi.org/10.1111/j.1365-2486.2006.01213.x)
- Jiang L, O'Neill BC 2017 Global urbanization projections for the Shared Socioeconomic Pathways. *Glob. Environ. Chang.* 42, 193–199. doi: [10.1016/j.gloenvcha.2015.03.008](https://doi.org/10.1016/j.gloenvcha.2015.03.008)
- Joffre RR, Rambal S 1993 How Tree Cover Influences the Water Balance of Mediterranean Rangelands. *Ecology* 74, 570–582. doi: [10.2307/1939317](https://doi.org/10.2307/1939317)
- Jones B, O'Neill BC 2016 Spatially explicit global population scenarios consistent with the Shared Socioeconomic Pathways. *Environ. Res. Lett.* 11, 84003. doi: [10.1088/1748-9326/11/8/084003](https://doi.org/10.1088/1748-9326/11/8/084003)
- Jones PD, Lister DH, Osborn TJ, Harpham C, Salmon M et al. 2012 Hemispheric and large-scale land surface air temperature variations: an extensive review and an update to 2010. *JGR Atmos.* 117, D05127. doi: [10.1029/2011JD017139](https://doi.org/10.1029/2011JD017139)
- Jordà G, Gomis D, Adloff F 2020 Vulnerability of marginal seas to sea-level rise: The case of the Mediterranean Sea. (in Rev.)
- Jordà G, Gomis D, Álvarez-Fanjul E 2012 The VANI2-ERA hind-cast of sea-level residuals: atmospheric forcing of sea-level variability in the Mediterranean Sea (1958–2008). *Sci. Mar.* 76S1, 133–146. doi: [10.3989/scimar.03612.19C](https://doi.org/10.3989/scimar.03612.19C)
- Jucker C, Lupi D 2011 Exotic Insects in Italy: An Overview on Their Environmental Impact. *Importance Biol. Interact. Study Biodivers.* doi: [10.5772/24263](https://doi.org/10.5772/24263)
- Justine J-L, Winsor L, Gey D, Gros P, Thévenot J 2018 Giant worms chez moi! Hammerhead flatworms (Platyhelminthes, Geoplanidae, *Bipalium* spp., *Diversipalium* spp.) in metropolitan France and overseas French territories. *PeerJ* 6, e4672. doi: [10.7717/peerj.4672](https://doi.org/10.7717/peerj.4672)
- Kairis O, Karavitis C, Salvati L, Kounalaki A, Kosmas K 2015 Exploring the Impact of Overgrazing on Soil Erosion and Land Degradation in a Dry Mediterranean Agro-Forest Landscape (Crete, Greece). *Arid L. Res. Manag.* 29, 360–374. doi: [10.1080/15324982.2014.968691](https://doi.org/10.1080/15324982.2014.968691)
- Kalabokas PD, Thouret V, Cammas J-P, Volz-Thomas A, Boulangier D et al. 2015 The geographical distribution of meteorological parameters associated with high and low summer ozone levels in the lower troposphere and the boundary layer over the eastern Mediterranean (Cairo case). *Tellus B Chem. Phys. Meteorol.* 67, 27853. doi: [10.3402/tellusb.v67.27853](https://doi.org/10.3402/tellusb.v67.27853)
- Kalimeris A, Ranieri E, Founda D, Norrant C 2017 Variability modes of precipitation along a Central Mediterranean area and their relations with ENSO, NAO, and other climatic patterns. *Atmos. Res.* 198, 56–80. doi: [10.1016/j.atmosres.2017.07.031](https://doi.org/10.1016/j.atmosres.2017.07.031)
- Kallos G, Solomos S, Kushta J, Mitsakou C, Spyrou C et al. 2014 Natural and anthropogenic aerosols in the Eastern Mediterranean and Middle East: Possible impacts. *Sci. Total Environ.* 488–489, 389–397. doi: [10.1016/j.scitotenv.2014.02.035](https://doi.org/10.1016/j.scitotenv.2014.02.035)
- Kambezidis HD, Psiloglou BE, Karagiannis D, Dumka UC, Kasikaoutis DG 2016 Recent improvements of the Meteorological Radiation Model for solar irradiance estimates under all-sky conditions. *Renew. Energy* 93, 142–158. doi: [10.1016/J.RENENE.2016.02.060](https://doi.org/10.1016/J.RENENE.2016.02.060)
- Kantiani L, Farré M, Asperger D, Rubio F, González S et al. 2008 Triclosan and methyl-triclosan monitoring study in the northeast of Spain using a magnetic particle enzyme immunoassay and confirmatory analysis by gas chromatography-mass spectrometry. *J. Hydrol.* 361, 1–9. doi: [10.1016/j.jhydrol.2008.07.016](https://doi.org/10.1016/j.jhydrol.2008.07.016)
- Karanasiou A, Querol X, Alastuey A, Perez N, Pey J et al. 2014 Particulate matter and gaseous pollutants in the Mediterranean Basin: Results from the MED-PARTICLES project. *Sci. Total Environ.* 488–489, 297–315. doi: [10.1016/j.scitotenv.2014.04.096](https://doi.org/10.1016/j.scitotenv.2014.04.096)
- Karanasiou AA, Siskos PA, Eleftheriadis K 2009 Assessment of source apportionment by Positive Matrix Factorization analysis on fine and coarse urban aerosol size fractions. *Atmos. Environ.* 43, 3385–3395. doi: [10.1016/j.atmosenv.2009.03.051](https://doi.org/10.1016/j.atmosenv.2009.03.051)
- Kark S, Solarz W, Chiron F, Clergeau P, Shirley S 2009 Alien Birds, Amphibians and Reptiles of Europe, in *Handbook of Alien Species in Europe*, ed. DAISIE (Springer Netherlands), 105–118. doi: [10.1007/978-1-4020-8280-1_8](https://doi.org/10.1007/978-1-4020-8280-1_8)
- Kaskaoutis DG, Dumka UC, Rashki A, Psiloglou BE, Gavriil A et al. 2019 Analysis of intense dust storms over the eastern Mediterranean in March 2018: Impact on radiative forcing and Athens air quality. *Atmos. Environ.* 209, 23–39. doi: [10.1016/J.ATMOSENV.2019.04.025](https://doi.org/10.1016/J.ATMOSENV.2019.04.025)
- Kassomenos PA, Kelessis A, Paschalidou AK, Petrakakis M 2011 Identification of sources and processes affecting particulate pollution in Thessaloniki, Greece. *Atmos. Environ.* 45, 7293–7300. doi: [10.1016/j.atmosenv.2011.08.034](https://doi.org/10.1016/j.atmosenv.2011.08.034)
- Katsanevakis S, Coll M, Piroddi C, Steenbeek J, Ben Rais Lasram F et al. 2014a Invading the Mediterranean Sea: biodi-

- versity patterns shaped by human activities. *Front. Mar. Sci.* 1. doi: [10.3389/fmars.2014.00032](https://doi.org/10.3389/fmars.2014.00032)
- Katsanevakis S, Rilov G, Edelist D 2018 Impacts of marine invasive alien species on European fisheries and aquaculture - plague or boon?, in *Engaging marine scientists and fishers to share knowledge and perceptions - early lessons*. CIESM Monograph n° 50, ed. Briand F [CIESM Publisher, Monaco and Paris], 125–132.
- Katsanevakis S, Tempeira F, Teixeira H 2016 Mapping the impact of alien species on marine ecosystems: The Mediterranean Sea case study. *Divers. Distrib.* 22, 694–707. doi: [10.1111/ddi.12429](https://doi.org/10.1111/ddi.12429)
- Katsanevakis S, Wallentinus I, Zenetos A, Leppäkoski E, Çınar ME et al. 2014b Impacts of invasive alien marine species on ecosystem services and biodiversity: a pan-European review. *Aquat. Invasions* 9, 391–423. doi: [10.3391/ai.2014.9.4.01](https://doi.org/10.3391/ai.2014.9.4.01)
- Katsanevakis S, Zenetos A, Belchior C, Cardoso AC 2013 Invading European Seas: Assessing pathways of introduction of marine aliens. *Ocean Coast. Manag.* 76, 64–74. doi: [10.1016/j.ocecoaman.2013.02.024](https://doi.org/10.1016/j.ocecoaman.2013.02.024)
- Kc S, Lutz W 2017 The human core of the shared socioeconomic pathways: Population scenarios by age, sex and level of education for all countries to 2100. *Glob. Environ. Chang.* 42, 181–192. doi: [10.1016/j.gloenvcha.2014.06.004](https://doi.org/10.1016/j.gloenvcha.2014.06.004)
- Kchih H, Perrino C, Cherif S 2015 Investigation of Desert Dust Contribution to Source Apportionment of PM10 and PM2.5 from a Southern Mediterranean Coast. *Aerosol Air Qual. Res.* 15, 454–464. doi: [10.4209/aaqr.2014.10.0255](https://doi.org/10.4209/aaqr.2014.10.0255)
- Keeley JE, Bond WJ, Bradstock RA, Pausas JG, Rundel PW 2012 *Fire in Mediterranean ecosystems: Ecology, evolution and management*. Cambridge, UK: Cambridge University Press doi: [10.1017/cbo9781139033091](https://doi.org/10.1017/cbo9781139033091)
- Keenan TF, Maria Serra J, Lloret F, Ninyerola M, Sabate S 2011 Predicting the future of forests in the Mediterranean under climate change, with niche- and process-based models: CO₂ matters! *Glob. Chang. Biol.* 17, 565–579. doi: [10.1111/j.1365-2486.2010.02254.x](https://doi.org/10.1111/j.1365-2486.2010.02254.x)
- Kelepertzis E 2014 Accumulation of heavy metals in agricultural soils of Mediterranean: Insights from Argolida basin, Peloponnese, Greece. *Geoderma* 221–222, 82–90. doi: [10.1016/j.geoderma.2014.01.007](https://doi.org/10.1016/j.geoderma.2014.01.007)
- Keller RP, Geist J, Jeschke JM, Kühn I 2011 Invasive species in Europe: ecology, status, and policy. *Environ. Sci. Eur.* 23, 23. doi: [10.1186/2190-4715-23-23](https://doi.org/10.1186/2190-4715-23-23)
- Kelley C, Ting M, Seager R, Kushnir Y 2012 The relative contributions of radiative forcing and internal climate variability to the late 20th century winter drying of the Mediterranean region. *Clim. Dyn.* 38, 2001–2015. doi: [10.1007/s00382-011-1221-z](https://doi.org/10.1007/s00382-011-1221-z)
- Kendon EJ, Roberts NM, Fowler HJ, Roberts MJ, Chan SC et al. 2014 Heavier summer downpours with climate change revealed by weather forecast resolution model. *Nat. Clim. Chang.* 4, 570–576. doi: [10.1038/nclimate2258](https://doi.org/10.1038/nclimate2258)
- Kenis M, Auger-Rozenberg M-A, Roques A, Timms L, Péré C et al. 2009 Ecological effects of invasive alien insects. *Biol. Invasions* 11, 21–45. doi: [10.1007/s10530-008-9318-y](https://doi.org/10.1007/s10530-008-9318-y)
- Kenis M, Branco M 2010 Impact of alien terrestrial arthropods in Europe. Chapter 5. *BioRisk* 4, 51–71. doi: [10.3897/biorisk.4.42](https://doi.org/10.3897/biorisk.4.42)
- Kerdelhué C, Boivin T, Burban C 2014 Contrasted invasion processes imprint the genetic structure of an invasive scale insect across southern Europe. *Heredity (Edinb)*. 113, 390–400. doi: [10.1038/hdy.2014.39](https://doi.org/10.1038/hdy.2014.39)
- Kirichenko N, Augustin S, Kenis M 2019 Invasive leafminers on woody plants: a global review of pathways, impact, and management. *J. Pest Sci.* (2004). 92, 93–106. doi: [10.1007/s10340-018-1009-6](https://doi.org/10.1007/s10340-018-1009-6)
- Klapfer R, Welch L 2011 Emerging contaminant threats and the Great Lakes. https://c.ymcdn.com/sites/www.productstewardship.us/resource/collection/FFDF28A1-9926-46E7-87F9-70C7BBD95491/Emerging_Contaminant_Threats_and_the_Great_Lakes.pdf
- Kleitou P, Hall-Spencer J, Rees S, Sfenthourakis S, Demetriou A et al. 2019 Tackling the lionfish invasion in the Mediterranean. The EU-LIFE RELIONMED Project: progress and results. in *Proceedings of the 1st Mediterranean Symposium on the Non-Indigenous Species* (Antalya, Turkey), 65–70.
- Kleitou DC, Hall-Spencer JM, Kleitou P 2016 A lionfish (*Pterois miles*) invasion has begun in the Mediterranean Sea. *Mar. Biodivers. Rec.* 9. doi: [10.1186/s41200-016-0065-y](https://doi.org/10.1186/s41200-016-0065-y)
- Knippertz P, Ulbrich U, Marques F, Corte-Real J 2003 Decadal changes in the link between El Niño and springtime North Atlantic oscillation and European–North African rainfall. *Int. J. Climatol.* 23, 1293–1311. doi: [10.1002/joc.944](https://doi.org/10.1002/joc.944)
- Knippertz P, Ulbrich U, Speth P 2000 Changing cyclones and surface wind speeds over the North Atlantic and Europe in a transient GHG experiment. *Clim. Res.* 15, 109–122. doi: [10.3354/cr015109](https://doi.org/10.3354/cr015109)
- Knutson TR, Zeng F, Wittenberg AT, Knutson TR, Zeng F et al. 2013 Multimodel Assessment of Regional Surface Temperature Trends: CMIP3 and CMIP5 Twentieth-Century Simulations. *J. Clim.* 26, 8709–8743. doi: [10.1175/JCLI-D-12-00567.1](https://doi.org/10.1175/JCLI-D-12-00567.1)
- Knutti R, Sedláček J 2012 Robustness and uncertainties in the new CMIP5 climate model projections. *Nat. Clim. Chang.* 3, 369–373. doi: [10.1038/nclimate1716](https://doi.org/10.1038/nclimate1716)
- Knutti R, Sedláček J, Sanderson BM, Lorenz R, Fischer EM et al. 2017 A climate model projection weighting scheme accounting for performance and interdependence. *Geophys. Res. Lett.* 44, 1909–1918.
- Koçak M, Mihalopoulos N, Kubilay NN 2007a Chemical composition of the fine and coarse fraction of aerosols in the north-eastern Mediterranean. *Atmos. Environ.* 41, 7351–7368. doi: [10.1016/j.atmosenv.2007.05.011](https://doi.org/10.1016/j.atmosenv.2007.05.011)
- Koçak M, Mihalopoulos N, Kubilay NN 2007b Contributions of natural sources to high PM10 and PM2.5 events in the eastern Mediterranean. *Atmos. Environ.* 41, 3806–3818. doi: [10.1016/j.atmosenv.2007.01.009](https://doi.org/10.1016/j.atmosenv.2007.01.009)
- Koelmans AA, Besseling E, Fockema E, Kooi M, Mintenig S et al. 2017 Risks of plastic debris: unravelling fact, opinion, perception, and belief. *Environ. Sci. Technol.* 51, 11513–11519. doi: [10.1021/acs.est.7b02219](https://doi.org/10.1021/acs.est.7b02219)
- Kok K, Pedde S, Gramberger M, Harrison PA, Holman IP 2019

- New European socio-economic scenarios for climate change research: Operationalising concepts to extend the shared socio-economic pathways. *Reg. Environ. Chang.* 19, 643–654. doi: [10.1007/s10113-018-1400-0](https://doi.org/10.1007/s10113-018-1400-0)
- Kok K, Rothman DS, Patel M 2006 Multi-scale narratives from an IA perspective: Part I. European and Mediterranean scenario development. *Futures* 38, 261–284. doi: [10.1016/j.futures.2005.07.001](https://doi.org/10.1016/j.futures.2005.07.001)
- Konstantinou IK, Hela DG, Albanis TA 2006 The status of pesticide pollution in surface waters (rivers and lakes) of Greece. Part I. Review on occurrence and levels. *Environ. Pollut.* 141, 555–570. doi: [10.1016/j.envpol.2005.07.024](https://doi.org/10.1016/j.envpol.2005.07.024)
- Kostopoulou E, Giannakopoulos C, Hatzaki M, Karali A, Hadjinicolaou P et al. 2014 Spatio-temporal patterns of recent and future climate extremes in the eastern Mediterranean and Middle East region. *Nat. Hazards Earth Syst. Sci.* 14, 1565–1577. doi: [10.5194/nhess-14-1565-2014](https://doi.org/10.5194/nhess-14-1565-2014)
- Kostopoulou E, Jones PD 2007 Comprehensive analysis of the climate variability in the eastern Mediterranean. Part II: relationships between atmospheric circulation patterns and surface climatic elements. *Int. J. Climatol.* 27, 1351–1371. doi: [10.1002/joc.1466](https://doi.org/10.1002/joc.1466)
- Kotlarski S, Szabó P, Herrera S, Räty O, Keuler K et al. 2019 Observational uncertainty and regional climate model evaluation: A pan-European perspective. *Int. J. Climatol.* 39, 3730–3749. doi: [10.1002/joc.5249](https://doi.org/10.1002/joc.5249)
- Kouroutzoglou J, Flocas HA, Keay K, Simmonds I, Hatzaki M 2010 Climatological aspects of explosive cyclones in the Mediterranean. *Int. J. Climatol.* 31, 1785–1802. doi: [10.1002/joc.2203](https://doi.org/10.1002/joc.2203)
- Kouroutzoglou J, Flocas HA, Keay K, Simmonds I, Hatzaki M 2011 Climatological aspects of explosive cyclones in the Mediterranean. *Int. J. Climatol.* 31, 1785–1802. doi: [10.1002/joc.2203](https://doi.org/10.1002/joc.2203)
- Koutavas A 2013 CO₂ fertilization and enhanced drought resistance in Greek firs from Cephalonia Island, Greece. *Glob. Chang. Biol.* 19, 529–539. doi: [10.1111/gcb.12053](https://doi.org/10.1111/gcb.12053)
- Koutsodendris A, Papatheodorou G, Kougiourouki O, Georgiadis M 2008 Benthic marine litter in four Gulfs in Greece, Eastern Mediterranean; abundance, composition and source identification. *Estuar. Coast. Shelf Sci.* 77, 501–512. doi: [10.1016/j.ecss.2007.10.011](https://doi.org/10.1016/j.ecss.2007.10.011)
- Kress N, Manca BB, Klein B, Deponte D 2003 Continuing influence of the changed thermohaline circulation in the eastern Mediterranean on the distribution of dissolved oxygen and nutrients: Physical and chemical characterization of the water masses. *JGR Ocean.* 108. doi: [10.1029/2002jc001397](https://doi.org/10.1029/2002jc001397)
- Kriticos DJ, Brunel S, Ota N, Fried G, Oude Lansink AGJM et al. 2015 Downscaling pest risk analyses: Identifying current and future potentially suitable habitats for *Parthenium hysterophorus* with particular reference to Europe and North Africa. *PLoS One* 10, e0132807. doi: [10.1371/journal.pone.0132807](https://doi.org/10.1371/journal.pone.0132807)
- Kuglitsch FG, Toreti A, Xoplaki E, Della-Marta PM, Zerefos CS et al. 2010 Heat wave changes in the eastern Mediterranean since 1960. *Geophys. Res. Lett.* 37, 1–5. doi: [10.1029/2009GL041841](https://doi.org/10.1029/2009GL041841)
- Kuper M, Ameur F, Hammani A 2017 *Unraveling the enduring paradox of increased pressure on groundwater through efficient drip irrigation*. www.routledge.com/Drip-Irrigation-for-Agriculture-Untold-Stories-of-Efficiency-Innovation/Venot-Kuper-Zwarteeveen/p/book/9781138687073
- Laabir M, Collos Y, Masseret E, Grzebyk D, Abadie E et al. 2013 Influence of environmental factors on the paralytic shellfish toxin content and profile of *Alexandrium catenella* (Dinophyceae) Isolated from the Mediterranean Sea. *Mar. Drugs* 11, 1583–1601. doi: [10.3390/md11051583](https://doi.org/10.3390/md11051583)
- Lacombe H 1990 A model of the world ocean, water, salt heat, and wind in the Mediterranean. *Oceanus* 33, 26–36.
- Lamb PJ, Peppler RA 1987 North Atlantic Oscillation: Concept and an Application. *Bull. Am. Meteorol. Soc.* 68, 1218–1225. doi: [10.1175/1520-0477\(1987\)068<1218:naocaa>2.0.co;2](https://doi.org/10.1175/1520-0477(1987)068<1218:naocaa>2.0.co;2)
- Lambdon PW, Hulme PE 2006 Predicting the invasion success of Mediterranean alien plants from their introduction characteristics. *Ecography (Cop.)* 29, 853–865. doi: [10.1111/j.2006.0906-7590.04614.x](https://doi.org/10.1111/j.2006.0906-7590.04614.x)
- Lambdon PW, Pyšek P, Basnou C, Hejda M, Arianoutsou M et al. 2008 Alien flora of Europe: species diversity, temporal trends, geographical patterns and research needs. *Preslia* 80, 101–149.
- Lapiedra O, Sol D, Traveset A, Vilà M 2015 Random processes and phylogenetic loss caused by plant invasions. *Glob. Ecol. Biogeogr.* 24, 774–785. doi: [10.1111/geb.12310](https://doi.org/10.1111/geb.12310)
- Lasanta T, Arnáez J, Pascual N, Ruiz-Flaño P, Errea MP et al. 2017 Space-time process and drivers of land abandonment in Europe. *Catena* 149, 810–823. doi: [10.1016/j.catena.2016.02.024](https://doi.org/10.1016/j.catena.2016.02.024)
- Laza-Martínez A, David H, Riobo P, Miguel I, Orive E et al. 2016 Characterization of a Strain of *Fukuyoa paulensis* (Dinophyceae) from the Western Mediterranean Sea. *J. Eukaryot. Microbiol.* 63, 481–497. doi: [10.1111/jeu.12292](https://doi.org/10.1111/jeu.12292)
- Le Cozannet G, Thieblemont R, Rohmer J, Idier D, Manceau J-C et al. 2019 Low-end probabilistic sea-level projections. *Water* 11, 1507.
- Le Houérou HN 1995 *Bioclimatologie et biogéographie des steppes arides du Nord de l'Afrique: diversité biologique, développement durable et désertification*. Montpellier: CIHEAM.
- Le Houérou HN 1996 Climate change, drought and desertification. *J. Arid Environ.* 34, 133–185.
- Leckebusch GC, Koffi B, Ulbrich U, Pinto JG, Spangehl T et al. 2006 Analysis of frequency and intensity of European winter storm events from a multi-model perspective, at synoptic and regional scales. *Clim. Res.* 31, 59–74. doi: [10.3354/cr031059](https://doi.org/10.3354/cr031059)
- Ledoux JB, Aurelle D, Bensoussan N, Marschal C, Féral JP et al. 2015 Potential for adaptive evolution at species range margins: Contrasting interactions between red coral populations and their environment in a changing ocean. *Ecol. Evol.* 5, 1178–1192. doi: [10.1002/ece3.1324](https://doi.org/10.1002/ece3.1324)
- Lehtijärvi A, Oskay F, Doğmuş Lehtijärvi HT, Aday Kaya AG, Pecori F et al. 2018 *Ceratocystis platani* is killing plane trees in Istanbul (Turkey). *For. Pathol.* 48, e12375. doi: [10.1111/efp.12375](https://doi.org/10.1111/efp.12375)
- Leiblein-Wild MC, Steinkamp J, Hickler T, Tackenberg O 2016

- Modelling the potential distribution, net primary production and phenology of common ragweed with a physiological model. *J. Biogeogr.* 43, 544–554. doi: [10.1111/jbi.12646](https://doi.org/10.1111/jbi.12646)
- Leimbach M, Kriegler E, Roming N, Schwanitz J 2017 Future growth patterns of world regions – A GDP scenario approach. *Glob. Environ. Chang.* 42, 215–225. doi: [10.1016/j.gloenvcha.2015.02.005](https://doi.org/10.1016/j.gloenvcha.2015.02.005)
- Lejeusne C, Chevaldonné P 2006 Brooding crustaceans in a highly fragmented habitat: The genetic structure of Mediterranean marine cave-dwelling mysid populations. *Mol. Ecol.* 15, 4123–4140. doi: [10.1111/j.1365-294x.2006.03101.x](https://doi.org/10.1111/j.1365-294x.2006.03101.x)
- Lekkas T, Kolokythas G, Nikolaou A, Kostopoulou M, Kotrikla A et al. 2004 Evaluation of the pollution of the surface waters of Greece from the priority compounds of list II, 76/464/EEC directive, and other toxic compounds. *Environ. Int.* 30, 995–1007. doi: [10.1016/j.envint.2004.04.001](https://doi.org/10.1016/j.envint.2004.04.001)
- Lelieveld J 2002 Global Air Pollution Crossroads over the Mediterranean. *Science (80-.).* 298, 794–799. doi: [10.1126/science.1075457](https://doi.org/10.1126/science.1075457)
- Lelieveld J, Hadjinicolaou P, Kostopoulou E, Chenoweth J, El Maayar M et al. 2012 Climate change and impacts in the Eastern Mediterranean and the Middle East. *Clim. Change* 114, 667–687. doi: [10.1007/s10584-012-0418-4](https://doi.org/10.1007/s10584-012-0418-4)
- Lelieveld J, Hadjinicolaou P, Kostopoulou E, Giannakopoulos C, Pozzer A et al. 2014 Model projected heat extremes and air pollution in the eastern Mediterranean and Middle East in the twenty-first century. *Reg. Environ. Chang.* 14, 1937–1949. doi: [10.1007/s10113-013-0444-4](https://doi.org/10.1007/s10113-013-0444-4)
- Lelieveld J, Proestos Y, Hadjinicolaou P, Tanarhte M, Tyrlis E et al. 2016 Strongly increasing heat extremes in the Middle East and North Africa (MENA) in the 21st century. *Clim. Change* 137, 245–260. doi: [10.1007/s10584-016-1665-6](https://doi.org/10.1007/s10584-016-1665-6)
- Lenderink G, Belušić D, Fowler HJ, Kjellström E, Lind P et al. 2019 Systematic increases in the thermodynamic response of hourly precipitation extremes in an idealized warming experiment with a convection-permitting climate model. *Environ. Res. Lett.* 14, 074012. doi: [10.1088/1748-9326/ab214a](https://doi.org/10.1088/1748-9326/ab214a)
- Lesieur V, Lombaert E, Guillemaud T, Courtial B, Strong W et al. 2019 The rapid spread of *Leptoglossus occidentalis* in Europe: a bridgehead invasion. *J. Pest Sci. (2004).* 92, 189–200. doi: [10.1007/s10340-018-0993-x](https://doi.org/10.1007/s10340-018-0993-x)
- Levesque B, Gervais M-C, Chevalier P, Gauvin D, Anassour-Laouan-Sidi E et al. 2014 Prospective study of acute health effects in relation to exposure to cyanobacteria. *Sci. Total Environ.* 466–467, 397–403. doi: [10.1016/j.scitotenv.2013.07.045](https://doi.org/10.1016/j.scitotenv.2013.07.045)
- Levin Z, Teller A, Ganor E, Graham B, Andreeae MO et al. 2003 Role of aerosol size and composition in nucleation scavenging within clouds in a shallow cold front. *JGR Atmos.* 108, 4700. doi: [10.1029/2003JD003647](https://doi.org/10.1029/2003JD003647)
- Li J, Carlson BE, Dubovnik O, Lacis AA 2014 Recent trends in aerosol optical properties derived from AERONET measurements. *Atmos. Chem. Phys.* 14, 12271–12289. doi: [10.5194/acp-14-12271-2014](https://doi.org/10.5194/acp-14-12271-2014)
- Licciardello F, Antoci ML, Brugaletta L, Cirelli GL 2011 Evaluation of groundwater contamination in a coastal area of south-eastern Sicily. *J. Environ. Sci. Heal. Part. B Pestic.* Food Contam. Agric. Wastes 46, 498–508. doi: [10.1080/03601234.2011.583870](https://doi.org/10.1080/03601234.2011.583870)
- Liebhold AM, Yamanaka T, Roques A, Augustin S, Chown SL et al. 2016 Global compositional variation among native and non-native regional insect assemblages emphasizes the importance of pathways. *Biol. Invasions* 18, 893–905. doi: [10.1007/s10530-016-1079-4](https://doi.org/10.1007/s10530-016-1079-4)
- Liebig MA, Tanaka DL, Wienhold BJ 2004 Tillage and cropping effects on soil quality indicators in the northern Great Plains. *Soil Tillage Res.* 78, 131–141. doi: [10.1016/j.still.2004.02.002](https://doi.org/10.1016/j.still.2004.02.002)
- Lillo F, Faraone FP, Lo Valvo M 2011 Can the introduction of Xenopus laevis affect native amphibian populations? Reduction of reproductive occurrence in presence of the invasive species. *Biol. Invasions* 13, 1533–1541. doi: [10.1007/s10530-010-9911-8](https://doi.org/10.1007/s10530-010-9911-8)
- Lionello P 2012 *The Climate of the Mediterranean Region: From the Past to the Future.* Elsevier Science. doi: [10.1016/C2011-0-06210-5](https://doi.org/10.1016/C2011-0-06210-5)
- Lionello P, Abrantes FG, Congedi L, Dulac F, Gačić M et al. 2012a Introduction: Mediterranean climate-background information, in *The Climate of the Mediterranean Region: From the Past to the Future*, ed. Lionello P [Elsevier Science]. doi: [10.1016/B978-0-12-416042-2.00012-4](https://doi.org/10.1016/B978-0-12-416042-2.00012-4)
- Lionello P, Cavalieri L, Nissen KM, Pino C, Raicich F et al. 2012b Severe marine storms in the Northern Adriatic: characteristics and trends. *Phys. Chem. Earth, Parts A/B/C* 40–41, 93–105. doi: [10.1016/j.pce.2010.10.002](https://doi.org/10.1016/j.pce.2010.10.002)
- Lionello P, Cogo S, Galati MB, Sanna A 2008 The Mediterranean surface wave climate inferred from future scenario simulations. *Glob. Planet. Change* 63, 152–162. doi: [10.1016/j.gloplacha.2008.03.004](https://doi.org/10.1016/j.gloplacha.2008.03.004)
- Lionello P, Conte D, Marzo L, Scarascia L 2017 The contrasting effect of increasing mean sea level and decreasing storminess on the maximum water level during storms along the coast of the Mediterranean Sea in the mid 21st century. *Glob. Planet. Change* 151, 80–91. doi: [10.1016/j.gloplacha.2016.06.012](https://doi.org/10.1016/j.gloplacha.2016.06.012)
- Lionello P, Dalan F, Elvini E 2002 Cyclones in the Mediterranean region: the present and the doubled CO₂ climate scenarios. *Clim. Res.* 22, 147–159. doi: [10.3354/cr022147](https://doi.org/10.3354/cr022147)
- Lionello P, Galati MB 2008 Links of the significant wave height distribution in the Mediterranean sea with the Northern Hemisphere teleconnection patterns. *Adv. Geosci.* 17, 13–18. doi: [10.5194/adgeo-17-13-2008](https://doi.org/10.5194/adgeo-17-13-2008)
- Lionello P, Giorgi F 2007 Winter precipitation and cyclones in the Mediterranean region: Future climate scenarios in a regional simulation. *Adv. Geosci.* 12, 153–158. doi: [10.5194/adgeo-12-153-2007](https://doi.org/10.5194/adgeo-12-153-2007)
- Lionello P, Malanotte-Rizzoli P, Boscolo R, Alpert P, Artale V et al. 2006 The Mediterranean climate: An overview of the main characteristics and issues, in *Mediterranean climate variability*, eds. Lionello P, Malanotte-Rizzoli P, Boscolo R [Elsevier], 1–26. doi: [10.1016/s1571-9197\(06\)80003-0](https://doi.org/10.1016/s1571-9197(06)80003-0)
- Lionello P, Sanna A 2005 Mediterranean wave climate variability and its links with NAO and Indian Monsoon. *Clim. Dyn.* 25, 611–623. doi: [10.1007/s00382-005-0025-4](https://doi.org/10.1007/s00382-005-0025-4)

- Lionello P, Scarascia L 2018 The relation between climate change in the Mediterranean region and global warming. *Reg. Environ. Chang.* 18, 1481–1493. doi: [10.1007/s10113-018-1290-1](https://doi.org/10.1007/s10113-018-1290-1)
- Lionello P, Scarascia L 2020 The relation of climate extremes with global warming in the Mediterranean region and its north versus south contrast. *Reg. Environ. Chang.* 20, 31. doi: [10.1007/s10113-020-01610-z](https://doi.org/10.1007/s10113-020-01610-z)
- Lionello P, Trigo IF, Gil V, Liberato MLR, Nissen KM et al. 2016 Objective climatology of cyclones in the Mediterranean region: A consensus view among methods with different system identification and tracking criteria. *Tellus A Dyn. Meteorol. Oceanogr.* 68, 29391. doi: [10.3402/tellusa.v68.29391](https://doi.org/10.3402/tellusa.v68.29391)
- Lioy S, Manino A, Porporato M, Laurino D, Romano A et al. 2019 Establishing surveillance areas for tackling the invasion of *Vespa velutina* in outbreaks and over the border of its expanding range. *NeoBiota* 46, 51–69. doi: [10.3897/neobiota.46.33099](https://doi.org/10.3897/neobiota.46.33099)
- Lippmann M, Chen L-C 2009 Health effects of concentrated ambient air particulate matter (CAPs) and its components. *Crit. Rev. Toxicol.* 39, 865–913. doi: [10.3109/10408440903300080](https://doi.org/10.3109/10408440903300080)
- Liubartseva S, Coppini G, Lecci R, Clementi E 2018 Tracking plastics in the Mediterranean: 2D Lagrangian model. *Mar. Pollut. Bull.* 129, 151–162. doi: [10.1016/j.marpolbul.2018.02.019](https://doi.org/10.1016/j.marpolbul.2018.02.019)
- Lloret F, Médail F, Brundu G, Camarda I, Moragues E et al. 2005 Species attributes and invasion success by alien plants on Mediterranean islands. *J. Ecol.* 93, 512–520. doi: [10.1111/j.1365-2745.2005.00979.x](https://doi.org/10.1111/j.1365-2745.2005.00979.x)
- Lloret F, Médail F, Brundu G, Hulme PE 2004a Local and regional abundance of exotic plant species on Mediterranean islands: are species traits important? *Glob. Ecol. Biogeogr.* 13, 37–45. doi: [10.1111/j.1466-882x.2004.00064.x](https://doi.org/10.1111/j.1466-882x.2004.00064.x)
- Lloret J, Palomera I, Salat J, Sole I 2004b Impact of freshwater input and wind on landings of anchovy (*Engraulis encrasicolus*) and sardine (*Sardina pilchardus*) in shelf waters surrounding the Ebre (Ebro) River delta (north-western Mediterranean). *Fish. Oceanogr.* 13, 102–110. doi: [10.1046/j.1365-2419.2003.00279.x](https://doi.org/10.1046/j.1365-2419.2003.00279.x)
- Loepfe L, Martínez-Vilalta J, Oliveres J, Piñol J, Lloret F 2010 Feedbacks between fuel reduction and landscape homogenisation determine fire regimes in three Mediterranean areas. *For. Ecol. Manage.* 259, 2366–2374. doi: [10.1016/j.foreco.2010.03.009](https://doi.org/10.1016/j.foreco.2010.03.009)
- Loepfe L, Martínez-Vilalta J, Piñol J 2011 An integrative model of human-influenced fire regimes and landscape dynamics. *Environ. Model. Softw.* 26, 1028–1040. doi: [10.1016/j.envsoft.2011.02.015](https://doi.org/10.1016/j.envsoft.2011.02.015)
- López-Parages J, Rodríguez-Fonseca B 2012 Multidecadal modulation of El Niño influence on the Euro-Mediterranean rainfall. *Geophys. Res. Lett.* 39. doi: [10.1029/2011GL050049](https://doi.org/10.1029/2011GL050049)
- Lopez B, Ollivier P, Togola A, Baran N, Ghement J-P 2015 Screening of French groundwater for regulated and emerging contaminants. *Sci. Total Environ.* 518–519, 562–573. doi: [10.1016/j.scitotenv.2015.01.110](https://doi.org/10.1016/j.scitotenv.2015.01.110)
- Löptien U, Zolina O, Gulev S, Latif M, Soloviov V 2008 Cyclone life cycle characteristics over the Northern Hemisphere in coupled GCMs. *Clim. Dyn.* 31, 507–532. doi: [10.1007/s00382-007-0355-5](https://doi.org/10.1007/s00382-007-0355-5)
- Louchart X, Voltz M, Andrieux P, Moussa R 2001 Herbicide transport to surface waters at field and watershed scales in a Mediterranean vineyard area. *J. Environ. Qual.* 30, 982–91. <http://www.ncbi.nlm.nih.gov/pubmed/11401289>
- Loumou A, Giourga C 2003 Olive groves: "The life and identity of the Mediterranean." *Agric. Human Values* 20, 87–95. doi: [10.1023/a:1022444005336](https://doi.org/10.1023/a:1022444005336)
- Lounibos LP 2002 Invasions by Insect Vectors of Human Disease. *Annu. Rev. Entomol.* 47, 233–266. doi: [10.1146/annurev.ento.47.091201.145206](https://doi.org/10.1146/annurev.ento.47.091201.145206)
- Lowenthal DH, Gertler AW, Labib MW 2014 Particulate matter source apportionment in Cairo: recent measurements and comparison with previous studies. *Int. J. Environ. Sci. Technol.* 11, 657–670. doi: [10.1007/s13762-013-0272-6](https://doi.org/10.1007/s13762-013-0272-6)
- Lu J, Vecchi GA, Reichler T 2007a Correction to "Expansion of the Hadley cell under global warming." *Geophys. Res. Lett.* 34. doi: [10.1029/2007gl030931](https://doi.org/10.1029/2007gl030931)
- Lu J, Vecchi GA, Reichler T 2007b Expansion of the Hadley cell under global warming. *Geophys. Res. Lett.* 34, L06805. doi: [10.1029/2006GL028443](https://doi.org/10.1029/2006GL028443)
- Ludwig W, Dumont E, Meybeck M, Heussner S 2009 River discharges of water and nutrients to the Mediterranean and Black Sea: Major drivers for ecosystem changes during past and future decades? *Prog. Oceanogr.* 80, 199–217. doi: [10.1016/j.pocean.2009.02.001](https://doi.org/10.1016/j.pocean.2009.02.001)
- Luterbacher J, Werner JP, Smerdon JE 2016 European summer temperatures since Roman times. *Environ. Res. Lett.* 11, 2 12. doi: [10.1088/1748-9326/11/2/024001](https://doi.org/10.1088/1748-9326/11/2/024001)
- Luterbacher J, Xoplaki E 2003 500-year Winter Temperature and Precipitation Variability over the Mediterranean area and its Connection to the Large-scale Atmospheric Circulation, in *Mediterranean Climate. Variability and Trends*, ed. Bolle HJ (Berlin: Springer Verlag), 133–153.
- Lutz W, Amran G, Bélanger A, Conte A, Gailey N et al. 2019 Demographic Scenarios for the EU - Migration, Population and Education. EUR 29739 EN, Publications Office, Luxembourg doi: [10.2760/590301](https://doi.org/10.2760/590301)
- M'Rabet C, Kéfi-Daly YO, Couet D, Gueroun SKM, Pringault O 2019 Consequences of a contaminant mixture of bisphenol A (BPA) and di-(2-ethylhexyl) phthalate (DEHP), two plastic-derived chemicals, on the diversity of coastal phytoplankton. *Mar. Pollut. Bull.* 138, 385–396. doi: [10.1016/j.marpolbul.2018.11.035](https://doi.org/10.1016/j.marpolbul.2018.11.035)
- Macías DM, García-Gorriz E, Stips A 2013 Understanding the causes of recent warming of Mediterranean waters. How much could be attributed to climate change? *PLoS One* 8, e81591. doi: [10.1371/journal.pone.0081591](https://doi.org/10.1371/journal.pone.0081591)
- Macías DM, García-Gorriz E, Stips A 2018 Deep winter convection and phytoplankton dynamics in the NW Mediterranean Sea under present climate and future (horizon 2030) scenarios. *Sci. Rep.* 8, 6626. doi: [10.1038/s41598-018-24965-0](https://doi.org/10.1038/s41598-018-24965-0)
- Maggi V, Colucci RR, Scoto F, Giudice G, Randazzo L 2018 Chapter 19 - Ice Caves in Italy, in, eds. Peršoiu A, Lauritzen S-EBT-IC (Elsevier), 399–423.

- doi: [10.1016/B978-0-12-811739-2.00019-X](https://doi.org/10.1016/B978-0-12-811739-2.00019-X)
- Magnusson M, Heimann K, Quayle P, Negri AP 2010 Additive toxicity of herbicide mixtures and comparative sensitivity of tropical benthic microalgae. *Mar. Pollut. Bull.* 60, 1978–1987. doi: [10.1016/j.marpolbul.2010.07.031](https://doi.org/10.1016/j.marpolbul.2010.07.031)
- Maheras P, Flocas HA, Patrikas I, Anagnostopoulou C 2001 A 40 year objective climatology of surface cyclones in the Mediterranean region: spatial and temporal distribution. *Int. J. Climatol.* 21, 109–130. doi: [10.1002/joc.599](https://doi.org/10.1002/joc.599)
- Mahmoud KF, Alfaro SC, Favez O, Abdel Wahab MM, Sciaro J 2008 Origin of black carbon concentration peaks in Cairo (Egypt). *Atmos. Res.* 89, 161–169. doi: [10.1016/J.ATMOSRES.2008.01.004](https://doi.org/10.1016/J.ATMOSRES.2008.01.004)
- Malagó A, Bouraoui F, Grizzetti B, De Roo A 2019 Modelling nutrient fluxes into the Mediterranean Sea. *J. Hydrol. Reg. Stud.* 22, 100592. doi: [10.1016/j.ejrh.2019.01.004](https://doi.org/10.1016/j.ejrh.2019.01.004)
- Malek Ž, Verburg PH 2018 Adaptation of land management in the Mediterranean under scenarios of irrigation water use and availability. *Mitig. Adapt. Strateg. Glob. Chang.* 23, 821–837. doi: [10.1007/s11027-017-9761-0](https://doi.org/10.1007/s11027-017-9761-0)
- Manachini B, Billeci N, Palla F 2013 Exotic insect pests: The impact of the Red Palm Weevil on natural and cultural heritage in Palermo (Italy). *J. Cult. Herit.* 14, e177–e182. doi: [10.1016/j.culher.2012.11.028](https://doi.org/10.1016/j.culher.2012.11.028)
- Manara V, Brunetti M, Celozzi M, Maugeri M, Sánchez-Lorenzo A et al. 2016 Detection of dimming / brightening in Italy from homogenized all-sky and clear-sky surface solar radiation records and underlying causes (1959–2013). *Atmos. Chem. Phys.* 16, 11145–11161. doi: [10.5194/acp-16-11145-2016](https://doi.org/10.5194/acp-16-11145-2016)
- Mandaric L, Kalogianni E, Skoulikidis N, Petrovic M, Sabater S 2019 Contamination patterns and attenuation of pharmaceuticals in a temporary Mediterranean river. *Sci. Total Environ.* 647, 561–569. doi: [10.1016/j.scitotenv.2018.07.308](https://doi.org/10.1016/j.scitotenv.2018.07.308)
- Manodori L, Gambaro A, Piazza R, Ferrari S, Stortini AM et al. 2006 PCBs and PAHs in sea-surface microlayer and sub-surface water samples of the Venice Lagoon (Italy). *Mar. Pollut. Bull.* 52, 184–192. doi: [10.1016/j.marpolbul.2005.08.017](https://doi.org/10.1016/j.marpolbul.2005.08.017)
- Mansui J, Molcard A, Ourmières Y 2015 Modelling the transport and accumulation of floating marine debris in the Mediterranean basin. *Mar. Pollut. Bull.* 91, 249–257. doi: [10.1016/j.marpolbul.2014.11.037](https://doi.org/10.1016/j.marpolbul.2014.11.037)
- Manzano S, Carrión JS, García-Murillo P, López-Merino L 2019 When dynamism is the baseline: long-term ecology of a Mediterranean seasonal wetland in the Doñana National Park (Southwestern Europe). *Biodivers. Conserv.* 28, 501–522. doi: [10.1007/s10531-018-1674-z](https://doi.org/10.1007/s10531-018-1674-z)
- Maraun D, Widmann M, Gutiérrez JM 2019 Statistical downscaling skill under present climate conditions: A synthesis of the VALUE perfect predictor experiment. *Int. J. Climatol.* 39, 3692–3703. doi: [10.1002/joc.5877](https://doi.org/10.1002/joc.5877)
- Marba N, Duarte CM 2010 Mediterranean warming triggers seagrass (*Posidonia oceanica*) shoot mortality. *Glob. Chang. Biol.* 16, 2366–2375.
- Marcelli M, Dayan AR, Langeneck J 2017 Finding Dory: first record of Paracanththurus hepatus (Perciformes: Acanthuridae) in the Mediterranean Sea. *Mar. Biodivers.* 47, 599–602. doi: [10.1007/s12526-016-0573-3](https://doi.org/10.1007/s12526-016-0573-3)
- Marcos M, Calafat FM, Berihuete A, Dangendorf S 2015 Long-term variations in global sea level extremes. *JGR Ocean.* 120, 8115–8134. doi: [10.1002/2015JC011173](https://doi.org/10.1002/2015JC011173)
- Marcos M, Jordà G, Gomis D, Pérez Gómez B 2011 Changes in storm surges in southern Europe from a regional model under climate change scenarios. *Glob. Planet. Change* 77, 116–128. <https://www.sciencedirect.com/science/article/pii/S0921818111000555> [Accessed September 17, 2019]
- Marcos M, Tsimplis MN 2008 Comparison of results of AOGCMs in the Mediterranean Sea during the 21st century. *JGR Ocean.* 113. doi: [10.1029/2008JC004820](https://doi.org/10.1029/2008JC004820)
- Marcos M, Tsimplis MN, Shaw AGP 2009 Sea level extremes in southern Europe. *JGR Ocean.* 114, C01007. doi: [10.1029/2008JC004912](https://doi.org/10.1029/2008JC004912)
- Mariotti A 2010 Recent changes in the Mediterranean water cycle: A pathway toward long-term regional hydroclimatic change? *J. Clim.* 23, 1513–1525. doi: [10.1175/2009JCLI3251.1](https://doi.org/10.1175/2009JCLI3251.1)
- Mariotti A, Ballabrera-Poy J, Zeng N 2005 Tropical influence on Euro-Asian autumn rainfall variability. *Clim. Dyn.* 24, 511–521.
- Mariotti A, Dell'Aquila A 2012 Decadal climate variability in the Mediterranean region: roles of large-scale forcings and regional processes. *Clim. Dyn.* 38, 1129–1145. doi: [10.1007/s00382-011-1056-7](https://doi.org/10.1007/s00382-011-1056-7)
- Mariotti A, Pan Y, Zeng N, Alessandri A 2015 Long-term climate change in the Mediterranean region in the midst of decadal variability. *Clim. Dyn.* 44, 1437–1456. doi: [10.1007/s00382-015-2487-3](https://doi.org/10.1007/s00382-015-2487-3)
- Mariotti A, Struglia MV, Zeng N, Lau K-M 2002a The Hydrological Cycle in the Mediterranean Region and Implications for the Water Budget of the Mediterranean Sea. *J. Clim.* 15, 1674–1690. doi: [10.1175/1520-0442\(2002\)015<1674:THCITM>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<1674:THCITM>2.0.CO;2)
- Mariotti A, Zeng N, Lau KM 2002b Euro-Mediterranean rainfall and ENSO-a seasonally varying relationship. *Geophys. Res. Lett.* 29, 1621. doi: [10.1029/2001GL014248](https://doi.org/10.1029/2001GL014248)
- Mariotti A, Zeng N, Yoon J-H, Artale V, Navarra A et al. 2008 Mediterranean water cycle changes: transition to drier 21st century conditions in observations and CMIP3 simulations. *Environ. Res. Lett.* 3, 044001. doi: [10.1088/1748-9326/3/4/044001](https://doi.org/10.1088/1748-9326/3/4/044001)
- Marmy A, Rajczak J, Delaloye R, Hilbich C, Hoelzle M et al. 2016 Semi-automated calibration method for modelling of mountain permafrost evolution in Switzerland. *Cryosph.* 10, 2693–2719. doi: [10.5194/tc-10-2693-2016](https://doi.org/10.5194/tc-10-2693-2016)
- Marty J-C, Chiavérini J 2010 Hydrological changes in the Ligurian Sea (NW Mediterranean, DYFAMED site) during 1995–2007 and biogeochemical consequences. *Biogeosciences* 7, 2117–2128. doi: [10.5194/bg-7-2117-2010](https://doi.org/10.5194/bg-7-2117-2010)
- Marullo S, Artale V, Santoleri R 2011 The SST Multidecadal Variability in the Atlantic–Mediterranean Region and Its Relation to AMO. *J. Clim.* 24, 4385–4401. doi: [10.1175/2011JCLI3884.1](https://doi.org/10.1175/2011JCLI3884.1)
- Marullo S, Santoleri R, Banzon V, Evans RH, Guaraccino M 2010 A diurnal-cycle resolving sea surface temperature product for the tropical Atlantic. *JGR Atmos.* 115, C05011.

- doi: [10.1029/2009JC005466](https://doi.org/10.1029/2009JC005466)
- Marvier M, Kareiva P, Neubert MG 2004 Habitat Destruction, Fragmentation, and Disturbance Promote Invasion by Habitat Generalists in a Multispecies Metapopulation. *Risk Anal.* 24, 869–878. doi: [10.1111/j.0272-4332.2004.00485.x](https://doi.org/10.1111/j.0272-4332.2004.00485.x)
- Matesanz S, Escudero A, Valladares F 2008 Additive effects of a potentially invasive grass and water stress on the performance of seedlings of gypsum specialists. *Appl. Veg. Sci.* 11, 287–296. doi: [10.3170/2008-7-18425](https://doi.org/10.3170/2008-7-18425)
- Matošević D, Pajač Živković I 2013 Alien phytophagous insect and mite species on woody plants in Croatia (in croatian). *Šumarski List*, 191–200.
- Matzrafi M 2019 Climate change exacerbates pest damage through reduced pesticide efficacy. *Pest Manag. Sci.* 75, 9–13. doi: [10.1002/ps.5121](https://doi.org/10.1002/ps.5121)
- McInnes KL, Erwin TA, Bathols JM 2011 Global Climate Model projected changes in 10 m wind speed and direction due to anthropogenic climate change. *Atmos. Sci. Lett.* 12, 325–333. doi: [10.1002/asl.341](https://doi.org/10.1002/asl.341)
- MCRIT 2015 ET2050. Territorial Scenarios and Visions for Europe. Volume 1 - Approach to Scenario Building and Storylines. https://www.espon.eu/sites/default/files/attachments/ET2050_FR-03_Volume_1_-_Approach_to_Scenario_Building_and_Storylines.pdf
- MEA 2005 *Ecosystems and human well-being: synthesis*. Island Press, Washington, DC <https://www.cifor.org/library/188/>
- Mediterranean Wetlands Observatory 2018 *Mediterranean Wetland Outlook 2: Solutions for sustainable Mediterranean Wetlands*. , eds. Geijzendorffer IR, Chazée L, Gaget E, Galewski T, Guelmami A et al. Arles, France <https://tourduvalat.org/en/actions/les-zones-humides-mediterraneennes-en-jeux-et-perspectives-2-solutions-pour-des-zones-humides-mediterraneennes-durables/>
- Meffe R, de Bustamante I 2014 Emerging organic contaminants in surface water and groundwater: A first overview of the situation in Italy. *Sci. Total Environ.* 481, 280–295. doi: [10.1016/j.scitotenv.2014.02.053](https://doi.org/10.1016/j.scitotenv.2014.02.053)
- Menchetti M, Mori E 2014 Worldwide impact of alien parrots (Aves Psittaciformes) on native biodiversity and environment: a review. *Ethol. Ecol. Evol.* 26, 172–194. doi: [10.1080/03949370.2014.905981](https://doi.org/10.1080/03949370.2014.905981)
- Menchetti M, Mori E, Angelici FM 2016 Effects of the recent world invasion by ring-necked parakeets *Psittacula krameri*, in *Problematic Wildlife*, ed. Angelici FM [Springer International Publishing], 253–266. doi: [10.1007/978-3-319-22246-2_12](https://doi.org/10.1007/978-3-319-22246-2_12)
- Mendel Z, Branco M, Battisti A 2016 Invasive Sap-Sucker Insects in the Mediterranean Basin. *Insects Dis. Mediterr. For. Syst.*, 261–291. doi: [10.1007/978-3-319-24744-1_10](https://doi.org/10.1007/978-3-319-24744-1_10)
- Menemenlis D, Fukumori I, Lee T 2007 Atlantic to Mediterranean Sea Level Difference Driven by Winds near Gibraltar Strait. *J. Phys. Oceanogr.* 37, 359–376. doi: [10.1175/JPO3015.1](https://doi.org/10.1175/JPO3015.1)
- Menendez M, García-Díez M, Fita L, Fernández J, Méndez FJ et al. 2014 High-resolution sea wind hindcasts over the Mediterranean area. *Clim. Dyn.* 42, 1857–1872. doi: [10.1007/s00382-013-1912-8](https://doi.org/10.1007/s00382-013-1912-8)
- Merhaby D, Net S, Halwani J, Ouddane B 2015 Organic pollution in surficial sediments of Tripoli harbour, Lebanon. *Mar. Pollut. Bull.* 93, 284–293. doi: [10.1016/j.marpolbul.2015.01.004](https://doi.org/10.1016/j.marpolbul.2015.01.004)
- Merhaby D, Ouddane B, Net S, Halwani J 2018 Assessment of trace metals contamination in surficial sediments along Lebanese Coastal Zone. *Mar. Pollut. Bull.* 133, 881–890. doi: [10.1016/j.marpolbul.2018.06.031](https://doi.org/10.1016/j.marpolbul.2018.06.031)
- Merkens J-L, Reimann L, Hinkel J, Vafeidis AT 2016 Gridded population projections for the coastal zone under the Shared Socioeconomic Pathways. *Glob. Planet. Change* 145, 57–66. doi: [10.1016/j.gloplacha.2016.08.009](https://doi.org/10.1016/j.gloplacha.2016.08.009)
- Meurisse N, Rassati D, Hurley BP, Brockerhoff EG, Haack RA 2019 Common pathways by which non-native forest insects move internationally and domestically. *J. Pest Sci. (2004)*, 92, 13–27. doi: [10.1007/s10340-018-0990-0](https://doi.org/10.1007/s10340-018-0990-0)
- Miglietta MM 2019 Mediterranean Tropical-Like Cyclones (Medicanes). *Atmosphere (Basel)*, 10, 206. doi: [10.3390/atmos10040206](https://doi.org/10.3390/atmos10040206)
- Miglietta MM, Laviola S, Malvaldi A, Conte D, Levizzani V et al. 2013 Analysis of tropical-like cyclones over the Mediterranean Sea through a combined modeling and satellite approach. *Geophys. Res. Lett.* 40, 2400–2405. doi: [10.1002/grl.50432](https://doi.org/10.1002/grl.50432)
- Miglietta MM, Mastrangelo E, Conte D 2015 Influence of physics parameterization schemes on the simulation of a tropical-like cyclone in the Mediterranean Sea. *Atmos. Res.* 153, 360–375. doi: [10.1016/j.atmosres.2014.09.008](https://doi.org/10.1016/j.atmosres.2014.09.008)
- Millán MM, Mantilla E, Salvador R, Carratalá A, Sanz MJ et al. 2000 Ozone Cycles in the Western Mediterranean Basin: Interpretation of Monitoring Data in Complex Coastal Terrain. *J. Appl. Meteorol.* 39, 487–508. doi: [10.1175/1520-0450\(2000\)039<0487:OCITWM>2.0.CO;2](https://doi.org/10.1175/1520-0450(2000)039<0487:OCITWM>2.0.CO;2)
- Millot C, Candela J, Fuda J-L, Tber Y 2006 Large warming and salinification of the Mediterranean outflow due to changes in its composition. *Deep Sea Res. Part I Oceanogr. Res. Pap.* 53, 656–666. doi: [10.1016/J.DSR.2005.12.017](https://doi.org/10.1016/J.DSR.2005.12.017)
- Mitsakou C, Kallos G, Papantoniou N, Spyrou C, Solomos S et al. 2008 Saharan dust levels in Greece and received inhalation doses. *Atmos. Chem. Phys.* 8, 7181–7192. doi: [10.5194/acp-8-7181-2008](https://doi.org/10.5194/acp-8-7181-2008)
- Moemken J, Reyers M, Feldmann H, Pinto JG 2018 Future Changes of Wind Speed and Wind Energy Potentials in EURO-CORDEX Ensemble Simulations. *JGR Atmos.* 123, 6373–6389. doi: [10.1029/2018JD028473](https://doi.org/10.1029/2018JD028473)
- Mohamed AH, Squires VR 2018 Drylands of the Mediterranean Basin: Challenges, Problems and Prospects, in *Climate Variability Impacts on Land Use and Livelihoods in Drylands*, eds. Gaur M, Squires V [Springer, Cham].
- Molle F, Tanouti O 2017 Squaring the circle: Agricultural intensification vs. Water conservation in Morocco. *Agric. Water Manag.* 192, 170–179. doi: [10.1016/j.agwat.2017.07.009](https://doi.org/10.1016/j.agwat.2017.07.009)
- Monceau K, Bonnard O, Thiéry D 2014 *Vespa velutina*: a new invasive predator of honeybees in Europe. *J. Pest Sci. (2004)*, 87, 1–16. doi: [10.1007/s10340-013-0537-3](https://doi.org/10.1007/s10340-013-0537-3)
- Moncer M, Hamza A, Feki-Sahnoun W, Mabrouk L, Hassen MB 2017 Variability patterns of epibenthic microalgae in eastern Tunisian coasts. *Sci. Mar.* 81, 487–498. doi: [10.3989/scimar.04651.17A](https://doi.org/10.3989/scimar.04651.17A)
- Montecchio L, Fanchin G, Simonato M, Faccoli M 2014 First

- record of thousand cankers disease fungal pathogen *Geosmithia morbida* and Walnut Twig Beetle *Pityophthorus juglandis* on *Juglans regia* in Europe. *Plant Dis.* 98, 1445. doi: [10.1094/pdis-07-14-0719-pdn](https://doi.org/10.1094/pdis-07-14-0719-pdn)
- Monteiro A, Russo M, Gama C, Borrego C 2018a How important are maritime emissions for the air quality: At European and national scale. *Environ. Pollut.* 242, 565–575. doi: [10.1016/j.envpol.2018.07.011](https://doi.org/10.1016/j.envpol.2018.07.011)
- Monteiro A, Sá E, Fernandes AP, Gama C, Sorte S et al. 2018b How healthy will be the air quality in 2050? *Air Qual. Atmos. Heal.* 11, 353–362. doi: [10.1007/s11869-017-0466-z](https://doi.org/10.1007/s11869-017-0466-z)
- Mooney HA, Hobbs RJ 2000 *Invasive Species in a Changing World*. Washington DC, USA: Island Press
- Moreira F, Viedma O, Arianoutsou M, Curt T, Koutsias N et al. 2011 Landscape – wildfire interactions in southern Europe: Implications for landscape management. *J. Environ. Manage.* 92, 2389–2402. doi: [10.1016/j.jenvman.2011.06.028](https://doi.org/10.1016/j.jenvman.2011.06.028)
- Moreno-González R, Rodríguez-Mozaz S, Gros M, Barceló D, León VM 2015 Seasonal distribution of pharmaceuticals in marine water and sediment from a mediterranean coastal lagoon (SE Spain). *Environ. Res.* 138, 326–344. doi: [10.1016/j.envres.2015.02.016](https://doi.org/10.1016/j.envres.2015.02.016)
- Moreno J, Arianoutsou M, González-Cabán A, Mouillot F, Oechel W et al. 2014 Forest fires under climate, social and economic changes in Europe, the Mediterranean and other fire-affected areas of the world.
- Morce CP, Kennedy JJ, Rayner NA, Jones PD 2012 Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: The HadCRUT4 data set. *JGR Atmos.* 117, D08101. doi: [10.1029/2011JD017187](https://doi.org/10.1029/2011JD017187)
- Moriondo M, Trombi G, Ferrise R, Brandani G, Dibari C et al. 2013 Olive trees as bio-indicators of climate evolution in the Mediterranean Basin. *Glob. Ecol. Biogeogr.* 22. doi: [10.1111/geb.12061](https://doi.org/10.1111/geb.12061)
- Moullec F, Barrier N, Drira S, Guilhaumon F, Marsaleix P et al. 2019 An End-to-End Model Reveals Losers and Winners in a Warming Mediterranean Sea. *Front. Mar. Sci.* 6. doi: [10.3389/fmars.2019.00345](https://doi.org/10.3389/fmars.2019.00345)
- Müller R-J, Kleeberg I, Deckwer W-D 2001 Biodegradation of polyesters containing aromatic constituents. *J. Biotechnol.* 86, 87–95. doi: [10.1016/S0168-1656\(00\)00407-7](https://doi.org/10.1016/S0168-1656(00)00407-7)
- Murakami D, Yamagata Y 2019 Estimation of Gridded Population and GDP Scenarios with Spatially Explicit Statistical Downscaling. *Sustainability* 11, 2106. doi: [10.3390/su11072106](https://doi.org/10.3390/su11072106)
- Mutinelli F, Montarsi F, Federico G, Granato A, Ponti AM et al. 2014 Detection of *Aethina tumida* Murray (Coleoptera: Nitidulidae) in Italy: outbreaks and early reaction measures. *J. Apic. Res.* 53, 569–575. doi: [10.3896/ibra.1.53.5.13](https://doi.org/10.3896/ibra.1.53.5.13)
- Mutke S, Calama R, Neaymeh EN, Roques A 2016 Impact of the dry cone syndrome on commercial kernel yield of stone pine cones. *Options Méditerranéennes* 122, 78–84.
- Nabat P, Somot S, Cassou C, Mallet M, Michou M et al. 2020 Modulation of radiative aerosols effects by atmospheric circulation over the Euro-Mediterranean region. *Atmos. Chem. Phys.* 20, 8315–8349.
- Nabat P, Somot S, Mallet M, Chiapello I, Morcrette JJ et al. 2013 A 4-D climatology (1979–2009) of the monthly tropospheric aerosol optical depth distribution over the Mediterranean region from a comparative evaluation and blending of remote sensing and model products. *Atmos. Meas. Tech.* 6. doi: [10.5194/amt-6-1287-2013](https://doi.org/10.5194/amt-6-1287-2013)
- Nabat P, Somot S, Mallet M, Michou M, Sevault F et al. 2015a Dust aerosol radiative effects during summer 2012 simulated with a coupled regional aerosol–atmosphere–ocean model over the Mediterranean. *Atmos. Chem. Phys.* 15, 3303–3326. doi: [10.5194/acp-15-3303-2015](https://doi.org/10.5194/acp-15-3303-2015)
- Nabat P, Somot S, Mallet M, Sánchez-Lorenzo A, Wild M 2014 Contribution of anthropogenic sulfate aerosols to the changing Euro-Mediterranean climate since 1980. *Geophys. Res. Lett.* 41, 5605–5611. doi: [10.1002/2014GL060798](https://doi.org/10.1002/2014GL060798)
- Nabat P, Somot S, Mallet M, Sevault F, Chiacchio M et al. 2015b Direct and semi-direct aerosol radiative effect on the Mediterranean climate variability using a coupled regional climate system model. *Clim. Dyn.* 44, 1127–1155. doi: [10.1007/s00382-014-2205-6](https://doi.org/10.1007/s00382-014-2205-6)
- Naidja L, Ali-Khodja H, Khardi S 2018 Sources and levels of particulate matter in North African and Sub-Saharan cities: a literature review. *Environ. Sci. Pollut. Res.* 25, 12303–12328. doi: [10.1007/s11356-018-1715-x](https://doi.org/10.1007/s11356-018-1715-x)
- Nakićenović N 2000 *Special report on emissions scenarios: A special report of Working Group III of the Intergovernmental Panel on Climate Change*. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press
- Naranjo C, García-Lafuente J, Sammartino S, Sánchez-Garrido JC, Sánchez-Leal R et al. 2017 Recent changes (2004–2016) of temperature and salinity in the Mediterranean outflow. *Geophys. Res. Lett.* 44, 5665–5672. doi: [10.1002/2017GL072615](https://doi.org/10.1002/2017GL072615)
- Nashwan MS, Shahid S, Abd Rahim N 2018 Unidirectional trends in annual and seasonal climate and extremes in Egypt. *Theor. Appl. Climatol.* 136, 457–473. doi: [10.1007/s00704-018-2498-1](https://doi.org/10.1007/s00704-018-2498-1)
- Nasrallah HA, Balling RC 1993 Spatial and temporal analysis of Middle Eastern temperature changes. *Clim. Change* 25, 153–161. doi: [10.1007/bf01661203](https://doi.org/10.1007/bf01661203)
- Nastos PT, Karavana Papadimou K, Matsangouras IT 2018 Mediterranean tropical-like cyclones: Impacts and composite daily means and anomalies of synoptic patterns. *Atmos. Res.* 208, 156–166. doi: [10.1016/J.ATMOSRES.2017.10.023](https://doi.org/10.1016/J.ATMOSRES.2017.10.023)
- Naumann G, Alfieri L, Wyser K, Mentaschi L, Betts RA et al. 2018 Global changes in drought conditions under different levels of warming. *Geophys. Res. Lett.* 45, 3285–3296. doi: [10.1002/2017GL076521](https://doi.org/10.1002/2017GL076521)
- Navarro-Pedreño J, Gómez I, Almendro-Candel MB, Meléndez-Pastor I 2008 Heavy metals in Mediterranean soils, in *Soil contamination research trends* (Nova Science Publishers Inc), 161–176. https://www.novapublishers.com/catalog/product_info.php?products_id=6860
- Naves P, Bonifácio L, de Sousa E 2016 The Pine Wood Nematode and Its Local Vectors in the Mediterranean Basin. *Insects Dis. Mediterr. For. Syst.*, 329–378. doi: [10.1007/978-3-319-24744-1_12](https://doi.org/10.1007/978-3-319-24744-1_12)
- Nehme N, Haydar C, Koubaissy B, Fakih M, Awad S et al. 2014 The distribution of heavy metals in the Lower River Basin,

- Lebanon. *Phys. Procedia* 55, 456–463.
doi: [10.1016/j.phpro.2014.07.066](https://doi.org/10.1016/j.phpro.2014.07.066)
- Nelson GC, Bennett EM, Berhe AA, Cassman KG, DeFries RS et al. 2006 Anthropogenic drivers of ecosystem change: An overview. *Ecol. Soc.* 11, 29. doi: [10.5751/ES-01826-110229](https://doi.org/10.5751/ES-01826-110229)
- Nepstad DC, Verssimo A, Alencar A, Nobre CA, Lima E et al. 1999 Large-scale impoverishment of Amazonian forests by logging and fire. *Nature* 398, 505–508. doi: [10.1038/19066](https://doi.org/10.1038/19066)
- Nieto R, Gimeno L, Trigo RM 2006 A Lagrangian identification of major sources of Sahel moisture. *Geophys. Res. Lett.* 33. doi: [10.1029/2006GL027232](https://doi.org/10.1029/2006GL027232)
- Nikulin G, Lennard C, Dosio A, Kjellström E, Chen Y et al. 2018 The effects of 1.5 and 2 degrees of global warming on Africa in the CORDEX ensemble. *Environ. Res. Lett.* 13, 65003. doi: [10.1088/1748-9326/aab1b1](https://doi.org/10.1088/1748-9326/aab1b1)
- Nissen KM, Leckebusch GC, Pinto JG, Renggli D, Ulbrich S et al. 2010 Cyclones causing wind storms in the Mediterranean: characteristics, trends and links to large-scale patterns. *Nat. Hazards Earth Syst. Sci.* 10, 1379–1391. doi: [10.5194/nhess-10-1379-2010](https://doi.org/10.5194/nhess-10-1379-2010)
- Nissen KM, Leckebusch GC, Pinto JG, Ulbrich U 2014 Mediterranean cyclones and windstorms in a changing climate. *Reg. Environ. Chang.* 14, 1873–1890. doi: [10.1007/s10113-012-0400-8](https://doi.org/10.1007/s10113-012-0400-8)
- Nödler K, Voutsas D, Licha T 2014 Polar organic micropollutants in the coastal environment of different marine systems. *Mar. Pollut. Bull.* 85, 50–59. doi: [10.1016/j.marpolbul.2014.06.024](https://doi.org/10.1016/j.marpolbul.2014.06.024)
- Nori M 2018 Migration, agriculture and rural territories in the Mediterranean, in *Migration and Inclusive Rural Development in the Mediterranean*, ed. MediTERRA C/ [Presses de Sciences Po].
- Novoa A, Le Roux JJ, Robertson MP, Wilson JRU, Richardson DM 2015 Introduced and invasive cactus species: a global review. *AoB Plants* 7. doi: [10.1093/aobpla/plu078](https://doi.org/10.1093/aobpla/plu078)
- Noyes PD, McElwee MK, Miller HD, Clark BW, Van Tiem LA et al. 2009 The toxicology of climate change: Environmental contaminants in a warming world. *Environ. Int.* 35, 971–986. doi: [10.1016/j.envint.2009.02.006](https://doi.org/10.1016/j.envint.2009.02.006)
- Nunes L, Gower ST, Peckham SD, Magalhães M, Lopes D et al. 2015 Estimation of productivity in pine and oak forests in northern Portugal using Biome-BGC. *For. An Int. J. For. Res.* 88, 200–212. doi: [10.1093/forestry/cpu044](https://doi.org/10.1093/forestry/cpu044)
- Nyngi W, Oguge N, Dziba L, Chandipo R, Didier TA et al. 2018 Direct and indirect drivers of change in biodiversity and nature's contributions to people, in *IPBES (2018): The IPBES regional assessment report on biodiversity and ecosystem services for Africa*, eds. Archer E, Dziba L, Mulongoy KJ, Maoela MA, Walters M (Bonn, Germany: Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services), 207–296.
- Nykjaer L 2009 Mediterranean Sea surface warming 1985–2006. *Clim. Res.* 39, 11–17. doi: [10.3354/cr00794](https://doi.org/10.3354/cr00794)
- O'Neill BC, Kriegler E, Ebi KL, Kemp-Benedict E, Riahi K et al. 2017 The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century. *Glob. Environ. Chang.* 42, 169–180. doi: [10.1016/j.gloenvcha.2015.01.004](https://doi.org/10.1016/j.gloenvcha.2015.01.004)
- O'Neill BC, Kriegler E, Riahi K, Ebi KL, Hallegatte S et al. 2014 A new scenario framework for climate change research: The concept of shared socioeconomic pathways. *Clim. Change* 122, 387–400. doi: [10.1007/s10584-013-0905-2](https://doi.org/10.1007/s10584-013-0905-2)
- Obermann-Hellund A, Conte D, Somot S, Torma CZ, Ahrens B 2018 Mistral and Tramontane wind systems in climate simulations from 1950 to 2100. *Clim. Dyn.* 50, 693–703. doi: [10.1007/s00382-017-3635-8](https://doi.org/10.1007/s00382-017-3635-8)
- Obermann A, Bastin S, Belamari S, Conte D, Gaertner MÁ et al. 2018 Mistral and Tramontane wind speed and wind direction patterns in regional climate simulations. *Clim. Dyn.* 51, 1059–1076. doi: [10.1007/s00382-016-3053-3](https://doi.org/10.1007/s00382-016-3053-3)
- Occhipinti-Ambrogi A 2007 Global change and marine communities: Alien species and climate change. *Mar. Pollut. Bull.* 55, 342–352. doi: [10.1016/j.marpolbul.2006.11.014](https://doi.org/10.1016/j.marpolbul.2006.11.014)
- Ochoa-Hueso R, Bell MD, Manrique E 2014 Impacts of increased nitrogen deposition and altered precipitation regimes on soil fertility and functioning in semiarid Mediterranean shrublands. *J. Arid Environ.* 104, 106–115. doi: [10.1016/j.jaridenv.2014.01.020](https://doi.org/10.1016/j.jaridenv.2014.01.020)
- Ochoa-Hueso R, Munzi S, Alonso R, Arróniz-Crespo M, Avila A et al. 2017 Ecological impacts of atmospheric pollution and interactions with climate change in terrestrial ecosystems of the Mediterranean Basin: Current research and future directions. Elsevier <https://www.sciencedirect.com/science/article/pii/S0269749116320760?via%3Dihub> [Accessed March 30, 2019]
- Olenycz M, Sokólski A, Niewińska A, Wołowicz M, Namieśnik J et al. 2015 Comparison of PCBs and PAHs levels in European coastal waters using mussels from the *Mytilus edulis* complex as biomonitor. *Oceanologia* 57, 196–211. doi: [10.1016/j.oceano.2014.12.001](https://doi.org/10.1016/j.oceano.2014.12.001)
- Oliva M, Gómez-Ortiz A, Salvador-Franch F, Salvà-Catarineu M, Palacios D et al. 2016 Inexistence of permafrost at the top of the Veleta peak (Sierra Nevada, Spain). *Sci. Total Environ.* 550, 484–494. doi: [10.1016/j.scitotenv.2016.01.150](https://doi.org/10.1016/j.scitotenv.2016.01.150)
- Oliva M, Žebre M, Guglielmin M, Hughes PD, Činer A et al. 2018 Permafrost conditions in the Mediterranean region since the Last Glaciation. *Earth-Science Rev.* 185, 397–436. doi: [10.1016/j.earscirev.2018.06.018](https://doi.org/10.1016/j.earscirev.2018.06.018)
- Oliveira MA, Tomlinson SJ, Carnell EJ, Dore AJ, Serrano HC et al. 2020 Nitrogen and sulfur deposition over a region in SW Europe based on a regional atmospheric chemical transport model. *Atmos. Environ.* 223, 117290. doi: [10.1016/j.atmosenv.2020.117290](https://doi.org/10.1016/j.atmosenv.2020.117290)
- Oliver ECJ, Donat MG, Burrows MT, Moore PJ, Smale DA et al. 2018 Longer and more frequent marine heatwaves over the past century. *Nat. Commun.* 9, 1324. doi: [10.1038/s41467-018-03732-9](https://doi.org/10.1038/s41467-018-03732-9)
- Onea F, Deleanu L, Rusu L, Georgescu C 2016 Evaluation of the wind energy potential along the Mediterranean Sea coasts. *Energy Explor. Exploit.* 34, 766–792. doi: [10.1177/0144598716659592](https://doi.org/10.1177/0144598716659592)
- Onorati G, di Meo T, Bussettini M, Fabiani C, Farrace MGG et al. 2006 Groundwater quality monitoring in Italy for the implementation of the EU water framework directive. *Phys. Chem. Earth, Parts A/B/C* 31, 1004–1014. doi: [10.1016/j.pce.2006.07.001](https://doi.org/10.1016/j.pce.2006.07.001)

- Oppenheimer M, Glavovic BC, Hinkel J, van de Wal R, Magnan AK et al. 2019 Sea Level Rise and Implications for Low-Lying Islands, Coasts and Communities, in *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, eds. Pörtner H-O, Roberts DC, Masson-Delmotte V, Zhai P, Tignor M et al. (in press).
- Osborn TJ, Briffa KR, Tett SFB, Jones PD, Trigo RM 1999 Evaluation of the North Atlantic Oscillation as simulated by a coupled climate model. *Clim. Dyn.* 15, 685–702. doi: [10.1007/s003820050310](https://doi.org/10.1007/s003820050310)
- Osio GC, Orio A, Millar CP 2015 Assessing the vulnerability of Mediterranean demersal stocks and predicting exploitation status of un-assessed stocks. *Fish. Res.* 171, 110–121. doi: [10.1016/j.fishres.2015.02.005](https://doi.org/10.1016/j.fishres.2015.02.005)
- Ouzeau G, Soubeiryroux J-M, Schneider M, Vautard R, Planton S 2016 Heat waves analysis over France in present and future climate: Application of a new method on the EURO-CORDEX ensemble. *Clim. Serv.* 4, 1–12. doi: [10.1016/j.cisler.2016.09.002](https://doi.org/10.1016/j.cisler.2016.09.002)
- Ozer T, Gertman I, Kress N, Silverman J, Herut B 2017 Interannual thermohaline (1979–2014) and nutrient (2002–2014) dynamics in the Levantine surface and intermediate water masses, SE Mediterranean Sea. *Glob. Planet. Change* 151, 60–67. doi: [10.1016/J.GLOPLACHA.2016.04.001](https://doi.org/10.1016/J.GLOPLACHA.2016.04.001)
- Ozturk T, Ceber ZP, Türkeş M, Kurnaz ML 2015 Projections of climate change in the Mediterranean Basin by using down-scaled global climate model outputs. *Int. J. Climatol.* 35, 4276–4292. doi: [10.1002/joc.4285](https://doi.org/10.1002/joc.4285)
- Paelrl HW, Paul VJ 2012 Climate change: Links to global expansion of harmful cyanobacteria. *Water Res.* 46, 1349–1363. doi: [10.1016/j.watres.2011.08.002](https://doi.org/10.1016/j.watres.2011.08.002)
- Palanques A, Lopez L, Guillén J, Puig P, Masqué P 2017 Decline of trace metal pollution in the bottom sediments of the Barcelona City continental shelf (NW Mediterranean). *Sci. Total Environ.* 579, 755–767. doi: [10.1016/j.scitotenv.2016.11.031](https://doi.org/10.1016/j.scitotenv.2016.11.031)
- Palmiéri J, Orr JC, Dutay J-C, Béranger K, Schneider A et al. 2015 Simulated anthropogenic CO₂ storage and acidification of the Mediterranean Sea. *Biogeosciences* 12, 781–802. doi: [10.5194/bg-12-781-2015](https://doi.org/10.5194/bg-12-781-2015)
- Paluselli A, Aminot Y, Galgani F, Net S, Sempéré R 2018a Occurrence of phthalate acid esters (PAEs) in the northwestern Mediterranean Sea and the Rhône River. *Prog. Oceanogr.* 163, 221–231. doi: [10.1016/j.pocean.2017.06.002](https://doi.org/10.1016/j.pocean.2017.06.002)
- Paluselli A, Fauvette V, Schmidt N, Galgani F, Net S et al. 2018b Distribution of phthalates in Marseille Bay (NW Mediterranean Sea). *Sci. Total Environ.* 621, 578–587. doi: [10.1016/j.scitotenv.2017.11.306](https://doi.org/10.1016/j.scitotenv.2017.11.306)
- Pant P, Harrison RM 2013 Estimation of the contribution of road traffic emissions to particulate matter concentrations from field measurements: A review. *Atmos. Environ.* 77, 78–97. doi: [10.1016/J.ATMOSENV.2013.04.028](https://doi.org/10.1016/J.ATMOSENV.2013.04.028)
- Papaconstantinou C, Farrugio H 2000 Fisheries in the Mediterranean. *Mediterr. Mar. Sci.* 1, 5. doi: [10.12681/mms.2](https://doi.org/10.12681/mms.2)
- Papanastasis VP 2004 Vegetation degradation and land use changes in agrosilvopastoral systems, in *Sustainability of Agrosilvopastoral Systems*, eds. Schnabel S, Ferreira A (Reiskirchen, Germany: Catena Verlag), 1–12.
- Papanastasis VP, Kazaklis A 1998 Land use changes and conflicts in the Mediterranean-Type ecosystems of Western Crete. *Ecol. Stud.*, 141–154. doi: [10.1007/978-3-662-03543-6_8](https://doi.org/10.1007/978-3-662-03543-6_8)
- Parravicini V, Mangialajo L, Mousseau L, Peirano A, Morri C et al. 2015 Climate change and warm-water species at the north-western boundary of the Mediterranean Sea. *Mar. Ecol.* 36, 897–909. doi: [10.1111/maec.12277](https://doi.org/10.1111/maec.12277)
- Pascual A, Vidal-Vijande E, Ruiz S, Somot S, Papadopoulos V 2014 Spatio-temporal variability of the surface circulation in the western Mediterranean: A comparative study using altimetry and modeling, in *The Mediterranean Sea: Temporal Variability and Spatial Patterns*, eds. Borzelli GLE, Gacic M, Lionello P, Malanotte-Rizzoli P, 5–24. <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/9781118847572#page=13>
- Pastor F, Valiente JA, Palau JL 2018 Sea surface temperature in the Mediterranean: trends and spatial patterns (1982–2016). *Pure Appl. Geophys.* 175, 4017–4029. doi: [10.1007/s00024-017-1739-z](https://doi.org/10.1007/s00024-017-1739-z)
- Pauly D, Zeller D 2016 Catch reconstructions reveal that global marine fisheries catches are higher than reported and declining. *Nat. Commun.* 7, 10244. doi: [10.1038/ncomms10244](https://doi.org/10.1038/ncomms10244)
- Pausas JG, Fernández-Muñoz S 2012 Fire regime changes in the Western Mediterranean Basin: from fuel-limited to drought-driven fire regime. *Clim. Change* 110, 215–226. doi: [10.1007/s10584-011-0060-6](https://doi.org/10.1007/s10584-011-0060-6)
- Pellet V, Aires F, Munier S, Fernández Prieto D, Jordà G et al. 2019 Integrating multiple satellite observations into a coherent dataset to monitor the full water cycle – application to the Mediterranean region. *Hydrol. Earth Syst. Sci.* 23, 465–491. doi: [10.5194/hess-23-465-2019](https://doi.org/10.5194/hess-23-465-2019)
- Peña-Angulo D, Vicente-Serrano SM, Domínguez-Castro F, Murphy C, Reig F et al. 2020 Long-term precipitation in Southwestern Europe reveals no clear trend attributable to anthropogenic forcing. *Environ. Res. Lett.* 15, 094070.
- Penna A, Bertozzini E, Battocchi C, Galluzzi L, Giacobbe MG et al. 2007 Monitoring of HAB species in the Mediterranean Sea through molecular methods. *J. Plankton Res.* 29, 19–38. doi: [10.1093/plankt/fbl053](https://doi.org/10.1093/plankt/fbl053)
- Peñuelas J, Canadell JG, Ogaya R 2011 Increased water-use efficiency during the 20th century did not translate into enhanced tree growth. *Glob. Ecol. Biogeogr.* 20, 597–608. doi: [10.1111/j.1466-8238.2010.00608.x](https://doi.org/10.1111/j.1466-8238.2010.00608.x)
- Perez J, Menendez M, Camus P, Mendez F, Fernando J, Losada IJ 2015 Statistical multi-model climate projections of surface ocean waves in Europe. *Ocean Model.* 96, 161–170. doi: [10.1016/j.ocemod.2015.06.001](https://doi.org/10.1016/j.ocemod.2015.06.001)
- Perović V, Jakšić D, Jaramaz D, Koković N, Čakmak D et al. 2018 Spatio-temporal analysis of land use/land cover change and its effects on soil erosion (Case study in the Oplenac wine-producing area, Serbia). *Environ. Monit. Assess.* 190, doi: [10.1007/s10661-018-7025-4](https://doi.org/10.1007/s10661-018-7025-4)
- Perry AL, Low PJ, Ellis JR, Reynolds JD 2005 Climate Change and Distribution Shifts in Marine Fishes. *Science (80-.).* 308, 1912–1915. doi: [10.1126/science.1111322](https://doi.org/10.1126/science.1111322)
- Petanidou T, Kizos T, Soulakellis N 2008 Socioeconomic Di-

- mensions of Changes in the Agricultural Landscape of the Mediterranean Basin: A Case Study of the Abandonment of Cultivation Terraces on Nisyros Island, Greece. *Environ. Manage.* 41, 250–266. doi: [10.1007/s00267-007-9054-6](https://doi.org/10.1007/s00267-007-9054-6)
- Peters K, Breitsameter L, Gerowitt B 2014 Impact of climate change on weeds in agriculture: a review. *Agron. Sustain. Dev.* 34, 707–721. doi: [10.1007/s13593-014-0245-2](https://doi.org/10.1007/s13593-014-0245-2)
- Petterssen S 1956 *Weather Analysis and Forecasting*. McGraw-Hill
- Pey J, Querol X, Alastuey A, Forastiere F, Stafoggia M 2013 African dust outbreaks over the Mediterranean Basin during 2001–2011: PM10 concentrations, phenomenology and trends, and its relation with synoptic and mesoscale meteorology. *Atmos. Chem. Phys.* 13, 1395–1410. doi: [10.5194/acp-13-1395-2013](https://doi.org/10.5194/acp-13-1395-2013)
- Pfeifroth U, Sánchez-Lorenzo A, Manara V, Trentmann J, Hollmann R 2018 Trends and variability of surface solar radiation in Europe based on surface- and satellite-based data records. *JGR Atmos.* 123, 1735–1754. doi: [10.1002/2017JD027418](https://doi.org/10.1002/2017JD027418)
- Pham MK, Chamizo E, Mas Balbuena JL, Miquel J-CC, Martín J et al. 2017 Impact of Saharan dust events on radionuclide levels in Monaco air and in the water column of the northwest Mediterranean Sea. *J. Environ. Radioact.* 166, 2–9. doi: [10.1016/j.jenvrad.2016.04.014](https://doi.org/10.1016/j.jenvrad.2016.04.014)
- Phelps LN, Kaplan JO 2017 Land use for animal production in global change studies: Defining and characterizing a framework. *Glob. Chang. Biol.* 23, 4457–4471. doi: [10.1111/gcb.13732](https://doi.org/10.1111/gcb.13732)
- Piazzi L, Balata D, Bulleri F, Gennaro P, Ceccherelli G 2016 The invasion of *Caulerpa cylindracea* in the Mediterranean: the known, the unknown and the knowable. *Mar. Biol.* 163. doi: [10.1007/s00227-016-2937-4](https://doi.org/10.1007/s00227-016-2937-4)
- Piazzi L, Ceccherelli G 2006 Persistence of biological invasion effects: Recovery of macroalgal assemblages after removal of *Caulerpa racemosa* var. *cylindracea*. *Estuar. Coast. Shelf Sci.* 68, 455–461. doi: [10.1016/j.ecss.2006.02.011](https://doi.org/10.1016/j.ecss.2006.02.011)
- Piazzi L, Meinesz A, Verlaque M, Akçali B, Antolic B et al. 2005 Invasion of *Caulerpa racemosa* var. *cylindracea* (Caulerpales, Chlorophyta) in the Mediterranean Sea: an assessment of the spread. *Cryptogam. Algol.* 26, 189–202.
- Pikitch EK, Santora C, Babcock EA, Bakun A, Bonfil R et al. 2004 Ecosystem-Based Fishery Management. *Science* (80-.). 305, 346–347. doi: [10.1126/science.1098222](https://doi.org/10.1126/science.1098222)
- Pinilla V, Ayuda MI, Saez LA 2008 Rural depopulation and the migration turnaround in Mediterranean Western Europe: a case study of Aragon. *J. Rural Community Dev.* 3.
- Pinnegar JK, Polunin NVC, Badalamenti F 2003 Long-term changes in the trophic level of western Mediterranean fishery and aquaculture landings. *Can. J. Fish. Aquat. Sci.* 60, 222–235. doi: [10.1139/f03-016](https://doi.org/10.1139/f03-016)
- Pinto-Correia T, Vos W 2004 Multifunctionality in Mediterranean landscapes - past and future. *New Dimens. Eur. Landsc.*, 135–164. doi: [10.1007/978-1-4020-2911-0_10](https://doi.org/10.1007/978-1-4020-2911-0_10)
- Pinto JG, Raible CC 2012 Past and recent changes in the North Atlantic oscillation. *Wiley Interdiscip. Rev. Clim. Chang.* 3, 79–90. doi: [10.1002/wcc.150](https://doi.org/10.1002/wcc.150)
- Pinto JG, Ulbrich U, Leckebusch GC, Spangehl T, Reyers M et al. 2007 Changes in storm track and cyclone activity in three SRES ensemble experiments with the ECHAM5/MPI-OM1 GCM. *Clim. Dyn.* 29, 195–210. doi: [10.1007/s00382-007-0230-4](https://doi.org/10.1007/s00382-007-0230-4)
- Pirazzoli PA, Tomasin A 2003 Recent near-surface wind changes in the central Mediterranean and Adriatic areas. *Int. J. Climatol.* 23, 963–973. doi: [10.1002/joc.925](https://doi.org/10.1002/joc.925)
- Pisano A, Nardelli BB, Tronconi C, Santoleri R 2016 The new Mediterranean optimally interpolated pathfinder AVHRR SST data set. *Remote Sens. Environ.* 176, 107–116. doi: [10.1016/j.rse.2016.01.019](https://doi.org/10.1016/j.rse.2016.01.019)
- Pisanu S, Farris E, Filigheddu R, García MB 2012 Demographic effects of large, introduced herbivores on a long-lived endemic plant. *Plant Ecol.* 213, 1543–1553. doi: [10.1007/s11258-012-0110-9](https://doi.org/10.1007/s11258-012-0110-9)
- Planton S, Lionello P, Artale V, Aznar R, Carrillo A et al. 2012 The climate of the Mediterranean region in future climate projections, in *The Climate of the Mediterranean Region: From the Past to the Future*, ed. Lionello P (Oxford: Elsevier), 449–502. doi: [10.1016/B978-0-12-416042-2.00008-2](https://doi.org/10.1016/B978-0-12-416042-2.00008-2)
- Poli JX 1791 Testacea utriusque Siciliae eorumque historia et anatomie tabulis aeneis illustrata. *Ex Reg. Typogr. Parmae* 1, i–lxiii, [1–6].
- Pomaro A, Cavalieri L, Lionello P 2017 Climatology and trends of the Adriatic Sea wind waves: Analysis of a 37-year long instrumental data set. *Int. J. Climatol.* 37, 4237–4250. doi: [10.1002/joc.5066](https://doi.org/10.1002/joc.5066)
- Por FD 1978 *Lessepsian migration: the influx of Red Sea biota into the Mediterranean by way of the Suez Canal*. Springer-Verlag, Berlin doi: [10.1016/0160-9327\(79\)90119-4](https://doi.org/10.1016/0160-9327(79)90119-4)
- Povinec PP, Holý K, Chudý M, Šivo A, Sýkora I et al. 2012 Long-term variations of ^{14}C and ^{137}Cs in the Bratislava air – implications of different atmospheric transport processes. *J. Environ. Radioact.* 108, 33–40. doi: [10.1016/J.JENVRAD.2011.08.004](https://doi.org/10.1016/J.JENVRAD.2011.08.004)
- Pozo-Vázquez D, Esteban-Parra MJ, Rodrigo FS, Castro-Díez Y 2001 The Association between ENSO and Winter Atmospheric Circulation and Temperature in the North Atlantic Region. *J. Clim.* 14, 3408–3420. doi: [10.1175/1520-0442\(2001\)014<3408:tabeaw>2.0.co;2](https://doi.org/10.1175/1520-0442(2001)014<3408:tabeaw>2.0.co;2)
- Prein AF, Gobiet A 2017 Impacts of uncertainties in European gridded precipitation observations on regional climate analysis. *Int. J. Climatol.* 37, 305–327. doi: [10.1002/joc.4706](https://doi.org/10.1002/joc.4706)
- Price C, Michaelides S, Pashiardis S, Alpert P 1999 Long term changes in diurnal temperature range in Cyprus. *Atmos. Res.* 51, 85–98. doi: [10.1016/S0169-8095\(99\)00022-8](https://doi.org/10.1016/S0169-8095(99)00022-8)
- Proia L, Lupini G, Osorio V, Pérez S, Barceló D et al. 2013 Response of biofilm bacterial communities to antibiotic pollutants in a Mediterranean river. *Chemosphere* 92, 1126–1135. doi: [10.1016/j.chemosphere.2013.01.063](https://doi.org/10.1016/j.chemosphere.2013.01.063)
- Puigdefábregas J, Mendizábal T 1998 Perspectives on desertification: western Mediterranean. *J. Arid Environ.* 39, 209–224. doi: [10.1006/jare.1998.0401](https://doi.org/10.1006/jare.1998.0401)
- Pujo-Pay M, Conan P, Oriol L, Cornet-Barthaux V, Falco C et al. 2011 Integrated survey of elemental stoichiometry (C, N, P) from the western to eastern Mediterranean Sea. *Biogeosciences* 8, 883–899. doi: [10.5194/bg-8-883-2011](https://doi.org/10.5194/bg-8-883-2011)

- Putaud J-P, Van Dingenen R, Alastuey A, Bauer H, Birmili W et al. 2010 A European aerosol phenomenology – 3: Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe. *Atmos. Environ.* 44, 1308–1320. doi: [10.1016/j.atmosenv.2009.12.011](https://doi.org/10.1016/j.atmosenv.2009.12.011)
- Pyrina M, Hatzianastassiou N, Matsoukas C, Fotiadi A, Papadimas CD et al. 2015 Cloud effects on the solar and thermal radiation budgets of the Mediterranean basin. *Atmos. Res.* 152, 14–28. doi: [10.1016/j.atmosres.2013.11.009](https://doi.org/10.1016/j.atmosres.2013.11.009)
- Quadrelli R, Pavan V, Molteni F 2001 Wintertime variability of Mediterranean precipitation and its links with large-scale circulation anomalies. *Clim. Dyn.* 17, 457–466. doi: [10.1007/s003820000121](https://doi.org/10.1007/s003820000121)
- Quan X-W, Hoerling MP, Perlitz J, Diaz HF 2018 On the Time of Emergence of Tropical Width Change. *J. Clim.* 31, 7225–7236. doi: [10.1175/jcli-d-18-0068.1](https://doi.org/10.1175/jcli-d-18-0068.1)
- Querol X, Alastuey A, Gangoiti G, Perez N, Lee HK et al. 2018 Phenomenology of summer ozone episodes over the Madrid Metropolitan Area, central Spain. *Atmos. Chem. Phys.* 18, 6511–6533. doi: [10.5194/acp-18-6511-2018](https://doi.org/10.5194/acp-18-6511-2018)
- Querol X, Alastuey A, Viana M-M, Rodríguez S, Artíñano B et al. 2004 Speciation and origin of PM10 and PM2.5 in Spain. *J. Aerosol Sci.* 35, 1151–1172. doi: [10.1016/j.jaerosci.2004.04.002](https://doi.org/10.1016/j.jaerosci.2004.04.002)
- Querol X, Pey J, Pandolfi M, Alastuey A, Cusack M et al. 2009 African dust contributions to mean ambient PM10 mass-levels across the Mediterranean Basin. *Atmos. Environ.* 43, 4266–4277. doi: [10.1016/j.atmosenv.2009.06.013](https://doi.org/10.1016/j.atmosenv.2009.06.013)
- Quesada B, Vautard R, Yiou P, Hirschi M, Seneviratne SI 2012 Asymmetric European summer heat predictability from wet and dry southern winters and springs. *Nat. Clim. Chang.* 2, 736–741. doi: [10.1038/nclimate1536](https://doi.org/10.1038/nclimate1536)
- Quintana-Seguí P, Martin E, Sánchez E, Zribi M, Vennetier M et al. 2016 Drought: Observed trends, future projections, in *The Mediterranean Region under Climate Change. A Scientific Update*, eds. Thiébault S, Moatti J-P (Marseille, France: Institut de Recherche pour le Développement), 123–132.
- Rabatel A, Letréguilly A, Dedieu J-P, Eckert N 2013 Changes in glacier equilibrium-line altitude in the western Alps from 1984 to 2010: evaluation by remote sensing and modeling of the morpho-topographic and climate controls. *Cryosphere* 7, 1455–1471. doi: [10.5194/tc-7-1455-2013](https://doi.org/10.5194/tc-7-1455-2013)
- Rabitsch W 2010 True Bugs (Hemiptera, Heteroptera). *BioRisk* 4, 407–433. doi: [10.3897/biorisk.4.44](https://doi.org/10.3897/biorisk.4.44)
- Raclot D, Le Bissonnais Y, Annabi M, Sabir M, Smetanova A 2018 Main issues for preserving Mediterranean soil resources from water erosion under global change. *L. Degrad. Dev.* 29, 789–799. doi: [10.1002/lrd.2774](https://doi.org/10.1002/lrd.2774)
- Rafael S, Martins H, Borrego C, Lopes M 2015 Urban vulnerability and resilience to climate change. *WIT Trans. Ecol. Environ.* 198, 379–390. doi: [10.2495/AIR150331](https://doi.org/10.2495/AIR150331)
- Raible CC, Saaroni H, Wild M 2010 Winter synoptic-scale variability over the Mediterranean Basin under future climate conditions as simulated by the ECHAM5. *Clim. Dyn.* 35, 473–488. doi: [10.1007/s00382-009-0678-5](https://doi.org/10.1007/s00382-009-0678-5)
- Räisänen J 2002 CO₂-induced changes in interannual temperature and precipitation variability in 19 CMIP2 experiments. *Bull. Am. Meteorol. Soc.* 15, 2395–2411. doi: [10.1175/1520-0442\[2002\]015%3C2395:CICIIT%3E2.0.CO;2](https://doi.org/10.1175/1520-0442[2002]015%3C2395:CICIIT%3E2.0.CO;2)
- Raitos DE, Beaugrand G, Georgopoulos D, Zenetos A, Pancucci-Papadopoulou AM et al. 2010 Global climate change amplifies the entry of tropical species into the eastern Mediterranean Sea. *Limnol. Oceanogr.* 55, 1478–1484. doi: [10.4319/lo.2010.55.4.1478](https://doi.org/10.4319/lo.2010.55.4.1478)
- Ramesh K, Matloob A, Aslam F, Florentine SK, Chauhan BS 2017 Weeds in a Changing Climate: Vulnerabilities, Consequences, and Implications for Future Weed Management. *Front. Plant Sci.* 8, 95. doi: [10.3389/fpls.2017.00095](https://doi.org/10.3389/fpls.2017.00095)
- Rascher KG, Große-Stoltenberg A, Mágua C, Meira-Neto JAA, Werner C 2011 Acacia longifolia invasion impacts vegetation structure and regeneration dynamics in open dunes and pine forests. *Biol. Invasions* 13, 1099–1113. doi: [10.1007/s10530-011-9949-2](https://doi.org/10.1007/s10530-011-9949-2)
- Rasheed T, Bilal M, Nabeel F, Adeel M, Iqbal HMN 2019 Environmentally-related contaminants of high concern: Potential sources and analytical modalities for detection, quantification, and treatment. *Env. Int.* 122, 52–66. doi: [10.1016/j.envint.2018.11.038](https://doi.org/10.1016/j.envint.2018.11.038)
- Rasmussen E, Zick C 1987 A subsynoptic vortex over the Mediterranean with some resemblance to polar lows. *Tellus A Dyn. Meteorol. Oceanogr.* 39 A, 408–425. doi: [10.1111/j.1600-0870.1987.tb00318.x](https://doi.org/10.1111/j.1600-0870.1987.tb00318.x)
- Rasmussen K, Thyrring J, Muscarella R, Borchsenius F 2017 Climate-change-induced range shifts of three allergenic ragweeds (*Ambrosia* L.) in Europe and their potential impact on human health. *PeerJ* 5, e3104. doi: [10.7717/peerj.3104](https://doi.org/10.7717/peerj.3104)
- Rasplus J-Y, Villemant C, Rosa Paiva M, Delvare G, Roques A 2010 Hymenoptera. *BioRisk* 4, 669–776. doi: [10.3897/biorisk.4.55](https://doi.org/10.3897/biorisk.4.55)
- Rassati D, Faccoli M, Petrucco Toffolo E, Battisti A, Marini L 2015 Improving the early detection of alien wood-boring beetles in ports and surrounding forests. *J. Appl. Ecol.* 52, 50–58. doi: [10.1111/1365-2664.12347](https://doi.org/10.1111/1365-2664.12347)
- Rassati D, Petrucco Toffolo E, Roques A, Battisti A, Faccoli M 2014 Trapping wood boring beetles in Italian ports: a pilot study. *J. Pest Sci.* (2004) 87, 61–69. doi: [10.1007/s10340-013-0499-5](https://doi.org/10.1007/s10340-013-0499-5)
- Ratola N, Cincinelli A, Alves A, Katsoyiannis A 2012 Occurrence of organic microcontaminants in the wastewater treatment process. A mini review. *J. Hazard. Mater.* 239–240, 1–18. doi: [10.1016/j.jhazmat.2012.05.040](https://doi.org/10.1016/j.jhazmat.2012.05.040)
- Rauthe M, Hense A, Paeth H 2004 A model intercomparison study of climate change-signals in extratropical circulation. *Int. J. Climatol.* 24, 643–662. doi: [10.1002/joc.1025](https://doi.org/10.1002/joc.1025)
- Reale M, Liberato MLR, Lionello P, Pinto JG, Salon S et al. 2019 A Global Climatology of Explosive Cyclones using a Multi-Tracking Approach. *Tellus A Dyn. Meteorol. Oceanogr.* 71, 1611340. doi: [10.1080/16000870.2019.1611340](https://doi.org/10.1080/16000870.2019.1611340)
- Reale M, Lionello P 2013 Synoptic climatology of winter intense precipitation events along the Mediterranean coasts. *Nat. Hazards Earth Syst. Sci.* 13, 1707–1722. doi: [10.5194/nhess-13-1707-2013](https://doi.org/10.5194/nhess-13-1707-2013)
- Reale O, Atlas R 2001 Tropical Cyclone-Like Vortices in the Extratropics: Observational Evidence and Synoptic Analysis. *Weather Forecast.* 16, 7–34. doi: [10.1175/1520-0434\[2001\]016<0007:tclvit>2.0.co;2](https://doi.org/10.1175/1520-0434[2001]016<0007:tclvit>2.0.co;2)

- Rees P, van der Gaag N, de Beer J, Heins F 2012 European Regional Populations: Current Trends, Future Pathways, and Policy Options. *Eur. J. Popul.* 28, 385–416.
doi: [10.1007/s10680-012-9268-z](https://doi.org/10.1007/s10680-012-9268-z)
- Reichard SH, Hamilton CW 1997 Predicting invasions of woody plants introduced into North America. *Conserv. Biol.* 11, 193–203. doi: [10.1046/j.1523-1739.1997.95473.x](https://doi.org/10.1046/j.1523-1739.1997.95473.x)
- Reimann L, Merkens J-L, Vafeidis AT 2018 Regionalized Shared Socioeconomic Pathways: Narratives and spatial population projections for the Mediterranean coastal zone. *Reg. Environ. Chang.* 18, 235–245. doi: [10.1007/s10113-017-1189-2](https://doi.org/10.1007/s10113-017-1189-2)
- Rejdali M 2004 Forest cover changes in the Maghreb countries with special reference to Morocco, in *Recent Dynamics of the Mediterranean Vegetation and Landscape*, eds. Mazzoleni S, Pasquale G Di, Mulligan M, Di Martino P, Rego F (John Wiley & Sons, Ltd), 21–31. doi: [10.1002/0470093714.ch3](https://doi.org/10.1002/0470093714.ch3)
- Rezg R, El-Fazaa S, Gharbi N, Mornagui B 2014 Bisphenol A and human chronic diseases: Current evidences, possible mechanisms, and future perspectives. *Environ. Int.* 64, 83–90. doi: [10.1016/j.envint.2013.12.007](https://doi.org/10.1016/j.envint.2013.12.007)
- Riahi K, Rao S, Krey V, Cho C, Chirkov V et al. 2011 RCP 8.5-A scenario of comparatively high greenhouse gas emissions. *Clim. Change* 109, 33–57. doi: [10.1007/s10584-011-0149-y](https://doi.org/10.1007/s10584-011-0149-y)
- Ribes A, Thao S, Vautard R, Dubuisson B, Somot S et al. 2019 Observed increase in extreme daily rainfall in the French Mediterranean. *Clim. Dyn.* 52, 1095–1114. <https://doi.org/10.1007/s00382-018-4179-2> [Accessed September 17, 2019].
- Ricart M, Guasch H, Barceló D, Brix R, Conceição MH et al. 2010 Primary and complex stressors in polluted mediterranean rivers: Pesticide effects on biological communities. *J. Hydrol.* 383, 52–61. doi: [10.1016/j.jhydrol.2009.08.014](https://doi.org/10.1016/j.jhydrol.2009.08.014)
- Rilov G 2016 Multi-species collapses at the warm edge of a warming sea. *Sci. Rep.* 6, 36897. doi: [10.1038/srep36897](https://doi.org/10.1038/srep36897)
- Rilov G, Crooks JA 2009 *Biological Invasions in Marine Ecosystems: Ecological, Management, and Geographic Perspectives*. Springer, Heidelberg, Germany.
- Rilov G, Galil BS 2009 Marine Bioinvasions in the Mediterranean Sea – History, Distribution and Ecology, in *Biological Invasions in Marine Ecosystems*, eds. Rilov G, Crooks JA (Springer-Verlag, Heidelberg, Germany.), 549–575. doi: [10.1007/978-3-540-79236-9_31](https://doi.org/10.1007/978-3-540-79236-9_31)
- Rilov G, Mazaris AD, Stelzenmüller V, Helmuth B, Wahl M et al. 2019a Adaptive marine conservation planning in the face of climate change: What can we learn from physiological, ecological and genetic studies? *Glob. Ecol. Conserv.* 17, e00566. doi: [10.1016/j.gecco.2019.e00566](https://doi.org/10.1016/j.gecco.2019.e00566)
- Rilov G, Peleg O, Guy-Haim T 2019b The Restructuring of Levant Reefs by Aliens, Ocean Warming and Overfishing: Implications to Species Interactions and Ecosystem Functions, in *Interactions in the Marine Benthos – Global Patterns and Processes*, eds. Hawkins SJ, Bohn K, Firth LB, Williams GA (Cambridge University Press, Cambridge).
- Rivetti I, Boero F, Fraschetti S, Zambianchi E, Lionello P 2017 Anomalies of the upper water column in the Mediterranean Sea. *Glob. Planet. Change* 151, 68–79. doi: [10.1016/j.gloplacha.2016.03.001](https://doi.org/10.1016/j.gloplacha.2016.03.001)
- Rivetti I, Fraschetti S, Lionello P, Zambianchi E, Boero F 2014 Global Warming and Mass Mortalities of Benthic Invertebrates in the Mediterranean Sea. *PLoS One* 9, e115655. doi: [10.1371/journal.pone.0115655](https://doi.org/10.1371/journal.pone.0115655)
- Rixen M, Beckers J-M, Levitus S, Antonov J, Boyer T et al. 2005 The Western Mediterranean Deep Water: A proxy for climate change. *Geophys. Res. Lett.* 32, n/a-n/a. doi: [10.1029/2005GL022702](https://doi.org/10.1029/2005GL022702)
- Roberts CM, O’Leary BC, McCauley DJ, Cury PM, Duarte CM et al. 2017 Marine reserves can mitigate and promote adaptation to climate change. *Proc. Natl. Acad. Sci. U. S. A.* 114, 6167–6175. doi: [10.1073/pnas.1701262114](https://doi.org/10.1073/pnas.1701262114)
- Robles-Molina J, Gilbert-López B, García-Reyes JF, Molina-Díaz A 2014 Monitoring of selected priority and emerging contaminants in the Guadalquivir River and other related surface waters in the province of Jaén, South East Spain. *Sci. Total Environ.* 479–480, 247–257. doi: [10.1016/j.scitotenv.2014.01.121](https://doi.org/10.1016/j.scitotenv.2014.01.121)
- Rochelle-Newall EJ, Delesalle B, Mari X, Rouchon C, Torréton JP et al. 2008 Zinc induces shifts in microbial carbon flux in tropical coastal environments. *Aquat. Microb. Ecol.* 52, 57–68. doi: [10.3354/ame01212](https://doi.org/10.3354/ame01212)
- Rodger HD, Henry L, Mitchell SO 2011 Non-infectious gill disorders of marine salmonid fish. *Rev. Fish Biol. Fish.* 21, 423–440. doi: [10.1007/s11160-010-9182-6](https://doi.org/10.1007/s11160-010-9182-6)
- Rodó X 2001 Reversal of three global atmospheric fields linking changes in SST anomalies in the Pacific, Atlantic and Indian oceans at tropical latitudes and midlatitudes. *Clim. Dyn.* 18, 203–217. doi: [10.1007/s003820100171](https://doi.org/10.1007/s003820100171)
- Rodó X, Baert E, Comin FA 1997 Variations in seasonal rainfall in Southern Europe during the present century: relationships with the North Atlantic Oscillation and the El Niño-Southern Oscillation. *Clim. Dyn.* 13, 275–284.
- Rodríguez-Fonseca B, de Castro M 2002 On the connection between winter anomalous precipitation in the Iberian Peninsula and North West Africa and the summer subtropical Atlantic sea surface temperature. *Geophys. Res. Lett.* 29, 10-1–10-4. doi: [10.1029/2001gl014421](https://doi.org/10.1029/2001gl014421)
- Rodríguez-Lázaro D, Cook N, Ruggeri FM, Sellwood J, Nasser A et al. 2012 Virus hazards from food, water and other contaminated environments. *FEMS Microbiol. Rev.* 36, 786–814. doi: [10.1111/j.1574-6976.2011.00306.x](https://doi.org/10.1111/j.1574-6976.2011.00306.x)
- Rodríguez S, Querol X, Alastuey A, de la Rosa J 2007 Atmospheric particulate matter and air quality in the Mediterranean: a review. *Environ. Chem. Lett.* 5, 1–7. doi: [10.1007/s10311-006-0071-0](https://doi.org/10.1007/s10311-006-0071-0)
- Rodríguez S, Querol X, Alastuey A, Viana M-M, Mantilla E 2003 Events Affecting Levels and Seasonal Evolution of Airborne Particulate Matter Concentrations in the Western Mediterranean. *Environ. Sci. Technol.* 37, 216–222. doi: [10.1021/es020106p](https://doi.org/10.1021/es020106p)
- Rodwell MJ, Hoskins BJ 1996 Monsoons and the dynamics of deserts. *Q. J. R. Meteorol. Soc.* 122, 1385–1404. doi: [10.1002/qj.49712253408](https://doi.org/10.1002/qj.49712253408)
- Roether W, Klein B, Manca BB, Theocharis A, Kioroglou S 2007 Transient Eastern Mediterranean deep waters in response to the massive dense-water output of the Aegean Sea in the 1990s. *Prog. Oceanogr.* 74, 540–571.

- doi: [10.1016/j.pocean.2007.03.001](https://doi.org/10.1016/j.pocean.2007.03.001)
- Rohat G, Flacke J, Dao H, van Maarseveen M 2018 Co-use of existing scenario sets to extend and quantify the shared socioeconomic pathways. *Clim. Change* 151, 619–636. doi: [10.1007/s10584-018-2318-8](https://doi.org/10.1007/s10584-018-2318-8)
- Rohling EJ, Bryden HL 1992 Man-induced salinity and temperature increases in Western Mediterranean deep water. *JGR Atmos.* 97, 11191–11198. doi: [10.1029/92JC00767](https://doi.org/10.1029/92JC00767)
- Roig N, Sierra J, Rovira J, Schuhmacher M, Domingo JL et al. 2013 In vitro tests to assess toxic effects of airborne PM10 samples. Correlation with metals and chlorinated dioxins and furans. *Sci. Total Environ.* 443, 791–797. doi: [10.1016/j.scitotenv.2012.11.022](https://doi.org/10.1016/j.scitotenv.2012.11.022)
- Rojas M, Li LZ, Kanakidou M, Hatzianastassiou N, Seze G et al. 2013 Winter weather regimes over the Mediterranean region: their role for the regional climate and projected changes in the twenty-first century. *Clim. Dyn.* 41, 551–571.
- Roll U, Dayan T, Simberloff D, Goren M 2007 Characteristics of the introduced fish fauna of Israel. *Biol. Invasions* 9, 813–824. doi: [10.1007/s10530-006-9083-8](https://doi.org/10.1007/s10530-006-9083-8)
- Romera R, Gaertner MÁ, Sánchez E, Domínguez M, González-Alemán JJ et al. 2017 Climate change projections of medicanes with a large multi-model ensemble of regional climate models. *Glob. Planet. Change* 151, 134–143. doi: [10.1016/j.gloplacha.2016.10.008](https://doi.org/10.1016/j.gloplacha.2016.10.008)
- Romero R, Emanuel K 2013 Medicane risk in a changing climate. *JGR Atmos.* 118, 5992–6001. doi: [10.1002/jgrd.50475](https://doi.org/10.1002/jgrd.50475)
- Romero R, Emanuel K 2017 Climate Change and Hurricane-Like Extratropical Cyclones: Projections for North Atlantic Polar Lows and Medicanes Based on CMIP5 Models. *J. Clim.* 30, 279–299. doi: [10.1175/jcli-d-16-0255.1](https://doi.org/10.1175/jcli-d-16-0255.1)
- Romo S, Soria J, Fernandez F, Ouahid Y, Baron-Sola A 2013 Water residence time and the dynamics of toxic cyanobacteria. *Freshw. Biol.* 58, 513–522. doi: [10.1111/j.1365-2427.2012.02734.x](https://doi.org/10.1111/j.1365-2427.2012.02734.x)
- Roques A 2010 Taxonomy, time and geographic patterns. Chapter 2. *BioRisk* 4, 11–26. doi: [10.3897/biorisk.4.70](https://doi.org/10.3897/biorisk.4.70)
- Roques A 2015 Drivers and pathways of forest insect invasions in Europe, can we predict the next arrivals? *Atti Accad. Naz. Ital. di Entomol.* 53, 145–150.
- Roques A, Auger-Rozenberg M-A, Blackburn TM, Garnas J, Pyšek P et al. 2016 Temporal and interspecific variation in rates of spread for insect species invading Europe during the last 200 years. *Biol. Invasions* 18, 907–920. doi: [10.1007/s10530-016-1080-y](https://doi.org/10.1007/s10530-016-1080-y)
- Roques A, Kenis M, Lees D, Lopez-Vaamonde C, Rabitsch W et al. 2010 Alien terrestrial arthropodes of Europe. *BIOISK*, 1028. doi: [10.1007/978-1-4020-8280-1_5](https://doi.org/10.1007/978-1-4020-8280-1_5)
- Roques A, Preda C, Augustin S, Auger-Rozenberg M-A 2018 Bugs, ants, wasps, moths and other insect species, in *Invasive species and human health*, eds. Mazzia G, Tricarico E (CABI, Wallingford, UK), 63–75. doi: [10.1079/9781786390981.0063](https://doi.org/10.1079/9781786390981.0063)
- Rotenberg E, Yakir D 2011 Distinct patterns of changes in surface energy budget associated with forestation in the semi-arid region. *Glob. Chang. Biol.* 17, 1536–1548. doi: [10.1111/j.1365-2486.2010.02320.x](https://doi.org/10.1111/j.1365-2486.2010.02320.x)
- Rotmans J, van Asselt M, Anastasi C, Greeuw S, Mellors J et al. 2000 Visions for a sustainable Europe. *Futures* 32, 809–831. doi: [10.1016/s0016-3287\(00\)00033-1](https://doi.org/10.1016/s0016-3287(00)00033-1)
- Rousi E, Anagnostopoulou C, Tolika K, Maher P 2015 Representing teleconnection patterns over Europe: A comparison of SOM and PCA methods. *Atmos. Res.* 152, 123–137. doi: [10.1016/j.atmosres.2013.11.010](https://doi.org/10.1016/j.atmosres.2013.11.010)
- Rousis NI, Bade R, Bijlsma L, Zuccato E, Sancho J V. et al. 2017 Monitoring a large number of pesticides and transformation products in water samples from Spain and Italy. *Environ. Res.* 156, 31–38. doi: [10.1016/j.envres.2017.03.013](https://doi.org/10.1016/j.envres.2017.03.013)
- Roy HE, Bacher S, Essl F, Adriaens T, Aldridge DC et al. 2019 Developing a list of invasive alien species likely to threaten biodiversity and ecosystems in the European Union. *Glob. Chang. Biol.* 25, gcb.14527. doi: [10.1111/gcb.14527](https://doi.org/10.1111/gcb.14527)
- Roy HE, Rabitsch W, Scalera R, Stewart A, Gallardo B et al. 2018 Developing a framework of minimum standards for the risk assessment of alien species. *J. Appl. Ecol.* 55, 526–538. doi: [10.1111/1365-2664.13025](https://doi.org/10.1111/1365-2664.13025)
- Roy HE, Rhule E, Harding S, Lawson Handley L-J, Poland RL et al. 2011 Living with the enemy: Parasites and pathogens of the ladybird *Harmonia axyridis*. *BioControl* 56, 663–679. doi: [10.1007/s10526-011-9387-1](https://doi.org/10.1007/s10526-011-9387-1)
- Ruffault J, Mouillet F, Peters DPC 2015 How a new fire-suppression policy can abruptly reshape the fire-weather relationship. *Ecosphere* 6, 1–19. doi: [10.1890/ES15-00182.1](https://doi.org/10.1890/ES15-00182.1)
- Rugman-Jones PF, Hoddle CD, Hoddle MS, Stouthamer R 2013 The Lesser of Two Weevils: Molecular genetics of pest palm weevil populations confirm *Rhynchophorus vulneratus* (Panzer 1798) as a valid species distinct from *R. ferrugineus* (Olivier 1790), and reveal the global extent of both. *PLoS One* 8, e78379. doi: [10.1371/journal.pone.0078379](https://doi.org/10.1371/journal.pone.0078379)
- Ruiz-Ramos M, Ferrise R, Rodríguez A, Lorite IJ, Bindi M et al. 2018 Adaptation response surfaces for managing wheat under perturbed climate and CO₂ in a Mediterranean environment. *Agric. Syst.* 159, 260–274. doi: [10.1016/j.jagsy.2017.01.009](https://doi.org/10.1016/j.jagsy.2017.01.009)
- Rusiñol M, Fernandez-Cassi X, Timoneda N, Carratalà A, Abril JF et al. 2015 Evidence of viral dissemination and seasonality in a Mediterranean river catchment: Implications for water pollution management. *J. Environ. Manage.* 159, 58–67. doi: [10.1016/j.jenvman.2015.05.019](https://doi.org/10.1016/j.jenvman.2015.05.019)
- Russo MA, Leitão J, Gama C, Ferreira J, Monteiro A 2018 Shipping emissions over Europe: A state-of-the-art and comparative analysis. *Atmos. Environ.* 177, 187–194. doi: [10.1016/j.atmosenv.2018.01.025](https://doi.org/10.1016/j.atmosenv.2018.01.025)
- Russo S, Dosio A, Graversen RG, Sillmann J, Carrao H et al. 2014 Magnitude of extreme heat waves in present climate and their projection in a warming world. *JGR Atmos.* 119, 12,500–12,512. doi: [10.1002/2014jd022098](https://doi.org/10.1002/2014jd022098)
- Ruti PM, Somot S, Giorgi F, Dubois C, Flaounas E et al. 2016 Med-CORDEX initiative for Mediterranean climate studies. *Bull. Am. Meteorol. Soc.* 97, 1187–1208. doi: [10.1175/BAMS-D-14-00176.1](https://doi.org/10.1175/BAMS-D-14-00176.1)
- Ryan PG, Moore CJ, van Franeker JA, Moloney CL 2009 Monitoring the abundance of plastic debris in the marine environment. *Philos. Trans. R. Soc. B Biol. Sci.* 364, 1999–2012. doi: [10.1098/rstb.2008.0207](https://doi.org/10.1098/rstb.2008.0207)

- Saadi S, Todorovic M, Tanasijevic L, Pereira LS, Pizzigalli C et al. 2015 Climate change and Mediterranean agriculture: Impacts on winter wheat and tomato crop evapotranspiration, irrigation requirements and yield. *Agric. Water Manag.* 147, 103–115. doi: [10.1016/J.AGWAT.2014.05.008](https://doi.org/10.1016/J.AGWAT.2014.05.008)
- Saaroni H, Ziv B, Edelson J, Alpert P 2003 Long-term variations in summer temperatures over the Eastern Mediterranean. *Geophys. Res. Lett.* 30. doi: [10.1029/2003gl017742](https://doi.org/10.1029/2003gl017742)
- Sabatés A, Martín P, Lloret J, Raya V 2006 Sea warming and fish distribution: The case of the small pelagic fish, *Sardinella aurita*, in the western Mediterranean. *Glob. Chang. Biol.* 12, 2209–2219. doi: [10.1111/j.1365-2486.2006.01246.x](https://doi.org/10.1111/j.1365-2486.2006.01246.x)
- Sabatés A, Martin P, Raya V, Sabatés A, Martín P et al. 2012 Changes in life-history traits in relation to climate change: bluefish (*Pomatomus saltatrix*) in the northwestern Mediterranean. *ICES J. Mar. Sci.* 69, 1000–1009. doi: [10.1093/icesjms/fss053](https://doi.org/10.1093/icesjms/fss053)
- Sabatés A, Salat J, Raya V, Emelianov M, Segura-Noguera M 2009 Spawning environmental conditions of *Sardinella aurita* at the northern limit of its distribution range, the western Mediterranean. *Mar. Ecol. Prog. Ser.* 385, 227–236. doi: [10.3354/meps08058](https://doi.org/10.3354/meps08058)
- Sabine CL, Feely RA, Gruber N, Key RM, Lee K et al. 2004 The Oceanic Sink for Anthropogenic CO₂. *Science* 305, 367–371. doi: [10.1126/science.1097403](https://doi.org/10.1126/science.1097403)
- Sáenz J, Rodríguez-Puebla C, Fernández J, Zubillaga J 2001 Interpretation of interannual winter temperature variations over southwestern Europe. *JGR Atmos.* 106, 20641–20651. doi: [10.1029/2001JD900247](https://doi.org/10.1029/2001JD900247)
- Saffioti C, Fischer EM, Scherrer SC, Knutti R 2016 Reconciling observed and modeled temperature and precipitation trends over Europe by adjusting for circulation variability. *Geophys. Res. Lett.* 43, 8189–8198. doi: [10.1002/2016GL069802](https://doi.org/10.1002/2016GL069802)
- Sahraoui I, Grami B, Bates SS, Bouchouicha D, Chikhaoui MA et al. 2012 Response of potentially toxic Pseudo-nitzschia (Bacillariophyceae) populations and domoic acid to environmental conditions in a eutrophied, SW Mediterranean coastal lagoon (Tunisia). *Estuar. Coast. Shelf Sci.* 102, 95–104. doi: [10.1016/j.ecss.2012.03.018](https://doi.org/10.1016/j.ecss.2012.03.018)
- Sahu LK, Saxena P 2015 High time and mass resolved PTR-TOF-MS measurements of VOCs at an urban site of India during winter: Role of anthropogenic, biomass burning, biogenic and photochemical sources. *Atmos. Res.* 164–165, 84–94. doi: [10.1016/J.ATMOSRES.2015.04.021](https://doi.org/10.1016/J.ATMOSRES.2015.04.021)
- Sahu LK, Yadav R, Pal D 2016 Source identification of VOCs at an urban site of western India: Effect of marathon events and anthropogenic emissions. *JGR Atmos.* 121, 2416–2433. doi: [10.1002/2015JD024454](https://doi.org/10.1002/2015JD024454)
- Sakka Hlaili A, Chikhaoui M-A, El Grami B, Mabrouk HH 2006 Effects of N and P supply on phytoplankton in Bizerte Lagoon (western Mediterranean). *J. Exp. Mar. Bio. Ecol.* 333, 79–96. doi: [10.1016/j.jembe.2005.12.049](https://doi.org/10.1016/j.jembe.2005.12.049)
- Sakka Hlaili A, Sahraoui I, Bouchouicha D, Meliti Garali S, Ksouri J et al. 2016 Toxic and potentially toxic diatoms blooms in SW Mediterranean waters: review of ten years investigations, in *Advances in Environmental Research*, ed. Daniels JA (Nova Science Publishers, Inc.).
- Sala E, Kizilkaya Z, Yildirim D, Ballesteros E 2011 Alien Marine Fishes Deplete Algal Biomass in the Eastern Mediterranean. *PLoS One* 6, e17356. doi: [10.1371/journal.pone.0017356](https://doi.org/10.1371/journal.pone.0017356)
- Saliba NA, Massoud R 2010 A Comparative Review of PM Levels, Sources, and Their Likely Fates in the Eastern Mediterranean Region, in *Environmental Science and Engineering (Subseries: Environmental Science)*, eds. Zereini F, Wiseman C (Springer, Berlin, Heidelberg), 3–17. doi: [10.1007/978-3-642-12278-1_1](https://doi.org/10.1007/978-3-642-12278-1_1)
- Saliba NA, Moussa S, Salame H, El-Fadel M 2006 Variation of selected air quality indicators over the city of Beirut, Lebanon: Assessment of emission sources. *Atmos. Environ.* 40, 3263–3268. doi: [10.1016/j.atmosenv.2006.01.054](https://doi.org/10.1016/j.atmosenv.2006.01.054)
- Sallustio L, di Cristofaro M, Hashmi M, Vizzarri M, Sitzia T et al. 2018 Evaluating the Contribution of Trees outside Forests and Small Open Areas to the Italian Landscape Diversification during the Last Decades. *Forests* 9, 701. doi: [10.3390/f9110701](https://doi.org/10.3390/f9110701)
- Salvador P, Artíñano B, Viana M-M, Alastuey A, Querol X 2012 Evaluation of the changes in the Madrid metropolitan area influencing air quality: Analysis of 1999–2008 temporal trend of particulate matter. *Atmos. Environ.* 57, 175–185. doi: [10.1016/J.ATMOSENV.2012.04.026](https://doi.org/10.1016/J.ATMOSENV.2012.04.026)
- Salvatí A, Monti P, Coch Roura H, Cecere C 2019 Climatic performance of urban textures: Analysis tools for a Mediterranean urban context. *Energy Build.* 185, 162–179. doi: [10.1016/j.enbuild.2018.12.024](https://doi.org/10.1016/j.enbuild.2018.12.024)
- Samaniego L, Thober S, Kumar R, Wanders N, Rakovec O et al. 2018 Anthropogenic warming exacerbates European soil moisture droughts. *Nat. Clim. Chang.* 8, 421–426. doi: [10.1038/s41558-018-0138-5](https://doi.org/10.1038/s41558-018-0138-5)
- San-Miguel-Ayanz J, Moreno JM, Camia A 2013 Analysis of large fires in European Mediterranean landscapes: Lessons learned and perspectives. *For. Ecol. Manage.* 294, 11–22. doi: [10.1016/j.foreco.2012.10.050](https://doi.org/10.1016/j.foreco.2012.10.050)
- Sánchez-Benítez A, García-Herrera R, Barriopedro D, Sousa PM, Trigo RM 2018 June 2017: The Earliest European Summer Mega-heatwave of Reanalysis Period. *Geophys. Res. Lett.* 45, 1955–1962. doi: [10.1002/2018GL077253](https://doi.org/10.1002/2018GL077253)
- Sánchez-Gómez E, Somot S, Déqué M 2009 Ability of an ensemble of regional climate models to reproduce weather regimes over Europe-Atlantic during the period 1961–2000. *Clim. Dyn.* 33, 723–736. doi: [10.1007/s00382-008-0502-7](https://doi.org/10.1007/s00382-008-0502-7)
- Sánchez-Lorenzo A, Brunetti M, Calbó J, Martín-Vide J 2007 Recent spatial and temporal variability and trends of sunshine duration over the Iberian Peninsula from a homogenized data set. *JGR Atmos.* 112, D20115. doi: [10.1029/2007JD008677](https://doi.org/10.1029/2007JD008677)
- Sánchez-Lorenzo A, Calbó J, Wild M 2013 Global and diffuse solar radiation in Spain: Building a homogeneous dataset and assessing their trends. *Glob. Planet. Change* 100, 343–352. doi: [10.1016/J.GLOPLACHA.2012.11.010](https://doi.org/10.1016/J.GLOPLACHA.2012.11.010)
- Sánchez-Lorenzo A, Enriquez-Alonso A, Calbó J, González J-A, Wild M et al. 2017 Fewer clouds in the Mediterranean: consistency of observations and climate simulations. *Sci. Rep.* 7, 41475. doi: [10.1038/srep41475](https://doi.org/10.1038/srep41475)

- Sánchez J, Bisquert M, Rubio E, Caselles V 2015 Impact of Land Cover Change Induced by a Fire Event on the Surface Energy Fluxes Derived from Remote Sensing. *Remote Sens.* 7, 14899–14915. doi: [10.3390/rs71114899](https://doi.org/10.3390/rs71114899)
- Sanna S, Le Tellier J 2013 Building on the Mediterranean scenario experiences: Cross-cutting approaches between regional foresight analysis and participatory prospective. Valbonne: Plan Bleu.
- Santi Delia A, Caruso G, Melcarne L, Caruso G, Parisi S et al. 2015 Biological Toxins from Marine and Freshwater Microalgae, in *Microbial Toxins and Related Contamination in the Food Industry*, eds. Caruso G, Caruso G, Laganà P, Santi Delia A, Parisi S et al. (Springer International Publishing).
- Santini A, Ghelardini L, de Pace C, Desprez-Loustau ML, Capretti P et al. 2013 Biogeographical patterns and determinants of invasion by forest pathogens in Europe. *New Phytol.* 197, 238–250. doi: [10.1111/j.1469-8137.2012.04364.x](https://doi.org/10.1111/j.1469-8137.2012.04364.x)
- Sarris D, Christopoulou A, Angelonidi E, Koutsias N, Fulé PZ et al. 2014 Increasing extremes of heat and drought associated with recent severe wildfires in southern Greece. *Reg. Environ. Chang.* 14, 1257–1268. doi: [10.1007/s10113-013-0568-6](https://doi.org/10.1007/s10113-013-0568-6)
- Saura S, Estreguil C, Mouton C, Rodriguez-Freire M 2011 Network analysis to assess landscape connectivity trends: application to European forests (1990–2000). *Ecol. Indic.* 11, 407–416. doi: [10.1016/j.ecolind.2010.06.011](https://doi.org/10.1016/j.ecolind.2010.06.011)
- Sauvé S, Desrosiers M 2014 A review of what is an emerging contaminant. *Chem. Cent. J.* 8. doi: [10.1186/1752-153X-8-15](https://doi.org/10.1186/1752-153X-8-15)
- Schär C, Vidale PL, Lüthi D, Frei C, Häberli C et al. 2004 The role of increasing temperature variability in European summer heatwaves. *Nature* 427, 332–336. doi: [10.1038/nature02300](https://doi.org/10.1038/nature02300)
- Schembri C, Cavalli F, Cuccia E, Hjorth J, Calzolai G et al. 2012 Impact of a European directive on ship emissions on air quality in Mediterranean harbours. *Atmos. Environ.* 61, 661–669. doi: [10.1016/j.atmosenv.2012.06.047](https://doi.org/10.1016/j.atmosenv.2012.06.047)
- Schneider A, Tanhua T, Körtzinger A, Wallace DWR 2010 High anthropogenic carbon content in the eastern Mediterranean. *J. Geophys. Res. Ocean.* 115. doi: [10.1029/2010JC006171](https://doi.org/10.1029/2010JC006171)
- Schroeder K, Chiggiato J, Bryden HL, Borghini M, Ben Ismail S 2016 Abrupt climate shift in the Western Mediterranean Sea. *Sci. Rep.* 6. doi: [10.1038/srep23009](https://doi.org/10.1038/srep23009)
- Schroeder K, Chiggiato J, Josey SA, Borghini M, Aracri S et al. 2017 Rapid response to climate change in a marginal sea. *Sci. Rep.* 7, 4065. doi: [10.1038/s41598-017-04455-5](https://doi.org/10.1038/s41598-017-04455-5)
- Schroeder K, García Lafuente J, Josey SA, Artale V, Buongiorno Nardelli B et al. 2012 Circulation of the Mediterranean Sea and its variability, in *The Climate of the Mediterranean Region: From the Past to the Future*, ed. Lionello P (Elsevier). doi: [10.1016/B978-0-12-416042-2.00012-4](https://doi.org/10.1016/B978-0-12-416042-2.00012-4)
- Scyphers SB, Powers SP, Akins JL, Drymon JM, Martin CW et al. 2015 The role of citizens in detecting and responding to a rapid marine invasion. *Conserv. Lett.* 8, 242–250. doi: [10.1111/conl.12127](https://doi.org/10.1111/conl.12127)
- Seager R, Liu H, Henderson N, Simpson IR, Kelley C et al. 2014 Causes of increasing aridification of the Mediterranean region in response to rising greenhouse gases. *J. Clim.* 27, 4655–4676. doi: [10.1175/jcli-d-13-00446.1](https://doi.org/10.1175/jcli-d-13-00446.1)
- Seebens H, Blackburn TM, Dyer EE, Genovesi P, Hulme PE et al. 2017 No saturation in the accumulation of alien species worldwide. *Nat. Commun.* 8, 14435. doi: [10.1038/ncomms14435](https://doi.org/10.1038/ncomms14435)
- Seebens H, Gastner MT, Blasius B 2013 The risk of marine bio-invasion caused by global shipping. *Ecol. Lett.* 16, 782–790. doi: [10.1111/ele.12111](https://doi.org/10.1111/ele.12111)
- Seinfeld JH, Pandis SN 2006 *Atmospheric chemistry and physics : from air pollution to climate change*. 2nd ed. Hoboken, N.J. : Wiley.
- Seljak G 2013 Dinamika vnosa tujerodnih fitofagnih žuželk in pršic v Slovenijo. *Acta Entomol. Slov.* 21, 85–122.
- Sellner KG, Doucette GJ, Kirkpatrick GJ 2003 Harmful algal blooms: Causes, impacts and detection. *J. Ind. Microbiol. Biotechnol.* 30, 383–406. doi: [10.1007/s10295-003-0074-9](https://doi.org/10.1007/s10295-003-0074-9)
- Sen OL, Ezber Y, Bozkurt D 2019 Euro-Mediterranean climate variability in boreal winter: A potential role of the East Asian trough. *Clim. Dyn.* 52, 7071–7084. doi: [10.1007/s00382-018-4573-9](https://doi.org/10.1007/s00382-018-4573-9)
- Senar JC, Domènech J, Arroyo L, Torre I, Gordo D 2016 An evaluation of monk parakeet damage to crops in the metropolitan area of Barcelona. *Anim. Biodivers. Conserv.* 39, 141–145. doi: [10.32800/abc.2016.39.0141](https://doi.org/10.32800/abc.2016.39.0141)
- Seneviratne SI, Lüthi D, Litschi M, Schär C 2006 Land-atmosphere coupling and climate change in Europe. *Nature* 443, 205–209. doi: [10.1038/nature05095](https://doi.org/10.1038/nature05095)
- Senouci R, Taibi N-E 2019 Impact of the urbanization on coastal dune: Case of Kharrouba, West of Algeria. *J. Sediment. Environ.* 4, 90–98. doi: [10.12957/jse.2019.39951](https://doi.org/10.12957/jse.2019.39951)
- Sevault F, Somot S, Alias A, Dubois C, Lebeaupin-Brossier C et al. 2014 A fully coupled Mediterranean regional climate system model: design and evaluation of the ocean component for the 1980–2012 period. *Tellus A Dyn. Meteorol. Oceanogr.* 66, 23967. doi: [10.3402/tellusa.v66.23967](https://doi.org/10.3402/tellusa.v66.23967)
- Severin T, Conan P, de Madron XD, Houpert L, Oliver MJ et al. 2014 Impact of open-ocean convection on nutrients, phytoplankton biomass and activity. *Deep Sea Res. Part I Oceanogr. Res. Pap.* 94, 62–71. doi: [10.1016/j.dsr.2014.07.015](https://doi.org/10.1016/j.dsr.2014.07.015)
- Shaltout M, Omstedt A 2014 Recent sea surface temperature trends and future scenarios for the Mediterranean Sea. *Oceanologia* 56, 411–443. doi: [10.5697/oc.56-3.411](https://doi.org/10.5697/oc.56-3.411)
- Shaw TA, Baldwin M, Barnes EA, Caballero R, Garfinkel CI et al. 2016 Storm track processes and the opposing influences of climate change. *Nat. Geosci.* 9, 656–664. doi: [10.1038/ngeo2783](https://doi.org/10.1038/ngeo2783)
- Sheffield J, Wood EF 2008 Global Trends and Variability in Soil Moisture and Drought Characteristics, 1950–2000, from Observation-Driven Simulations of the Terrestrial Hydrologic Cycle. *J. Clim.* 21, 432–458. doi: [10.1175/2007JCLI1822.1](https://doi.org/10.1175/2007JCLI1822.1)
- Shepherd TG 2014 Atmospheric circulation as a source of uncertainty in climate change projections. *Nat. Geosci.* 7, 703–708. doi: [10.1038/NGEO2253](https://doi.org/10.1038/NGEO2253)
- Shindell DT, Faluvegi G, Koch DM, Schmidt GA, Unger N et al. 2009 Improved Attribution of Climate Forcing to Emissions. *Science (80-.).* 326, 716–718. doi: [10.1126/science.1174760](https://doi.org/10.1126/science.1174760)
- Shindell DT, Lamarque J-F, Schulz M, Flanner M, Jiao C et al. 2013 Radiative forcing in the ACCMIP historical and future

- climate simulations. *Atmos. Chem. Phys.* 13, 2939–2974. doi: [10.5194/acp-13-2939-2013](https://doi.org/10.5194/acp-13-2939-2013)
- Shomar BH, Müller G, Yahya A 2006 Occurrence of Pesticides in Groundwater and Topsoil of the Gaza Strip. *Water, Air, Soil Pollut.* 171, 237–251. doi: [10.1007/s11270-005-9038-1](https://doi.org/10.1007/s11270-005-9038-1)
- Sicard P, de Marco A, Troussier F, Renou C, Vas N et al. 2013 Decrease in surface ozone concentrations at Mediterranean remote sites and increase in the cities. *Atmos. Environ.* 79, 705–715. doi: [10.1016/j.atmosenv.2013.07.042](https://doi.org/10.1016/j.atmosenv.2013.07.042)
- Sillmann J, Kharin VV, Zwiers FW, Zhang X, Bronaugh D 2013 Climate extremes indices in the CMIP5 multimodel ensemble: Part 2. Future climate projections. *JGR Atmos.* 118, 2473–2493. doi: [10.1002/jgrd.50188](https://doi.org/10.1002/jgrd.50188)
- Simpson IR, Seager R, Shaw TA, Ting M 2015 Mediterranean summer climate and the importance of middle East topography. *J. Clim.* 28, 1977–1996. doi: [10.1175/jcli-d-14-00298.1](https://doi.org/10.1175/jcli-d-14-00298.1)
- Singh AN, Kumar P 2017 Emerging Contaminants – An Concise Overview.
- Sinović G, Franičević M, Zorica B, Čikeš-Keč V 2004 Short Communication Length-weight and length-length relationships for 10 pelagic fish species from the Adriatic Sea (Croatia). *J. Appl. Ichthyol.* 20, 156–158. doi: [10.1046/j.1439-0426.2003.00519.x](https://doi.org/10.1046/j.1439-0426.2003.00519.x)
- Sire A, Amouroux I 2016 Détermination de Valeurs Guides Environnementales (VGE) mollusques alternatives aux Normes de Qualité Environnementale (NQE) définies dans la DCE. <https://archimer.ifremer.fr/doc/0033/44378/> [Accessed April 17, 2019].
- Sisma-Ventura G, Yam R, Shemesh A 2014 Recent unprecedented warming and oligotrophy of the eastern Mediterranean Sea within the last millennium. *Geophys. Res. Lett.* 41, 5158–5166. doi: [10.1002/2014GL060393](https://doi.org/10.1002/2014GL060393)
- Skejic S, Car A, Marasovic I, Jozic S, Buzancic M et al. 2017 Morphology and ecology of the poorly known dinoflagellate *Prorocentrum arcuatum* (Dinophyceae) from the Medulin Bay (eastern Adriatic Sea). *Acta Adriat.* 58, 41–52.
- Skliris N, Zika JD, Herold L, Josey SA, Marsh R 2018 Mediterranean sea water budget long-term trend inferred from salinity observations. *Clim. Dyn.* 51, 2857–2876. doi: [10.1007/s00382-017-4053-7](https://doi.org/10.1007/s00382-017-4053-7)
- Slanger ABA, Adloff F, Jevrejeva S, Leclercq PW, Marzeion B et al. 2017 A review of recent updates of sea-level projections at global and regional scales. *Surv. Geophys.* 38, 395–416. doi: [10.1007/978-3-319-56490-6_17](https://doi.org/10.1007/978-3-319-56490-6_17)
- Slimani H, Aidoud A 2004 Desertification in the Maghreb: A case study of an Algerian high-plain steppe, in *Environmental Challenges in the Mediterranean 2000–2050*, ed. Marquina A (Dordrecht, Netherlands: Springer).
- Smith IM 1979 EPPO: the work of a regional plant protection organization, with particular reference to phytosanitary regulations, in *Plant Health: The Scientific Basis for Administrative Control of Plant Diseases and Pests*, eds. Ebbels DL, King JE (Oxford, UK: Blackwell Scientific Publications).
- Smouni A, Ater M, Auguy F, Laplaze L, El Mzibri M et al. 2010 Assessment of contamination by metallic trace elements in a mining area of eastern Morocco. *Cah. Agric.* 19, 273–279. doi: [10.1684/agr.2010.0413](https://doi.org/10.1684/agr.2010.0413)
- Sobrino E, Sanz-Elorza M, Dana ED, González-Moreno A 2002 Invasibility of a coastal strip in NE Spain by alien plants. *J. Veg. Sci.* 13, 585–594. doi: [10.1111/j.1654-1103.2002.tb02085.x](https://doi.org/10.1111/j.1654-1103.2002.tb02085.x)
- Somot S, Houpert L, Sevault F, Testor P, Bosse A et al. 2018 Characterizing, modelling and understanding the climate variability of the deep water formation in the North-Western Mediterranean Sea. *Clim. Dyn.* 51, 1179–1210.
- Somot S, Sevault F, Déqué M 2006 Transient climate change scenario simulation of the Mediterranean Sea for the twenty-first century using a high-resolution ocean circulation model. *Clim. Dyn.* 27, 851–879. doi: [10.1007/s00382-006-0167-z](https://doi.org/10.1007/s00382-006-0167-z)
- Somot S, Sevault F, Déqué M, Crépon M 2008 21st century climate change scenario for the Mediterranean using a coupled atmosphere-ocean regional climate model. *Glob. Planet. Change* 63, 112–126. doi: [10.1016/j.gloplacha.2007.10.003](https://doi.org/10.1016/j.gloplacha.2007.10.003)
- Soto-Navarro J, Jordà G, Amores A, Cabos W, Somot S et al. 2020 Evolution of Mediterranean Sea water properties under climate change scenarios in the Med-CORDEX ensemble. *Clim. Dyn.* 54, 2135–2165. doi: [10.1007/s00382-019-05105-4](https://doi.org/10.1007/s00382-019-05105-4)
- Soto-Navarro J, Somot S, Sevault F, Beuvier J, Criado-Aldeanueva F et al. 2015 Evaluation of regional ocean circulation models for the Mediterranean Sea at the Strait of Gibraltar: volume transport and thermohaline properties of the outflow. *Clim. Dyn.* 44, 1277–1292. doi: [10.1007/s00382-014-2179-4](https://doi.org/10.1007/s00382-014-2179-4)
- Spinoni J, Naumann G, Vogt J, Barbosa P 2015 European drought climatologies and trends based on a multi-indicator approach. *Glob. Planet. Change* 127, 50–57. doi: [10.1016/J.GLOPLACHA.2015.01.012](https://doi.org/10.1016/J.GLOPLACHA.2015.01.012)
- Spinoni J, Vogt J V., Naumann G, Barbosa P, Dosio A 2018 Will drought events become more frequent and severe in Europe? *Int. J. Climatol.* 38, 1718–1736. doi: [10.1002/joc.5291](https://doi.org/10.1002/joc.5291)
- Squadrone S, Brizio P, Stella C, Prearo M, Pastorino P et al. 2016 Presence of trace metals in aquaculture marine ecosystems of the northwestern Mediterranean Sea (Italy). *Environ. Pollut.* 215, 77–83. doi: [10.1016/j.envpol.2016.04.096](https://doi.org/10.1016/j.envpol.2016.04.096)
- Stachowicz JJ, Terwin JR, Whitlatch RB, Osman RW 2002 Linking climate change and biological invasions: Ocean warming facilitates nonindigenous species invasions. *Proc. Natl. Acad. Sci. U. S. A.* 99, 15497–15500. doi: [10.1073/pnas.242437499](https://doi.org/10.1073/pnas.242437499)
- Stagge JH, Kohn I, Tallaksen LM, Stahl K 2015 Modeling drought impact occurrence based on meteorological drought indices in Europe. *J. Hydrol.* 530, 37–50. doi: [10.1016/j.jhydrol.2015.09.039](https://doi.org/10.1016/j.jhydrol.2015.09.039)
- Stamatis N, Hela D, Triantafyllidis V, Konstantinou I 2013 Spatiotemporal variation and risk assessment of pesticides in water of the lower catchment basin of Acheloos River, Western Greece. *ScientificWorldJournal.* 2013, 231610. doi: [10.1155/2013/231610](https://doi.org/10.1155/2013/231610)
- Statuto D, Cillis G, Picuno P 2016 Analysis of the effects of agricultural land use change on rural environment and

- landscape through historical cartography and GIS tools. *J. Agric. Eng.* 47, 468.
- Stehle S, Schulz R 2015 Agricultural insecticides threaten surface waters at the global scale. *Proc. Natl. Acad. Sci. U. S. A.* 112, 5750–5755. doi: [10.1073/pnas.1500232112](https://doi.org/10.1073/pnas.1500232112)
- Stephenson DB, Pavan V, Collins M, Junge MM, Quadrelli R 2006 North Atlantic Oscillation response to transient greenhouse gas forcing and the impact on European winter climate: A CMIP2 multi-model assessment. *Clim. Dyn.* 27, 401–420. doi: [10.1007/s00382-006-0140-x](https://doi.org/10.1007/s00382-006-0140-x)
- Stergiou KI, Somarakis S, Triantafyllou G, Tsiaras KP, Giannoulaiki M et al. 2016 Trends in productivity and biomass yields in the Mediterranean Sea Large Marine Ecosystem during climate change. *Environ. Dev.* 17, 57–74. doi: [10.1016/j.envdev.2015.09.001](https://doi.org/10.1016/j.envdev.2015.09.001)
- Stergiou KI, Tsikliras AC 2011 Fishing down, fishing through and fishing up: fundamental process versus technical details. *Mar. Ecol. Prog. Ser.* 441, 295–301. doi: [10.3354/meps09377](https://doi.org/10.3354/meps09377)
- Stinca A, Chirico GB, Incerti G, Bonanomi G 2015 Regime shift by an exotic nitrogen-fixing shrub mediates plant facilitation in primary succession. *PLoS One* 10, e0123128. doi: [10.1371/journal.pone.0123128](https://doi.org/10.1371/journal.pone.0123128)
- Stohl A, Aamaas B, Amann M, Baker LH, Bellouin N et al. 2015 Evaluating the climate and air quality impacts of short-lived pollutants. *Atmos. Chem. Phys.* 15, 10529–10566. doi: [10.5194/acp-15-10529-2015](https://doi.org/10.5194/acp-15-10529-2015)
- Storkey J, Strattonovitch P, Chapman DS, Vidotto F, Semenov MA 2014 A Process-Based Approach to Predicting the Effect of Climate Change on the Distribution of an Invasive Allergenic Plant in Europe. *PLoS One* 9, e88156. doi: [10.1371/journal.pone.0088156](https://doi.org/10.1371/journal.pone.0088156)
- Strode SA, Ott LE, Pawson S, Bowyer TW 2012 Emission and transport of cesium-137 from boreal biomass burning in the summer of 2010. *JGR Atmos.* 117, n/a-n/a. doi: [10.1029/2011JD017382](https://doi.org/10.1029/2011JD017382)
- Sturaro E, Thiene M, Cocca G, Mrad M, Tempesta T et al. 2013 Factors influencing summer farms management in the Alps. *Ital. J. Anim. Sci.* 12, e25. doi: [10.4081/ijas.2013.e25](https://doi.org/10.4081/ijas.2013.e25)
- Suffert M, Wilstermann A, Petter F, Schrader G, Grousset F 2018 Identification of new pests likely to be introduced into Europe with the fruit trade. *Bull. OEPP/EPPO* 48, 144–154.
- Sumaila UR, Cheung WWL, Lam VWY, Pauly D, Herrick S 2011 Climate change impacts on the biophysics and economics of world fisheries. *Nat. Clim. Chang.* 1, 449–456. doi: [10.1038/nclimate1301](https://doi.org/10.1038/nclimate1301)
- Sumaila UR, Tai TC, Lam VWY, Cheung WWL, Bailey M et al. 2019 Benefits of the Paris Agreement to ocean life, economies, and people. *Sci. Adv.* 5, eaau3855. doi: [10.1126/sciadv.aau3855](https://doi.org/10.1126/sciadv.aau3855)
- Sun X, Ren G, You Q, Ren Y, Xu W et al. 2019 Global diurnal temperature range (DTR) changes since 1901. *Clim. Dyn.* 52, 3343–3356. doi: [10.1007/s00382-018-4329-6](https://doi.org/10.1007/s00382-018-4329-6)
- Taillefumier F, Piégay H 2003 Contemporary land use changes in Prealpine Mediterranean mountains. A multivariate GIS-based approach applied to two municipalities in the Southern French Prealps. *Catena* 51, 267–296. doi: [10.1016/S0341-8162\(02\)00168-6](https://doi.org/10.1016/S0341-8162(02)00168-6)
- Talbi A, Kerchich Y, Kerbachi R, Boughedaoui M 2018 Assessment of annual air pollution levels with PM1, PM2.5, PM10 and associated heavy metals in Algiers, Algeria. *Environ. Pollut.* 232, 252–263. doi: [10.1016/j.envpol.2017.09.041](https://doi.org/10.1016/j.envpol.2017.09.041)
- Tanarhte M, Hadjinicolaou P, Lelieveld J 2012 Intercomparison of temperature and precipitation data sets based on observations in the Mediterranean and the Middle East. *JGR Atmos.* 117, n/a-n/a. doi: [10.1029/2011jd017293](https://doi.org/10.1029/2011jd017293)
- Tanasićević L, Todorović M, Pereira LS, Pizzigalli C, Lionello P 2014 Impacts of climate change on olive crop evapotranspiration and irrigation requirements in the Mediterranean region. *Agric. Water Manag.* 144, 54–68. doi: [10.1016/J.AGWAT.2014.05.019](https://doi.org/10.1016/J.AGWAT.2014.05.019)
- Tanhua T, Hainbucher D, Schroeder K, Cardin V, Alvarez M et al. 2013 The Mediterranean Sea system: a review and an introduction to the special issue. *Ocean Sci.* 9, 789–803. doi: [10.5194/os-9-789-2013](https://doi.org/10.5194/os-9-789-2013)
- Tarabon S, Bertrand R, Lavoie C, Vigouroux T, Isselin-Nondedeu F 2018 The effects of climate warming and urbanised areas on the future distribution of Cortaderia selloana, pampas grass, in France. *Weed Res.* 58, 413–423. doi: [10.1111/wre.12330](https://doi.org/10.1111/wre.12330)
- Taylor KE, Stouffer RJ, Meehl GA 2012 An Overview of CMIP5 and the Experiment Design. *Bull. Am. Meteorol. Soc.* 93, 485–498. doi: [10.1175/BAMS-D-11-00094.1](https://doi.org/10.1175/BAMS-D-11-00094.1)
- Taylor PD, Fahrig L, Henein K, Merriam G 1993 Connectivity is a vital element of landscape structure. *Oikos* 68, 571–573.
- Tchounwou PB, Yedjou CG, Patlolla AK, Sutton DJ 2012 Heavy metal toxicity and the environment. *Exp. Suppl.* 101, 133–164. doi: [10.1007/978-3-7643-8340-4_6](https://doi.org/10.1007/978-3-7643-8340-4_6)
- Testor P, Bosse A, Houpert L, Margirier F, Mortier L et al. 2018 Multiscale Observations of Deep Convection in the Northwestern Mediterranean Sea During Winter 2012–2013 Using Multiple Platforms. *J. Geophys. Res. Ocean.* 123, 1745–1776. doi: [10.1002/2016JC012671](https://doi.org/10.1002/2016JC012671)
- Testor P, de Madron XD, Mortier L, D'Ortenzio F, Legoff H et al. 2019 LION observatory data. *SEANOE*. doi: [10.17882/44411](https://doi.org/10.17882/44411)
- Thiéblemont R, Le Cozannet G, Toimil A, Meyssignac B, Losada IJ 2019 Likely and High-End Impacts of Regional Sea-Level Rise on the Shoreline Change of European Sandy Coasts Under a High Greenhouse Gas Emissions Scenario. *Water* 11, 2607.
- Thiery W, Davin EL, Lawrence DM, Hirsch AL, Hauser M et al. 2017 Present-day irrigation mitigates heat extremes. *J. Geophys. Res. Atmos.* 122, 1403–1422. doi: [10.1002/2016JD025740](https://doi.org/10.1002/2016JD025740)
- Thornes JB 2009 Land degradation, in *The Physical Geography of the Mediterranean*, ed. Woodward J (New York: Oxford University Press), 563–581.
- Thorpe A, Harrison RM 2008 Sources and properties of non-exhaust particulate matter from road traffic: A review. *Sci. Total Environ.* 400, 270–282. doi: [10.1016/J.SCITOTENV.2008.06.007](https://doi.org/10.1016/J.SCITOTENV.2008.06.007)
- Thorpe RB, Bigg GR 2000 Modelling the sensitivity of Mediterranean outflow to anthropogenically forced climate change. *Clim. Dyn.* 16, 355–368. doi: [10.1007/s003820050333](https://doi.org/10.1007/s003820050333)
- Titelboim D, Almogi-Labin A, Herut B, Kucera M, Aschner-Polivoda S et al. 2019 Thermal tolerance and range

- expansion of invasive foraminifera under climate changes. *Sci. Rep.* 9, 1–5. doi: [10.1038/s41598-019-40944-5](https://doi.org/10.1038/s41598-019-40944-5)
- Titelboim D, Sadekov A, Almogi-Labin A, Herut B, Kucera M et al. 2017 Geochemical signatures of benthic foraminiferal shells from a heat-polluted shallow marine environment provide field evidence for growth and calcification under extreme warmth. *Glob. Chang. Biol.* 23, 4346–4353. doi: [10.1111/gcb.13729](https://doi.org/10.1111/gcb.13729)
- Tiwari D, Kamble J, Chilgunde S, Patil P, Maru G et al. 2012 Clastogenic and mutagenic effects of bisphenol A: An endocrine disruptor. *Mutat. Res. - Genet. Toxicol. Environ. Mutagen.* 743, 83–90. doi: [10.1016/j.mrgentox.2011.12.023](https://doi.org/10.1016/j.mrgentox.2011.12.023)
- Tobin I, Vautard R, Balog I, Bréon F-M, Jerez S et al. 2015 Assessing climate change impacts on European wind energy from ENSEMBLES high-resolution climate projections. *Clim. Change* 128, 99–112. doi: [10.1007/s10584-014-1291-0](https://doi.org/10.1007/s10584-014-1291-0)
- Tolika K 2019 Assessing Heat Waves over Greece Using the Excess Heat Factor (EHF). *Climate* 7, 9. doi: [10.3390/climatology7010009](https://doi.org/10.3390/climatology7010009)
- Toreti A, Xoplaki E, Maraun D, Kuglitsch FG, Wanner H et al. 2010 Characterisation of extreme winter precipitation in Mediterranean coastal sites and associated anomalous atmospheric circulation patterns. *Nat. Hazards Earth Syst. Sci.* 10, 1037–1050. doi: [10.5194/nhess-10-1037-2010](https://doi.org/10.5194/nhess-10-1037-2010)
- Torma CZ 2019 Detailed validation of EURO-CORDEX and Med-CORDEX regional climate model ensembles over the Carpathian Region. *Q. J. Hungarian Meteorol. Serv.* 123, 217–240.
- Torma CZ, Giorgi F, Coppola E 2015 Added value of regional climate modeling over areas characterized by complex terrain-Precipitation over the Alps. *JGR Atmos.* 120, 3957–3972. doi: [10.1002/2014JD022781](https://doi.org/10.1002/2014JD022781)
- Touaylia S, Ghannem S, Toumi H, Bejaoui M 2016 Assessment of heavy metals status in northern Tunisia using contamination indices: Case of the Ichkeul steams system. 3, 209–217.
- Touratier F, Goyet C 2011 Impact of the Eastern Mediterranean Transient on the distribution of anthropogenic CO₂ and first estimate of acidification for the Mediterranean Sea. *Deep Sea Res. Part I Oceanogr. Res. Pap.* 58, 1–15. doi: [10.1016/j.dsr.2010.10.002](https://doi.org/10.1016/j.dsr.2010.10.002)
- Tous M, Romero R 2013 Meteorological environments associated with medicane development. *Int. J. Climatol.* 33, 1–14. doi: [10.1002/joc.3428](https://doi.org/10.1002/joc.3428)
- Tous M, Zappa G, Romero R, Shaffrey L, Vidale PL 2016 Projected changes in medicanes in the HadGEM3 N512 high-resolution global climate model. *Clim. Dyn.* 47, 1913–1924. doi: [10.1007/s00382-015-2941-2](https://doi.org/10.1007/s00382-015-2941-2)
- Tovar-Sánchez A, Basterretxea G, Ben Omar M, Jordi A, Sánchez-Quiles D et al. 2016 Nutrients, trace metals and B-vitamin composition of the Moulouya River: A major North African river discharging into the Mediterranean Sea. *Estuar. Coast. Shelf Sci.* 176, 47–57. doi: [10.1016/j.ecss.2016.04.006](https://doi.org/10.1016/j.ecss.2016.04.006)
- Traina A, Bono G, Bonsignore M, Falco F, Giuga M et al. 2019 Heavy metals concentrations in some commercially key species from Sicilian coasts (Mediterranean Sea): Potential human health risk estimation. *Ecotoxicol. Environ. Saf.* 168, 466–478. doi: [10.1016/j.ecoenv.2018.10.056](https://doi.org/10.1016/j.ecoenv.2018.10.056)
- Tramblay Y, Feki H, Quintana-Seguí P, Guijarro JA 2019 The SAFRAN daily gridded precipitation product in Tunisia (1979–2015). *Int. J. Climatol.*, joc.6181. doi: [10.1002/joc.6181](https://doi.org/10.1002/joc.6181)
- Tribouillois H, Constantin J, Justes E 2018 Cover crops mitigate direct greenhouse gases balance but reduce drainage under climate change scenarios in temperate climate with dry summers. *Glob. Chang. Biol.* 24, 2513–2529. doi: [10.1111/gcb.14091](https://doi.org/10.1111/gcb.14091)
- Trigo IF, Bigg GR, Davies TD 2002a Climatology of Cyclogenesis Mechanisms in the Mediterranean. *Mon. Weather Rev.* 130, 549–569. doi: [10.1175/1520-0493\(2002\)130<0549:COCMIT>2.0.CO;2](https://doi.org/10.1175/1520-0493(2002)130<0549:COCMIT>2.0.CO;2)
- Trigo IF, Davies TD, Bigg GR 1999 Objective Climatology of Cyclones in the Mediterranean Region. *J. Clim.* 12, 1685–1696. doi: [10.1175/1520-0442\(1999\)012<1685:OCOCIT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<1685:OCOCIT>2.0.CO;2)
- Trigo IF, Davies TD, Bigg GR 2000 Decline in Mediterranean rainfall caused by weakening of Mediterranean cyclones. *Geophys. Res. Lett.* 27, 2913–2916. doi: [10.1029/2000GL011526](https://doi.org/10.1029/2000GL011526)
- Trigo R, Xoplaki E, Zorita E, Luterbacher J, Krichak SO et al. 2006 Chapter 3 Relations between variability in the Mediterranean region and mid-latitude variability, in *Mediterranean*, eds. Lionello P, Malanotte-Rizzoli P, Boscolo RBT-D in E and ES [Elsevier], 179–226. doi: [10.1016/S1571-9197\(06\)80006-6](https://doi.org/10.1016/S1571-9197(06)80006-6)
- Trigo RM, Osborn TJ, Corte-Real JM 2002b The North Atlantic Oscillation influence on Europe: climate impacts and associated physical mechanisms. *Clim. Res.* 20, 9–17. doi: [10.3354/cr02009](https://doi.org/10.3354/cr02009)
- Trigo RM, Pozo-Vázquez D, Osborn TJ, Castro-Díez Y, Gámiz-Fortis S et al. 2004 North Atlantic oscillation influence on precipitation, river flow and water resources in the Iberian Peninsula. *Int. J. Climatol.* 24, 925–944. doi: [10.1002/joc.1048](https://doi.org/10.1002/joc.1048)
- Trouet V 2014 A Tree-Ring Based Late Summer Temperature Reconstruction (AD 1675–1980) for the Northeastern Mediterranean. *Radiocarbon* 56, S69–S78. doi: [10.1017/s0033822200050372](https://doi.org/10.1017/s0033822200050372)
- Tsikliras AC 2008 Climate-related geographic shift and sudden population increase of a small pelagic fish (*Sardinella aurita*) in the eastern Mediterranean Sea. *Mar. Biol. Res.* 4, 477–481. doi: [10.1080/17451000802291292](https://doi.org/10.1080/17451000802291292)
- Tsikliras AC 2014 Fisheries Mismanagement in the Mediterranean: A Greek Tragedy. *Fish. Aquac. J.* 05. doi: [10.4172/2150-3508.1000e113](https://doi.org/10.4172/2150-3508.1000e113)
- Tsikliras AC, Dinouli A, Tsalkou E 2013a Exploitation trends of the Mediterranean and Black Sea fisheries. *Acta Adriat.* 54, 273–282.
- Tsikliras AC, Dinouli A, Tsilos V-Z, Tsalkou E 2015 The Mediterranean and Black Sea Fisheries at Risk from Overexploitation. *PLoS One* 10, e0121188. doi: [10.1371/journal.pone.0121188](https://doi.org/10.1371/journal.pone.0121188)
- Tsikliras AC, Licandro P, Pardalou A, McQuinn IH, Gröger JP et al. 2019 Synchronization of Mediterranean pelagic fish populations with the North Atlantic climate variability. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 159, 143–151.

- doi: [10.1016/j.dsr2.2018.07.005](https://doi.org/10.1016/j.dsr2.2018.07.005)
- Tsikliras AC, Stergiou KI 2014 Mean temperature of the catch increases quickly in the Mediterranean Sea. *Mar. Ecol. Prog. Ser.* 515, 281–284. doi: [10.3354/meps11005](https://doi.org/10.3354/meps11005)
- Tsikliras AC, Tsilos VZ, Stergiou KI 2013b Assessing the state of Greek marine fisheries resources. *Fish. Manag. Ecol.* 20, 34–41. doi: [10.1111/j.1365-2400.2012.00863.x](https://doi.org/10.1111/j.1365-2400.2012.00863.x)
- Tsimplis MN, Álvarez-Fanjul E, Gomis D, Fenoglio-Marc L, Pérez B 2005 Mediterranean Sea level trends: Atmospheric pressure and wind contribution. *Geophys. Res. Lett.* 32, L20602. doi: [10.1029/2005GL023867](https://doi.org/10.1029/2005GL023867)
- Tsimplis MN, Zervakis V, Josey S, Peneva E, Struglia MV et al. 2006 Variability of the Mediterranean sea level and oceanic circulation and their relation to climate patterns, in *Mediterranean climate variability*, eds. Lionello P, Malanotte-Rizzoli P, Boscolo R (Elsevier).
- Tsopelas P, Santini A, Wingfield MJ, de Beer ZW 2017 Canker stain: A lethal disease destroying iconic plane trees. *Plant Dis.* 101, 645–658. doi: [10.1094/PDIS-09-16-1235-FE](https://doi.org/10.1094/PDIS-09-16-1235-FE)
- Turbé A, Strubbe D, Mori E, Carrete M, Chiron F et al. 2017 Assessing the assessments: evaluation of four impact assessment protocols for invasive alien species. *Divers. Distrib.* 23, 297–307. doi: [10.1111/ddi.12528](https://doi.org/10.1111/ddi.12528)
- Turco M, Bedía J, di Liberto F, Fiorucci P, Von Hardenberg J et al. 2016 Decreasing fires in Mediterranean Europe. *PLoS One* 11. doi: [10.1371/journal.pone.0150663](https://doi.org/10.1371/journal.pone.0150663)
- Turco M, Jerez S, Augusto S, Tarín-Carrasco P, Ratola N et al. 2019 Climate drivers of the 2017 devastating fires in Portugal. *Sci. Rep.* 9, 13886. doi: [10.1038/s41598-019-50281-2](https://doi.org/10.1038/s41598-019-50281-2)
- Turco M, Llasat MC, von Hardenberg J, Provenzale A 2014 Climate change impacts on wildfires in a Mediterranean environment. *Clim. Change* 125, 369–380. doi: [10.1007/s10584-014-1183-3](https://doi.org/10.1007/s10584-014-1183-3)
- Turco M, Rosa-Cánovas JJ, Bedía J, Jerez S, Montávez JP et al. 2018 Exacerbated fires in Mediterranean Europe due to anthropogenic warming projected with non-stationary climate-fire models. *Nat. Commun.* 9, 3821. doi: [10.1038/s41467-018-06358-z](https://doi.org/10.1038/s41467-018-06358-z)
- Türkeş M, Erlat E 2003 Precipitation changes and variability in Turkey linked to the North Atlantic oscillation during the period 1930–2000. *Int. J. Climatol.* 23, 1771–1796. doi: [10.1002/joc.962](https://doi.org/10.1002/joc.962)
- Tyrlis E, Hoskins BJ 2008 Aspects of a northern hemisphere atmospheric blocking climatology. *J. Atmos. Sci.* 65, 1638–1652. doi: [10.1175/2007JAS2337.1](https://doi.org/10.1175/2007JAS2337.1)
- Tyrlis E, Lelieveld J 2013 Climatology and Dynamics of the Summer Etesian Winds over the Eastern Mediterranean*. *J. Atmos. Sci.* 70, 3374–3396. doi: [10.1175/jas-d-13-0351](https://doi.org/10.1175/jas-d-13-0351)
- Tzanatos E, Raitsos DE, Triantafyllou G, Somarakis S, Tsonis AA 2014 Indications of a climate effect on Mediterranean fisheries. *Clim. Change* 122, 41–54. doi: [10.1007/s10584-013-0972-4](https://doi.org/10.1007/s10584-013-0972-4)
- Ulbrich U, Christoph M 1999 A shift of the NAO and increasing storm track activity over Europe due to anthropogenic greenhouse gas forcing. *Clim. Dyn.* 15, 551–559. doi: [10.1007/s003820050299](https://doi.org/10.1007/s003820050299)
- Ulbrich U, Leckebusch GC, Pinto JG 2009 Extra-tropical cyclones in the present and future climate: A review. *Theor. Appl. Climatol.* 96, 117–131. doi: [10.1007/s00704-008-0083-8](https://doi.org/10.1007/s00704-008-0083-8)
- Ulbrich U, Lionello P, Belušić D, Jacobbeit J, Knippertz P et al. 2012 Climate of the Mediterranean: Synoptic patterns, temperature, precipitation, winds and their extremes, in *The Climate of the Mediterranean Region: From the Past to the Future*, ed. Lionello P (Elsevier), 1–61.
- Ulbrich U, Pinto JG, Kupfer H, Leckebusch GC, Spangehl T et al. 2008 Changing Northern Hemisphere Storm Tracks in an Ensemble of IPCC Climate Change Simulations. *J. Clim.* 21, 1669–1679. doi: [10.1175/2007jcli1992.1](https://doi.org/10.1175/2007jcli1992.1)
- Ullah S, Zuberi A, Alagawany M, Farag MR, Dadar M et al. 2018 Cypermethrin induced toxicities in fish and adverse health outcomes: Its prevention and control measure adaptation. *J. Environ. Manage.* 206, 863–871. doi: [10.1016/j.jenvman.2017.11.076](https://doi.org/10.1016/j.jenvman.2017.11.076)
- UN-Water 2014 *The United Nations World Water Development Report* 3. doi: [10.4324/9781849773355](https://doi.org/10.4324/9781849773355)
- UN 2015 Transforming our world: the 2030 Agenda for Sustainable Development. New York https://www.un.org/ga/search/view_doc.asp?symbol=A/RES/70/1&Lang=E
- Underwood EC, Viers JH, Klausmeyer KR, Cox RL, Shaw MR 2009 Threats and biodiversity in the Mediterranean biome. *Divers. Distrib.* 15, 188–197. doi: [10.1111/j.1472-4642.2008.00518.x](https://doi.org/10.1111/j.1472-4642.2008.00518.x)
- UNEP-GRID Arendal 2013 State of the Mediterranean Marine and Coastal Environment.
- UNEP/MAP 2012a Initial integrated assessment of the Mediterranean sea: fulfilling step 3 of the ecosystem approach process.
- UNEP/MAP 2012b State of the Mediterranean Marine and Coastal Environment. UNEP/MAP – Barcelona Convention, Athens.
- UNEP/MAP 2017 2017 Mediterranean Quality Status Report.
- Valladares F, Matesanz S, Guilhaumon F, Araújo MB, Balaguer L et al. 2014 The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change. *Ecol. Lett.* 17, 1351–1364. doi: [10.1111/ele.12348](https://doi.org/10.1111/ele.12348)
- Van der Sluis T, Pedroli B, Kristensen SBP, Cosor GL, Pavlis E 2016 Changing land use intensity in Europe—Recent processes in selected case studies. *Land use policy* 57, 777–785. doi: [10.1016/j.landusepol.2014.12.005](https://doi.org/10.1016/j.landusepol.2014.12.005)
- Van Dingenen R, Raes F, Putaud J-P, Baltensperger U, Charron A et al. 2004 A European aerosol phenomenology—1: physical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe. *Atmos. Environ.* 38, 2561–2577. doi: [10.1016/J.ATMOSENV.2004.01.040](https://doi.org/10.1016/J.ATMOSENV.2004.01.040)
- Van Kleunen M, Essl F, Pergl J, Brundu G, Carboni M et al. 2018 The changing role of ornamental horticulture in alien plant invasions. *Biol. Rev.* 93, 1421–1437. doi: [10.1111/brv.12402](https://doi.org/10.1111/brv.12402)
- van Ruijen BJ, Levy MA, Agrawal A, Biermann F, Birkmann J et al. 2014 Enhancing the relevance of Shared Socioeconomic Pathways for climate change impacts, adaptation and vulnerability research. *Clim. Change* 122, 481–494. doi: [10.1007/s10584-013-0931-0](https://doi.org/10.1007/s10584-013-0931-0)
- van Vuuren DP, Edmonds JE, Kainuma M, Riahi K, Thomson A

- et al. 2011 The representative concentration pathways: An overview. *Clim. Change* 109, 5–31.
doi: [10.1007/s10584-011-0148-z](https://doi.org/10.1007/s10584-011-0148-z)
- Vanderhoeven S, Branquart E, Casaer J, D'hondt B, Hulme PE et al. 2017 Beyond protocols: improving the reliability of expert-based risk analysis underpinning invasive species policies. *Biol. Invasions* 19, 2507–2517.
doi: [10.1007/s10530-017-1434-0](https://doi.org/10.1007/s10530-017-1434-0)
- Vargas-Yáñez M, García-Martínez MC, Moya F, Balbín R, López-Jurado JL et al. 2017 Updating temperature and salinity mean values and trends in the Western Mediterranean: The RADMED project. *Prog. Oceanogr.* 157, 27–46.
doi: [10.1016/J.POCEAN.2017.09.004](https://doi.org/10.1016/J.POCEAN.2017.09.004)
- Vargas-Yáñez M, Zunino P, Benali A, Delpy M, Pastre F et al. 2010 How much is the western Mediterranean really warming and salting? *JGR Atmos.* 115, C04001.
doi: [10.1029/2009jc005816](https://doi.org/10.1029/2009jc005816)
- Vasilakopoulos P, Maravelias CD, Tserpes G 2014 The alarming decline of mediterranean fish stocks. *Curr. Biol.* 24, 1643–1648. doi: [10.1016/j.cub.2014.05.070](https://doi.org/10.1016/j.cub.2014.05.070)
- Vasilakopoulos P, Raitsos DE, Tzanatos E, Maravelias CD 2017 Resilience and regime shifts in a marine biodiversity hotspot. *Sci. Rep.* 7. doi: [10.1038/s41598-017-13852-9](https://doi.org/10.1038/s41598-017-13852-9)
- Vautard R, Cattiaux J, Yiou P, Thépaut J-N, Ciais P 2010 Northern Hemisphere atmospheric stilling partly attributed to an increase in surface roughness. *Nat. Geosci.* 3, 756–761.
doi: [10.1038/ngeo979](https://doi.org/10.1038/ngeo979)
- Vautard R, Gobiet A, Sobolowski SP, Kjellström E, Stegehuis A et al. 2014 The European climate under a 2°C global warming. *Environ. Res. Lett.* 9, 34006.
doi: [10.1088/1748-9326/9/3/034006](https://doi.org/10.1088/1748-9326/9/3/034006)
- Vayreda J, Martínez-Vilalta J, Gracia M, Retana J 2012 Recent climate changes interact with stand structure and management to determine changes in tree carbon stocks in Spanish forests. *Glob. Chang. Biol.* 18, 1028–1041.
doi: [10.1111/j.1365-2486.2011.02606.x](https://doi.org/10.1111/j.1365-2486.2011.02606.x)
- Velaoras D, Kassis D, Perivoliotis L, Pagonis P, Hondronasios A et al. 2013 Temperature and salinity variability in the Greek Seas based on POSEIDON stations time series: preliminary results. *Mediterr. Mar. Sci.* 14, 5–18. doi: [10.12681/mms.446](https://doi.org/10.12681/mms.446)
- Verdura J, Linares C, Ballesteros E, Coma R, Uriz MJ et al. 2019 Biodiversity loss in a Mediterranean ecosystem due to an extreme warming event unveils the role of an engineering gorgonian species. *Sci. Rep.* 9.
doi: [10.1038/s41598-019-41929-0](https://doi.org/10.1038/s41598-019-41929-0)
- Verfaillie D, Lafaysse M, Déqué M, Eckert N, Lejeune Y et al. 2018 Multi-component ensembles of future meteorological and natural snow conditions for 1500 m altitude in the Chartreuse mountain range, Northern French Alps. *Cryosph.* 12, 1249–1271. doi: [10.5194/tc-12-1249-2018](https://doi.org/10.5194/tc-12-1249-2018)
- Vergés A, Doropoulos C, Malcolm HA, Skye M, García-Pizá M et al. 2016 Long-term empirical evidence of ocean warming leading to tropicalization of fish communities, increased herbivory, and loss of kelp. *Proc. Natl. Acad. Sci. U. S. A.* 113, 13791–13796. doi: [10.1073/pnas.1610725113](https://doi.org/10.1073/pnas.1610725113)
- Vergés A, Steinberg PD, Hay ME, Poore AGB, Campbell AH et al. 2014a The tropicalization of temperate marine ecosystems: climate-mediated changes in herbivory and community phase shifts. *Proc. R. Soc. B Biol. Sci.* 281, 20140846.
doi: [10.1098/rspb.2014.0846](https://doi.org/10.1098/rspb.2014.0846)
- Vergés A, Tomas F, Cebrán E, Ballesteros E, Kızılıkaya Z et al. 2014b Tropical rabbitfish and the deforestation of a warming temperate sea. *J. Ecol.* 102, 1518–1527.
doi: [10.1111/1365-2745.12324](https://doi.org/10.1111/1365-2745.12324)
- Verheyen J, Stoks R 2019 Current and future daily temperature fluctuations make a pesticide more toxic: Contrasting effects on life history and physiology. *Environ. Pollut.* 248, 209–218. doi: [10.1016/j.envpol.2019.02.022](https://doi.org/10.1016/j.envpol.2019.02.022)
- Verkerk PJ, Sánchez A, Libbrecht S, Broekman A, Bruggeman A et al. 2017 A participatory approach for adapting river basins to climate change. *Water* 9, 958.
doi: [10.3390/w9120958](https://doi.org/10.3390/w9120958)
- Vetraino AM, Roques A, Yart A, Fan J, Sun J et al. 2015 Sentinel trees as a tool to forecast invasions of alien plant pathogens. *PLoS One* 10, e0120571. doi: [10.1371/journal.pone.0120571](https://doi.org/10.1371/journal.pone.0120571)
- Vicente-Serrano SM, Domínguez-Castro F, Murphy C, Hannaford J, Reig F et al. 2020 Long-term variability and trends in meteorological droughts in Western Europe (1851–2018). *Int. J. Climatol.* n/a. doi: [10.1002/joc.6719](https://doi.org/10.1002/joc.6719)
- Vicente-Serrano SM, López Moreno JI, Beguería S, Lorenzo-Lacruz J, Sánchez-Lorenzo A et al. 2014 Evidence of increasing drought severity caused by temperature rise in southern Europe. *Environ. Res. Lett.* 9.
doi: [10.1088/1748-9326/9/4/044001](https://doi.org/10.1088/1748-9326/9/4/044001)
- Vicente-Serrano SM, López Moreno JI, Gimeno L, Nieto R, Morán-Tejeda E et al. 2011 A multiscalar global evaluation of the impact of ENSO on droughts. *JGR Atmos.* 116, D20109. doi: [10.1029/2011JD016039](https://doi.org/10.1029/2011JD016039)
- Vicente-Serrano SM, Van der Schrier G, Beguería S, Azorin-Molina C, López Moreno JI 2015 Contribution of precipitation and reference evapotranspiration to drought indices under different climates. *J. Hydrol.* 526, 42–54.
doi: [10.1016/j.jhydrol.2014.11.025](https://doi.org/10.1016/j.jhydrol.2014.11.025)
- Vieira G, Mora C, Fahle A 2017 New observations indicate the possible presence of permafrost in North Africa (Djebel Toubkal, High Atlas, Morocco). *Cryosph.* 11, 1691–1705.
doi: [10.5194/tc-11-1691-2017](https://doi.org/10.5194/tc-11-1691-2017)
- Vilà-Cabrera A, Coll L, Martínez-Vilalta J, Retana J 2018 Forest management for adaptation to climate change in the Mediterranean basin: A synthesis of evidence. *For. Ecol. Manage.* 407, 16–22. doi: [10.1016/j.foreco.2017.10.021](https://doi.org/10.1016/j.foreco.2017.10.021)
- Vila M, Camp J, Garces E, Maso M, Delgado M 2001 High resolution spatio-temporal detection of potentially harmful dinoflagellates in confined waters of the NW Mediterranean. *J. Plankton Res.* 23, 497–514. doi: [10.1093/plankt/23.5.497](https://doi.org/10.1093/plankt/23.5.497)
- Vilà M, Espinar JL, Hejda M, Hulme PE, Jarošík V et al. 2011 Ecological impacts of invasive alien plants: a meta-analysis of their effects on species, communities and ecosystems. *Ecol. Lett.* 14, 702–708.
doi: [10.1111/j.1461-0248.2011.01628.x](https://doi.org/10.1111/j.1461-0248.2011.01628.x)
- Vilà M, Hulme PE 2017 *Impact of Biological Invasions on Ecosystem Services*. Springer
- Vilà M, Pino J, Font X 2007 Regional assessment of plant inva-

- sions across different habitat types. *J. Veg. Sci.* 18, 35–42. doi: [10.1111/j.1654-1103.2007.tb02513.x](https://doi.org/10.1111/j.1654-1103.2007.tb02513.x)
- Vilà M, Pino J, Montero A, Font X 2010 Are island plant communities more invaded than their mainland counterparts? *J. Veg. Sci.* 21, 438–446. <https://www.jstor.org/stable/40925501>
- Vilà M, Pujadas J 2001 Land-use and socio-economic correlates of plant invasions in European and North African countries. *Biol. Conserv.* 100, 397–401. doi: [10.1016/s0006-3207\(01\)00047-7](https://doi.org/10.1016/s0006-3207(01)00047-7)
- Vilà M, Tessier M, Suehs CM, Brundu G, Carta L et al. 2006 Local and regional assessments of the impacts of plant invaders on vegetation structure and soil properties of Mediterranean islands. *J. Biogeogr.* 33, 853–861. doi: [10.1111/j.1365-2699.2005.01430.x](https://doi.org/10.1111/j.1365-2699.2005.01430.x)
- Vilibić I, Čikeš Keč V, Zorica B, Šepić J, Matijević S et al. 2016 Hydrographic conditions driving sardine and anchovy populations in a land-locked sea. *Mediterr. Mar. Sci.* 17, 1–12. doi: [10.12681/mms.1120](https://doi.org/10.12681/mms.1120)
- von Schuckmann K, Le Traon P-Y, Smith N, Pascual A, Brasseur P et al. 2018 Copernicus Marine Service Ocean State Report. *J. Oper. Oceanogr.* 11, S1–S142. doi: [10.1080/1755876X.2018.1489208](https://doi.org/10.1080/1755876X.2018.1489208)
- Vousdoukas MI, Mentaschi L, Voukouvalas E, Verlaan M, Feyen L 2017 Extreme sea levels on the rise along Europe's coasts. *Earth's Futur.* 5, 304–323. doi: [10.1002/2016EF000505](https://doi.org/10.1002/2016EF000505)
- Wahl M, Buchholz B, Winde V, Golomb D, Guy-Haim T et al. 2015 A mesocosm concept for the simulation of near-natural shallow underwater climates: The Kiel Outdoor Benthocosms (KOB). *Limnol. Oceanogr. Methods* 13, 651–663. doi: [10.1002/lom3.10055](https://doi.org/10.1002/lom3.10055)
- Waldman R, Somot S, Herrmann M, Sevault F, Isachsen PE 2018 On the chaotic variability of deep convection in the Mediterranean Sea. *Geophys. Res. Lett.* 45, 2433–2443. doi: [10.1002/2017GL076319](https://doi.org/10.1002/2017GL076319)
- Walsh K, Giorgi F, Coppola E 2014 Mediterranean warm-core cyclones in a warmer world. *Clim. Dyn.* 42, 1053–1066. doi: [10.1007/s00382-013-1723-y](https://doi.org/10.1007/s00382-013-1723-y)
- Walther GR, Roques A, Hulme PE, Sykes MT, Pyšek P et al. 2009 Alien species in a warmer world: risks and opportunities. *Trends Ecol. Evol.* 24, 686–693. doi: [10.1016/j.tree.2009.06.008](https://doi.org/10.1016/j.tree.2009.06.008)
- Wang F, Polcher J 2019 Assessing the freshwater flux from the continents to the Mediterranean Sea. *Sci. Rep.* 9, 8024. doi: [10.1038/s41598-019-44293-1](https://doi.org/10.1038/s41598-019-44293-1)
- Watt MS, Kriticos DJ, Lamoureux SL, Bourdôt GW 2011 Climate change and the potential global distribution of serrated tussock (*Nassella trichotoma*). *Weed Sci.* 59, 538–545. doi: [10.1614/WS-D-11-00032.1](https://doi.org/10.1614/WS-D-11-00032.1)
- Werner C, Zumkier U, Beyschlag W, Mágus C 2010 High competitiveness of a resource demanding invasive *Acacia* under low resource supply. *Plant Ecol.* 206, 83–96. doi: [10.1007/s11258-009-9625-0](https://doi.org/10.1007/s11258-009-9625-0)
- Wernli H, Schwierz C 2006 Surface Cyclones in the ERA-40 Dataset (1958–2001). Part I: Novel Identification Method and Global Climatology. *J. Atmos. Sci.* 63, 2486–2507. doi: [10.1175/jas3766.1](https://doi.org/10.1175/jas3766.1)
- Wibig J 1999 Precipitation in Europe in relation to circulation patterns at the 500 hPa level. *Int. J. Climatol.* 19, 253–269. doi: [10.1002/\(SICI\)1097-0088\(19990315\)19:3<253::AID-JOC366>3.0.CO;2-0](https://doi.org/10.1002/(SICI)1097-0088(19990315)19:3<253::AID-JOC366>3.0.CO;2-0)
- Wild M 2009 Global dimming and brightening: A review. *JGR Atmos.* 114, D00D16. doi: [10.1029/2008JD011470](https://doi.org/10.1029/2008JD011470)
- Wild M 2012 Enlightening Global Dimming and Brightening. *Bull. Am. Meteorol. Soc.* 93, 27–37. doi: [10.1175/BAMS-D-11-00074.1](https://doi.org/10.1175/BAMS-D-11-00074.1)
- Wild RJ, Dubé WP, Aikin KC, Eilerman SJ, Neuman JA et al. 2017 On-road measurements of vehicle NO₂/NO_x emission ratios in Denver, Colorado, USA. *Atmos. Environ.* 148, 182–189. doi: [10.1016/J.ATMOSENV.2016.10.039](https://doi.org/10.1016/J.ATMOSENV.2016.10.039)
- Wilks DS 2016 “The stippling shows statistically significant grid points”: How research results are routinely overstated and overinterpreted, and what to do about it. *Bull. Am. Meteorol. Soc.* 97, 2263–2273. doi: [10.1175/BAMS-D-15-00267.1](https://doi.org/10.1175/BAMS-D-15-00267.1)
- Witkowski F, Andral B, Derolez V, Tomasino C 2017 Campagne de surveillance DCE 2015 en Méditerranée française Districts « Rhône et Cotiers Méditerranéens » et « Corse ».
- WMO 2017 WMO Guidelines on the Calculation of Climate Normals - 2017 edition. Geneva, Switzerland.
- Wood S, Sebastian K, Scherr SJ 2000 Pilot analysis of global ecosystems: Agroecosystems.
- Woollings T, Blackburn M 2012 The North Atlantic Jet Stream under Climate Change and Its Relation to the NAO and EA Patterns. *J. Clim.* 25, 886–902. doi: [10.1175/jcli-d-11-00087.1](https://doi.org/10.1175/jcli-d-11-00087.1)
- Wöppelmann G, Marcos M 2012 Coastal sea level rise in Southern Europe and the nonclimate contribution of vertical land motion. *JGR Ocean.* 117. doi: [10.1029/2011JC007469](https://doi.org/10.1029/2011JC007469)
- World Economic Forum 2011 Scenarios for the Mediterranean Region. 40.
- Wu J, Zha J, Zhao D, Yang Q 2018a Changes in terrestrial near-surface wind speed and their possible causes: an overview. *Clim. Dyn.* 51, 2039–2078. doi: [10.1007/s00382-017-3997-y](https://doi.org/10.1007/s00382-017-3997-y)
- Wu N, Wang C, Ausseil AG, Alhafedh Y, Broadhurst L et al. 2018b Direct and indirect drivers of change in biodiversity and nature's contributions to people, in *IPBES (2018): The IPBES regional assessment report on biodiversity and ecosystem services for Asia and the Pacific*, eds. Karki M, Senaratna Sellamuttu S, Okayasu S, Suzuki W [Bonn, Germany: Secretariat of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services], 265–370.
- Wurl O, Obbard JP 2004 A review of pollutants in the sea-surface microlayer (SML): a unique habitat for marine organisms. *Mar. Pollut. Bull.* 48, 1016–1030. doi: [10.1016/j.marpolbul.2004.03.016](https://doi.org/10.1016/j.marpolbul.2004.03.016)
- Xoplaki E, González-Rouco JF, Gyalistras D, Luterbacher J, Rickli R et al. 2003a Interannual summer air temperature variability over Greece and its connection to the large-scale atmospheric circulation and Mediterranean SSTs 1950–1999. *Clim. Dyn.* 20, 537–554. doi: [10.1007/s00382-002-0291-3](https://doi.org/10.1007/s00382-002-0291-3)
- Xoplaki E, Gonzalez-Rouco JF, Luterbacher J, Wanner H 2004 Wet season Mediterranean precipitation variability: influence of large-scale dynamics and trends. *Clim. Dyn.* 23,

- 63–78. doi: [10.1007/s00382-004-0422-0](https://doi.org/10.1007/s00382-004-0422-0)
- Xoplaki E, González-Rouco JF, Luterbacher J, Wanner H 2003b Mediterranean summer air temperature variability and its connection to the large-scale atmospheric circulation and SSTs. *Clim. Dyn.* 20, 723–739.
doi: [10.1007/s00382-003-0304-x](https://doi.org/10.1007/s00382-003-0304-x)
- Yeruham E, Rilov G, Shpigel M, Abelson A 2015 Collapse of the echinoid *Paracentrotus lividus* populations in the Eastern Mediterranean - result of climate change? *Sci. Rep.* 5, 13479. doi: [10.1038/srep13479](https://doi.org/10.1038/srep13479)
- Yin JH 2005 A consistent poleward shift of the storm tracks in simulations of 21st century climate. *Geophys. Res. Lett.* 32, {L18}701. doi: [10.1029/2005GL023684](https://doi.org/10.1029/2005GL023684)
- Youssef L, Younes G, Kouzayha A, Jaber F 2015 Occurrence and levels of pesticides in South Lebanon water. *Chem. Speciat. Bioavailab.* 27, 62–70. doi: [10.1080/09542299.2015.1023092](https://doi.org/10.1080/09542299.2015.1023092)
- Yurtkuran Z, Saygi Y 2013 Assessment of Pesticide Residues in Karabogaz Lake from Kizilirmak Delta, Turkey. *Bull. Environ. Contam. Toxicol.* 91, 165–170.
doi: [10.1007/s00128-013-1037-0](https://doi.org/10.1007/s00128-013-1037-0)
- Zaghdene H, Kallel M, Louati A, Elleuch B, Oudot J et al. 2005 Hydrocarbons in surface sediments from the Sfax coastal zone, (Tunisia) Mediterranean Sea. *Mar. Pollut. Bull.* 11, 1287–1294. doi: [10.1016/j.marpolbul.2005.04.045](https://doi.org/10.1016/j.marpolbul.2005.04.045)
- Zahn A, Rainho A, Rodrigues L, Palmeirim JM 2009 Low macro-arthropod abundance in exotic Eucalyptus plantations in the Mediterranean. *Appl. Ecol. Environ. Res.* 7, 297–301.
- Zalidis GC, Stamatiadis S, Takavagoglou V, Eskridge K, Misopolinos N 2002 Impacts of agricultural practices on soil and water quality in the Mediterranean region and proposed assessment methodology. *Agric. Ecosyst. Environ.* 88, 137–146.
- Zampieri M, Lionello P 2011 Anthropic land use causes summer cooling in Central Europe. *Clim. Res.* 46, 255–268.
doi: [10.3354/cr00981](https://doi.org/10.3354/cr00981)
- Zanis P, Hadjinicolaou P, Pozzer A, Tyrllis E, Dafka S et al. 2014 Summertime free-tropospheric ozone pool over the eastern Mediterranean/Middle East. *Atmos. Chem. Phys.* 14, 115–132. doi: [10.5194/acp-14-115-2014](https://doi.org/10.5194/acp-14-115-2014)
- Zappa G, Hoskins BJ, Shepherd TG 2015 The dependence of wintertime Mediterranean precipitation on the atmospheric circulation response to climate change. *Environ. Res. Lett.* 10, 104012.
- Zappa G, Shaffrey LC, Hodges KI, Sansom PG, Stephenson DB et al. 2013 A Multimodel Assessment of Future Projections of North Atlantic and European Extratropical Cyclones in the CMIP5 Climate Models. *J. Clim.* 26, 5846–5862.
doi: [10.1175/JCLI-D-12-00573.1](https://doi.org/10.1175/JCLI-D-12-00573.1)
- Zemp M, Paul F, Hoelzle M, Haeberli W 2008 Glacier fluctuations in the European Alps, 1850–2000, in *Darkening Peaks: Glacier Retreat, Science, and Society*, eds. Orlove B, Wiegandt E, Luckman BH (Berkeley, Los Angeles, London: University of California Press), 152–167.
- Zenetos A 2019 Mediterranean Sea: 30 Years of Biological Invasions (1988–2017). in *Proceedings of the 1st Mediterranean Symposium on the Non-Indigenous Species* (Tunis, Antalya: SPA/RAC Publications), 13–19.
- Zenetos A, Çınar ME, Crocetta F, Golani D, Rosso A et al. 2017 Uncertainties and validation of alien species catalogues: The Mediterranean as an example. *Estuar. Coast. Shelf Sci.* 191, 171–187. doi: [10.1016/j.ecss.2017.03.031](https://doi.org/10.1016/j.ecss.2017.03.031)
- Zenetos A, Mačić V, Jaklin A, Lipej L, Poursanidis D et al. 2016 Adriatic ‘opisthobranchs’ (Gastropoda, Heterobranchia): shedding light on biodiversity issues. *Mar. Ecol.* 37, 1239–1255. doi: [10.1111/maec.12306](https://doi.org/10.1111/maec.12306)
- Zerefos CS, Eleftheratos K, Meleti C, Kazadzis S, Romanou A et al. 2009 Solar dimming and brightening over Thessaloniki, Greece, and Beijing, China. *Tellus B Chem. Phys. Meteorol.* 61, 657–665. doi: [10.1111/j.1600-0889.2009.00425.x](https://doi.org/10.1111/j.1600-0889.2009.00425.x)
- Zhou JL, Maskooui K 2003 Distribution of polycyclic aromatic hydrocarbons in water and surface sediments from Daya Bay, China. *Environ. Pollut.* 121, 269–281.
doi: [10.1016/S0269-7491\(02\)00215-4](https://doi.org/10.1016/S0269-7491(02)00215-4)
- Zittis G 2018 Observed rainfall trends and precipitation uncertainty in the vicinity of the Mediterranean, Middle East and North Africa. *Theor. Appl. Climatol.* 134, 1207–1230.
doi: [10.1007/s00704-017-2333-0](https://doi.org/10.1007/s00704-017-2333-0)
- Zittis G, Hadjinicolaou P 2017 The effect of radiation parameterization schemes on surface temperature in regional climate simulations over the MENA-CORDEX domain. *Int. J. Climatol.* 37, 3847–3862. doi: [10.1002/joc.4959](https://doi.org/10.1002/joc.4959)
- Zittis G, Hadjinicolaou P, Fnais M, Lelieveld J 2016 Projected changes in heat wave characteristics in the eastern Mediterranean and the Middle East. *Reg. Environ. Chang.* 16, 1863–1876. doi: [10.1007/s10113-014-0753-2](https://doi.org/10.1007/s10113-014-0753-2)
- Zittis G, Hadjinicolaou P, Klangidou M, Proestos Y, Lelieveld J 2019 A multi-model, multi-scenario, and multi-domain analysis of regional climate projections for the Mediterranean. *Reg. Environ. Chang.* 19, 2621–2635.
doi: [10.1007/s10113-019-01565-w](https://doi.org/10.1007/s10113-019-01565-w)
- Zittis G, Hadjinicolaou P, Lelieveld J 2014 Role of soil moisture in the amplification of climate warming in the eastern Mediterranean and the Middle East. *Clim. Res.* 59, 27–37.
doi: [10.3354/cr01205](https://doi.org/10.3354/cr01205)
- Zubler EM, Folini D, Lohmann U, Lüthi D, Schär C et al. 2011 Simulation of dimming and brightening in Europe from 1958 to 2001 using a regional climate model. *JGR Atmos.* 116, D18205. doi: [10.1029/2010JD015396](https://doi.org/10.1029/2010JD015396)
- Zurek MB, Henrichs T 2007 Linking scenarios across geographical scales in international environmental assessments. *Technol. Forecast. Soc. Change* 74, 1282–1295.
doi: [10.1016/j.techfore.2006.11.005](https://doi.org/10.1016/j.techfore.2006.11.005)

Information about authors

Coordinating Lead Authors

Semia Cherif:

Tunis El Manar University, Higher Institute for Applied Biological Sciences of Tunis (UTM-ISSBAT), Tunis, Tunisia

Enrique Doblas-Miranda:

CREAF, Bellaterra (Barcelona), Spain & Universitat Autònoma de Barcelona, Bellaterra (Barcelona), Spain

Piero Lionello:

University of Salento, Lecce, Italy

Lead Authors

Carlos Borrego:

Department of Environment and Planning (DAO), Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, Portugal

Filippo Giorgi:

Abdus Salam International Centre for Theoretical Physics (ICTP), Trieste, Italy

Ana Iglesias:

Universidad Politécnica de Madrid (UPM), Madrid, Spain

Sihem Jebari:

Tunisian National Research Institute for Rural Engineering, Water and Forestry (INRGREF), Tunis, Tunisia

Ezzeddine Mahmoudi:

University of Carthage, Bizerte, Tunisia

Marco Moriondo:

Institute of BioEconomy, Italian National Research Council (CNR-IBE), Florence, Italy

Olivier Pringault:

Mediterranean Institute of Oceanography (MIO), Institute of Research for Development (IRD), Marseille, France

Gil Rilov:

National Institute of Oceanography, Israel Oceanographic and Limnological Research (IOLR), Haifa, Israel

Samuel Somot:

Centre National de Recherches Météorologiques (CNRM, Université de Toulouse, Météo-France, CNRS), Toulouse, France

Athanassios Tsikliras:

Aristotle University of Thessaloniki, Thessaloniki, Greece

Monsterrat Vila:

Spanish National Research Council, Doñana Biological Station (EBD-CSIC), Sevilla, Spain

Georgios Zittis:

The Cyprus Institute, Nicosia, Cyprus

Contributing Authors

Giovanni Argenti:

University of Florence, Florence, Italy

Marie-Anne Auger-Rozenberg:

Orléans Forest Zoology research unit, French National Research Institute for Agriculture, Food and Environment (INRAE), Orléans, France

Ernesto Azzurro:

Institute for Marine Biological Resources and Biotech-

nology of the National Research Council (CNR-IRBIM), Ancona, Italy

Corina Basnou:

CREAF, Bellaterra (Barcelona), Spain

Sophie Bastin:

Laboratoire ATmosphères Milieux Observations Spatiales/Institut Pierre Simon Laplace (LATMOS/IPSL), UVSQ Université Paris-Saclay, SU, CNRS, Guyancourt, France

Mustapha Béjaoui:

University of Carthage, Bizerte, Tunisia

Lorenzo Brilli:

Institute of BioEconomy, Italian National Research Council (CNR-IBE), Florence, Italy

Martina Carrete:

University Pablo de Olavide, Sevilla, Spain

Emma Cebria:

Blanes Centre for Advanced Studies (CEAB), Spanish National Research Council (CSIC), Blanes, Girona, Spain

Hanene Chaabane :

Laboratory of Pest control and Integrated Protection in Agriculture / LMI NAILA, National Institute of Agronomy of Tunisia (INAT), University of Carthage, Tunis, Tunisia

Sílvia Coelho:

Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, Portugal

Renato Colucci:

Institute of Polar Sciences, Marine Science Institute, National Research Council (CNR-ISPI), Trieste, Italy

Styliani Dafka:

Justus-Liebig University, Giessen, Germany

Sofia Darmaraki:

Dalhousie University, Canada

Camilla Dibari:

University of Florence, Florence, Italy

Donna Dimarchopoulou:

Aristotle University of Thessaloniki, Thessaloniki, Greece

Jean-Claude Dutay:

Laboratory of Climate and Environmental Sciences (LSCE), Pierre Simon Laplace Institute (IPSL), Gif-sur-Yvette, France

Monia El Bour:

National Institute of Marine Sciences and Technology (INSTMI), Carthage - Salammbô, Tunisia

Antonietta Elia:

University of Santiago de Compostela, Santiago de Compostela, Spain

Elena Georgopoulou:

National Observatory of Athens, Athens, Greece

Sylvaine Giakoumi:

Zoological Station Anton Dohrn, Naples, Italy & School of Biological Sciences, University of Queensland, Australia

Juan Jesus Gonzalez Aleman:

Complutense University of Madrid, Madrid, Spain

Pablo González-Moreno:

CABI, Egham, United Kingdom

Madeleine Goutx:

Mediterranean Institute of Oceanography (MIO), CNRS, Marseille, France

CHAPTER 2 - DRIVERS OF CHANGE

- Olivier Grunberger:
National Agricultural Institute of Tunisia (INAT), French National Research Institute for Sustainable Development (IRD), Tunis, Tunisia
- Ivan Gütler:
Croatian Meteorological and Hydrological Service, Zagreb, Croatia
- Nathalie Hilmi:
Monaco Scientific Centre (MSC), Monaco
- Gabriel Jorda:
Spanish Institute of Oceanography, Balearic Islands Oceanographic Centre (COB-IEO), Palma, Spain
- Stelios Katsanevakis:
University of the Aegean, Mitilini, Greece
- Mehdi Lahlou:
National Institute of Statistics and Applied Economics (INSEA), Rabat, Morocco
- Manfred A. Lange:
The Cyprus Institute, Nicosia, Cyprus
- Luisa Leolini:
University of Florence, Florence, Italy
- Myriam Lopes:
Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, Portugal
- Annarita Mariotti:
National Oceanic and Atmospheric Administration (NOAA), Silver Spring, United States of America
- Ana Isabel Miranda:
Centre for Environmental and Marine Studies (CESAM), Department of Environment and Planning, University of Aveiro, Aveiro, Portugal
- Meryem Mojtabah:
University of Angers, Angers, France
- Alexandra Monteiro:
Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, Portugal
- Samuel Morin:
Université Grenoble Alpes, Université de Toulouse, Météo-France, CNRS, CNRM, Centre d'Études de la Neige, Grenoble, France
- Pierre Nabat:
Centre National de Recherches Météorologiques (CNRM, Université de Toulouse, Météo-France, CNRS), Toulouse, France
- Anika Obermann-Hellhund:
Goethe University, Frankfurt am Main, Germany
- Tuğba ÖzTÜRK:
İşık University, Istanbul, Turkey
- Androniki Pardalou:
Aristotle University of Thessaloniki, Thessaloniki, Greece
- Sandra Rafael:
Centre for Environmental and Marine Studies (CESAM), University of Aveiro, Aveiro, Portugal
- Francesca Raffaele:
Abdus Salam International Centre for Theoretical Physics (ICTP), Trieste, Italy
- Lena Reimann:
Christian-Albrechts University, Kiel, Germany
- Alain Roques:
Orléans Forest Zoology research unit, French National Research Institute for Agriculture, Food and Environment (INRAE), Orléans, France
- Asma Sakka Hlaili:
University of Carthage, Bizerte, Tunisia
- Alberto Santini:
Institute for Sustainable Plant Protection, Italian National Research Council (CNR), Sesto Fiorentino, Italy
- Giuseppe Scarcella:
National Research Council, Institute of Marine Biological Resources and Biotechnologies (CNR-IRBIM), Ancona, Italy
- Katrin Schroeder:
Italian National Research Council, Institute of Marine Science (CNR-ISMAR), Venice, Italy
- Isla Simpson:
Climate and Global Dynamics Laboratory of the National Center for Atmospheric Research in Boulder, Colorado, United States of America
- Nicolina Staglianò:
University of Florence, Florence, Italy
- Meryem Tanarhte:
Hassan II Mohammedia University, Casablanca, Morocco
- Rob Tanner:
European and Mediterranean Plant Protection Organisation (EPPO), Paris, France
- Rémi Thiéblemont:
French Geological Survey, Orléans, France
- Yves Tramblay:
HydroSciences Montpellier (University Montpellier, CNRS, IRD), Montpellier, France
- Marco Turco:
University of Murcia, Murcia, Spain
- Athanasiос Vafeidis:
Christian-Albrechts University, Kiel, Germany
- Martin Wild:
Institute of Atmospheric and Climate Science, ETH Zürich, Zürich, Switzerland
- Elena Xoplaki:
Department of Geography & Center for international Development and Environmental Research, Justus Liebig University, Giessen, Germany
- Argyro Zenetos:
Institute of Marine Biological Resources and Inland Waters, Hellenic Centre for Marine Research (HCMR), Argiroupoli, Greece